Companion Book For The Pathology Prof Buster Course On medLIVE By Dr. Priyanka

Companion Book For The Pathology Prof Buster Course On medLIVE By Dr. Priyanka

medLIVE

PATHOLOGY
PROF BUSTER

PATHOLOGY DEMYSTIFIED
DISTINCTION AMPLIFIED

Volume-1

PRIYANKA SACHDEV, MD

Ed Uthman

PAPER 1

GENERAL PATHOLOGY

Chapter 1 | Cell Adaptations, Cell Injury, Cell Death

(1.01) Cellular adaptations
(1.02) Mechanisms of Cell Injury
(1.03) Morphological changes – Reversible versus irreversible cell injury
(1.04) Apoptosis and Efferocytosis
(1.05) Necrosis
(1.06) Necrotosis, Pyroptosis, Ferroptosis
(1.07) Gangrene
(1.08) Pathological calcification
(1.09) Pigments
(1.10) Cellular Ageing

Chapter 2 | Inflammation

(2.01) Vascular events of Acute inflammation
(2.02) Cellular events of Acute inflammation
(2.03) Chronic inflammation or Granulomatous inflammation
(2.04) Chemical mediators of Inflammation

Chapter 3 | Wound Healing and Repair

(3.01) Wound healing (Regeneration and Repair)
(3.02) Healing of skin wounds (Primary and Secondary intention)
(3.03) Factors affecting wound healing and Complications of Wound Healing

Chapter 4 | Hemodynamic Disorders

(4.01) Edema and Effusion
(4.02) Hyperemia and Congestion (CVC lung, CVC Liver, CVC Spleen)
(4.03) Thrombosis
(4.04) Embolism
(4.05) Ischemia
(4.06) Infarction
(4.07) Shock

Chapter 5 | Genetic Disorders

(5.01) Genetic inheritance of Single gene disorders (AD, AR, XD, XR)
(5.02) Cytogenetic Abnormalities

Chapter 6 | Immunopathology

(6.01) Hypersensitivity reactions
(6.02) Transplant rejection
(6.03) Autoimmunity
(6.04) Systemic Lupus Erythematosus (SLE)
(6.05) HIV / AIDS
(6.06) Amyloidosis

Chapter 7 | Neoplasia

(7.01) Neoplasia (Definition, Nomenclature and Predisposing conditions)
(7.02) Teratoma, Hamartoma and Choristoma
(7.03) Benign versus Malignant tumours

PAPER 1

(7.04) Spread of tumours (Metastasis)
(7.05) Proto-oncogenes – RAS, ABL- BCR
(7.06) Tumour suppressor genes - RB, P53
(7.07) Carcinogenesis (PHYSICAL, CHEMICAL, Biological)
(7.08) Clinical Features (Tumour Lysis Syndrome and Paraneoplastic syndromes)
(7.09) Grading and staging of cancers
(7.10) Laboratory diagnosis of cancer (Tumour markers)

Chapter 8 | Environmental and Nutritional Disorders

(8.01) Obesity

Chapter 9 | Cancers of Infancy and Childhood

(9.01) Cancers of infancy and childhood

Hematology

Chapter 10 | RBC Disorders

(10.01) Bone marrow aspiration and biopsy
(10.02) Reticulocytes
(10.03) RBC indices and ESR
(10.04) Anemia→ Definition, classification, c/f
(10.05) Hemolytic anemias → Definition, classification, lab diagnosis
(10.06) Hereditary spherocytosis
(10.07) G6PD deficiency Anemia
(10.08) Paroxysmal Noctural Hematuria (PNH)
(10.09) Autoimmune Hemolytic Anemia (AIHA)
(10.10) Sickle cell Anemia
(10.11) Thalassemia
(10.12) Iron deficiency Anemia (IDA)
(10.13) Sideroblastic Anemia
(10.14) Anemia of Chronic disease (AOCD)
(10.15) Megaloblastic Anemia
(10.16) Pernecious Anemia
(10.17) Aplastic Anemia and Pure Red cell Aplasia (PRCA)
(10.18) Pancytopenia

Chapter 11 | WBC Disorders

(11.01) Interpretation of DLC
(11.02) Leukemoid Reaction
(11.03) Leukemias and Lymphomas (Definition and Classification)
(11.04) Chronic Myeloid Leukemia (CML)
(11.05) Acute Myeloid Leukemia (AML)
(11.06) Acute Lymphoblastic Leukemia (ALL)
(11.07) Chronic Lymphocytic Leukemia (CLL)
(11.08) Polycythemia Vera
(11.09) Myelodysplastic syndrome
(11.10) Hodgkin's Lymphoma (HL)
(11.11) Non-Hodgkin's Lymphoma (NHL)
(11.12) Plasma cell disorders → Multiple Myeloma
(11.13) Spleenomegaly

PAPER 1

Chapter 12 | Platelet Disorders

(12.01) Bleeding disorders (Definition and Classification)
(12.02) Thrombocytopenia (ITP, TTP, HUS)
(12.03) Functional platelet disorders (Bernard Soulier syndrome, Glanzmann's syndrome)
(12.04) Coagulation Disorders- Hemophilia and Von Willebrand's disease
(12.04) Disseminated Intravascular Coagulation (DIC)

PAPER 2

SYSTEMIC PATHOLOGY

Chapter 13 | Vascular System

(13.01) Atherosclerosis
(13.02) Aneurysms & Aortic Dissection
(13.03) Hypertensive Vascular Disease
(13.04) Vasculitis
(13.05) Raynaud's phenomenon
(13.06) Tumours of Blood Vessels

Chapter 14 | Heart

(14.01) congenital heart diseases
(14.02) ischemic heart diseases / coronary artery disease
(14.03) Rheumatic fever and Rheumatic heart disease
(14.04) Infective Endocarditis
(14.05) NON-INFECTIVE Endocarditis (NBTE and Libman-sacks Endocarditis)
(14.06) Cardiomyopathies
(14.07) Pericarditis
(14.08) Cor Pulmonale

Chapter 15 | Respiratory System

(15.01) Normal anatomy and histology
(15.02) Acute respiratory distress syndrome (ARDS)
(15.03) COPD – Chronic Bronchitis
(15.04) COPD – Emphysema
(15.05) COPD – Asthma
(15.06) COPD – Bronchiectesis
(15.07) Pneumoconiosis → Asbestosis, Silicosis, Bagassosis
(15.08) Pneumonias
(15.09) Pulmonary Tuberculosis
(15.10) Lung abscess
(15.11) Lung carcinoma

Chapter 16 | Gastrointestinal System

(16.01) Barret esophagus
(16.02) Achalasia (Cardiospasm)

PAPER 2

(16.03) Esophageal lacerations
(16.04) Hiatus hernia
(16.05) Esophageal tumours
(16.06) Pyloric stenosis
(16.07) Gastritis
(16.08) Peptic ulcer
(16.09) Gastric carcinoma
(16.10) Meckel's diverticulum
(16.11) Intussusception
(16.12) Inflammatory Bowel Disease (Crohn's disease, Ulcerative colitis)
(16.13) Infectious Enterocolitis (Typhoid, TB, Amoebic ulcer)
(16.14) Malabsorption syndrome (Celiac sprue, Topical sprue, Whipple's disease)
(16.15) Carcinoid tumours
(16.16) Appendicitis
(16.17) Hirschsprung's disease
(16.18) Colorectal carcinoma

Chapter 17 | Hepatobiliary and Pancreatic System

(17.01) Jaundice
(17.02) Neonatal Jaundice / Congenital hyperbilirubinemia
(17.03) Viral hepatitis
(17.04) Cirrhosis (DEFINITION, PATHOGENESIS, CLASSIFICATION)
(17.05) Alcoholic Liver Disease (Alcoholic Cirrhosis)
(17.06) Non-Alcoholic Fatty liver Disease (NAFLD)
(17.07) Wilson Disease
(17.08) Hemochromatosis
(17.09) Portal Hypertension
(17.10) CVC Liver / Nutmeg liver
(17.11) Liver abcess (Pyogenic, Amoebic, Hydatid cyst)
(17.12) Malignant tumours of Liver
(17.13) Cholelithiasis (Gall stones)
(17.14) Cholecystitis
(17.15) Gall bladder carcinoma
(17.16) Pancreatitis
(17.17) Pancreatic tumours

Chapter 18 | Urinary System

(18.01) Normal anatomy and histology
(18.02) Nephritic versus Nephrotic syndrome
(18.03) Glomerulonephritis Pathogenesis and Classification
(18.04) Primary Glomerulonephritis (APGN, RPGN, MCD, MGN, MPGN, FSGN, FSGS, IgA nephropathy, Chronic GN)
(18.05) Diabetic Nephropathy
(18.06) Pyelonephritis
(18.07) Renal calculus and hydronephrosis
(18.08) Renal changes in Hypertension (Nephrosclerosis)
(18.09) Polycystic kidney disease
(18.10) Tumours of kidney
(18.11) Bladder carcinoma

Chapter 19 | Male Genital System

(19.01) Cryptorchidism
(19.02) Testicular tumours
(19.03) Benign Prostate Hyperplasia & Prostate carcinoma

PAPER 2

(19.04) Male infertility

Chapter 20 | Female Genital System

(20.01) Cervical Precancers (CIN / SIL)
(20.02) Cervix cancer
(20.03) Endometriosis
(20.04) Adenomyosis
(20.05) Endometrial hyperplasia
(20.06) Endometrial carcinoma
(20.07) Tumours of Myometrium → Leiomyoma
(20.08) Ovarian tumours
(20.09) Gestational trophoblastic Disease

Chapter 21 | Breast

(21.01) Stromal Breast Tumors (Fibrosdenoma and Phyllodes)
(21.02) Breast Carcinoma

Chapter 22 | Endocrine System

(22.01) Autoimmune disorders of thyroid (Hashimoto thyroiditis, Graves' disease)
(22.02) Goiter
(22.03) Thyroid neoplasm
(22.04) Diabetes mellitus
(22.05) Zollinger – Ellison syndrome (Gastrinoma)
(22.06) Cushing syndrome
(22.07) Pheochromocytoma
(22.08) Multiple Endocrine Neoplasia (MEN) syndromes

Chapter 23 | Muscoloskeletal System

(23.01) Osteomyelitis (Pyogenic, Tubercular)
(23.02) Bone tumours
(23.03) GOUT

Chapter 24 | Central Nervous System

(24.01) Meningitis
(24.02) Tumours of CNS

Chapter 25 | Salivary Glands

(25.01) Tumors of Salivary Glands

Chapter 26 | Skin

(26.01) Tumours of Skin

(1.01) CELLULAR ADAPTATIONS

University Exams

Long questions

1. Define Cellular adaptations. Write its types with mechanisms and examples

Short questions

1. Definition, types, mechanism and examples of hypertrophy.
2. Definition, types, mechanism and examples of Atrophy.
3. Definition, types, mechanism and examples of Metaplasia.
4. Definition, types, mechanism and examples of Dysplasia.
5. Definition, types, mechanism and examples of Hyperplasia
6. Differentiate Hypertrophy and hyperplasia.
7. Differences between atrophy, hypertrophy and hyperplasia.
8. Differences between hypertrophy, hyperplasia and metaplasia.
9. Differentiate metaplasia and Dysplasia.

Very short questions

1. Define hypertrophy.
2. Define Atrophy.
3. Define Metaplasia.
4. Define Dysplasia.
5. Define Hyperplasia.
6. Write two examples of hypertrophy.
7. Write two examples of Atrophy.
8. Write two examples of Metaplasia.
9. Write two examples of Dysplasia.
10. Write two examples of Hyperplasia.

Overview

- Introduction
- Definition
- Types →

    1. Atrophy
    2. Hypertrophy
    3. Hyperplasia
    4. Metaplasia
    5. Dysplasia

Introduction



Definition

Adaptations →
- Reversible
- Structural responses to physiologic stress (e.g., pregnancy) and pathologic stress
- During which new but altered steady states are achieved
- Allowing cell to survive and continue to function

Types

1. Atrophy → Decrease in size of cells / shrinkage of cells
2. Hypertrophy→ Increase in size of cells
3. Hyperplasia → Increase in number of cells
4. Metaplasia→ Transformation of one adult cell type with another
5. Dysplasia → Disordered cellular development



(3) Hyperplasia

Definition →

Increase in number of cells, not size of cell



Mechanism

- Increased production of growth factors, growth factor receptors



Types

A. Physiologic hyperplasia
B. Pathologic hyperplasia

(A) Physiologic hyperplasia

1. Hyperplasia of female breast at puberty, pregnancy and lactation
2. Enlarged size of the uterus in pregnancy → physiologic hypertrophy as well as hyperplasia.
3. Compensatory hyperplasia →
4. Following nephrectomy on one side → hyperplasia of nephrons of other kidney











(B) Pathologic hyperplasia

1. Endometrial hyperplasia → following oestrogen excess
2. Benign prostatic hyperplasia → due to DHT (dihydrotestosterone)
3. Skin warts → from hyperplasia of epidermis due to HPV





(2) Hypertrophy

Definition →

Increase in size of cells, not number of cells



Mechanism

Increased size is due to more structural components / cellular proteins (not due to cellular swelling)



Types

(A) Physiologic hypertrophy
(B) Pathologic hypertrophy

(A) Physiologic hypertrophy

1. Enlarged size of uterus in pregnancy → physiologic hypertrophy as well as hyperplasia.
2. Hypertrophy of skeletal muscle → in athletes and manual labourers.



(B) Pathologic hypertrophy

1. Hypertrophy of cardiac muscle→ occurs in Systemic hypertension, Aortic valve disease
2. Hypertrophy of smooth muscle→ e.g. Cardiac achalasia (in oesophagus), Pyloric stenosis (in stomach).





(1) Atrophy

Definition →

Decrease in cell size and number of cells

- Atrophied cells are smaller than normal but they are still viable



Mechanism

Shrinkage in cell size is due to reduction in cell organelles

Types

(A) Physiologic atrophy
(B) Pathologic atrophy

(A) Physiologic atrophy

1. Atrophy of thymus in adult life.
2. Atrophy of ovary after menopause.
3. Atrophy of brain with ageing.
4. Atrophy of uterus after parturition





(B) Pathologic atrophy

1. Starvation atrophy → cachexia seen in cancer and severely ill patients.
2. Ischaemic atrophy → atrophic kidney in atherosclerosis of renal artery, Atrophy of brain in cerebral atherosclerosis.
3. Neuropathic atrophy → e.g. Poliomyelitis
4. Disuse atrophy → e.g. Wasting of muscles of limb immobilized in cast, Atrophy of the pancreas in obstruction of pancreatic duct.



5. Endocrine atrophy → eg. Hypopituitarism may lead to atrophy of thyroid, adrenal and gonads
6. Pressure atrophy → eg. Erosion of skull by meningioma, Erosion of the sternum by aneurysm of arch of aorta
7. Idiopathic atrophy → no cause present





(4) Metaplasia

Definition →

One mature/adult cell type is replaced by another mature/adult cell type



Mechanism

- Metaplasia does not result from a change in phenotype of an already differentiated cell type
- Result of a reprogramming of stem cells that are known to exist in normal tissues.



Types

(A) Squamous (Col → Sq)
(B) Columnar (Sq → Col)

(A) Squamous metaplasia Examples

1. In bronchus in chronic smokers.
2. In gallbladder in cholelithiasis.
3. In prostate in chronic prostatitis
4. In uterine endocervix in prolapse of uterus



(B) Columnar metaplasia Examples

1. Barrett's oesophagus → due to GERD
2. Cervical erosion → Ectocervix → due to increased exposure of cervical epithelium to estrogen





(5) Dysplasia

Definition →

- Disordered cellular development
- Characterised by cellular cytologic changes

Mechanism

Cytological changes →
1. Increased number of layers
2. Disorderly arrangement of cells from basal layer to surface layer
3. Loss of basal polarity i.e. nuclei lying away from basement membrane
4. Cellular and nuclear pleomorphism
5. Increased nucleocytoplasmic (N/C) ratio
6. Nuclear hyperchromatism
7. Increased mitotic activity



NORMAL ENDOCERVICAL EPITHELIUM

- No. of layers normal
- Ordered arrangement
- Basal Polarity ⊕
- Pleomorphism ⊖
- $\frac{N}{C}$ Normal
- Hyper chromatism ⊖
- Mitosis normal



DYSPLASIA

- No. of layers ↑
- Disordered arrangement
- Basal Polarity ⊖
- Pleomorphism ⊕
- $\frac{N}{C}$ ↑
- Hyper chromatism ⊕
- Mitosis ↑

- Full thickness dysplasia is known as carcinoma in situ (Precurssor of cancer)



NORMAL ENDOCERVICAL EPITHELIUM | SQUAMOUS METAPLASIA | DYSPLASIA | CARCINOMA IN SITU

	Hyperplasia	Hypertrophy	Atrophy	Metaplasia	Dysplasia
Definition	↑ Number	↑ Size	↓ Number ↓ Size	Transformation one cell → Another	Disordered development
Diagram					
Mechanism	↑ In GR and GF	↑ In cell organelles	↓ In cell organelles	Reprogramming of Stem Cells	- No. of layers ↑ - Disordered arrangement - Basal Polarity ⊖ - Pleomorphism ⊕ - $\frac{N}{C}$ ↑ - Hyper chromatism ⊕ - Mitosis ↑
Types with e.g.	Physio Patho	Physio Patho	Physio Patho	Epi → Sq Mesen → Col	

(1.02) MECHANISMS OF CELL INJURY

University Exams

Long questions

1. Describe biochemical mechanisms of cell injury
2. What are free radicals? Describe free radical-induced cell injury.

Short questions

1. Role of free radicals in cell injury
2. Role of ATP depletion in cell injury

Very short questions

1. Types of free radicals
2. Scavangers of free radicals
3. Antioxidants

Overview

4 mechanisms →

1. ATP depletion
2. Mitochondrial Damage
3. Loss of calcium homeostasis
4. Free radical injury →

- Introduction
- Generation
- Types
- Effects
- Removal

(1) ATP depletion



Ischemic
↓
Mitochondria - reduced oxidative phosphorylation
↓
ATP depletion

ATP depletion
↓
Cell membrane - reduced sodium pump
↓
Sodium, water enter the cell and potassium exits
↓
Endoplasmic reticulum dilates
Cell swells Blebs appear



ATP depletion
↓
Anaerobic glycolysis occurs
↓
Loss of glycogen
↓
Accumulation of lactic acid
↓
Acid pH
↓
Clumping of nuclear chromatin

ATP depletion
↓
RER loses ribosomes
↓
Protein synthesis falls
↓
Structural proteins (membranes, cytoskeleton) and enzymes depletes



(2) Mitochondrial Damage

- Mitochondria supply life-sustaining energy by producing ATP (Power house of cell)

Mitochondrial damage
↓
Formation of a high-conductance channel in mitochondrial membrane → mitochondrial permeability transition pore
↓
Loss of mitochondrial membrane
↓
Failure of NORMAL oxidative phosphorylation
↓
Depletion of ATP

Mitochondrial damage
↓
Abnormal oxidative phosphorylation
↓
Formation of reactive oxygen species (ROS)
↓
Deleterious effects

Mitochondrial damage
↓
Increased permeability of outer mitochondrial membrane
↓
Leakage of cytochrome c into cytosol
(Mitochondria sequester cytochrome c between their outer and inner membranes)
↓
Apoptosis





(3) Loss of calcium homeostasis

- Intracellular free calcium is very low compared with extracellular levels
- Most intracellular calcium is sequestered in mitochondria and ER.

Injury
↓
Increased influx and Release of Ca 2+ from intracellular stores ie. mitochondria and ER
↓
Increased cytosolic Ca 2+
↓
Activates many enzyme
↓
Phospholipases (membrane damage)
Proteases (breakdown proteins)
Endonucleases (DNA fragmentation)
ATPases (hastening ATP depletion)



(4) Free Radical Injury

Introduction→

- Normally generation of ATP by oxidative process→ oxygen (O2) combines with hydrogen atom (H), water (H2O) is formed
- This reaction of O2 to H2O involves **'four electron donation'** in four steps involving transfer of one electron at each step.



Generation

- Free radicals are intermediate chemical species having a single unpaired electron in its outer orbit.
- Unpaired electrons are highly reactive and "attack" inorganic or organic chemicals → proteins, lipids, carbohydrates, nucleic acids
- Generated within mitochondrial inner membrane



Types

3 types →

1. Superoxide oxygen (O_2^-): one electron
2. Hydrogen peroxide (H_2O_2): two electrons
3. Hydroxyl radical (OH^-): three electrons

Fenton reaction → iron is converted from ferrous to ferric form.

$$(H_2O_2 + Fe^{2+} \quad Fe^{3+} + OH^+ + OH^-)$$

Haber- Weiss reaction → generation of free radical (OH) during normal metabolic process

$$(H_2O_2 + O_2 \quad 2OH^- + O_2)$$

Effects

1. Damage proteins, lipids, carbohydrates, nucleic acids
2. Lipid peroxidation in cell membranes
3. Oxidation of amino acids and proteins resulting in fragmentation
4. Protein-protein cross linkages.



Removal

Scavengers of free radicals →
1. Superoxide dismutase
2. Glutathione peroxidase
3. Catalase
4. Vitamin E, vitamin C
5. Transferrin, ferritin and ceruloplasmin
   (Iron and copper catalyze free radical formation)



Superoxide dismutase (SOD)

- Present in mitochondria (manganese-SOD) and in cytosol (copper-zinc- SOD)
- Converts superoxide to hydrogen peroxide

$$20^- + 2H \rightarrow 2\ H_20_2 + 0_2$$

Glutathione peroxidase

- Present in mitochondria and cytosol
- Converts hydrogen peroxide to hydroxyl ion

$$H_20_2 \rightarrow OH^-$$

Catalase

- Present in peroxisomes
- Converts hydrogen peroxide to water

$$2H_20_2 \rightarrow 0_2 + 2H_20$$

(1.03) MORPHOLOGICAL CHANGES – REVERSIBLE VERSUS IRREVERSIBLE CELL INJURY

University Exams

Long questions

1. Compare and contrast structural changes in reversible and irreversible cell injury
2. Differences between reversible and irreversible cell injury
3. Morphological features of reversible and Irreversible cell injury

Short questions

1. Describe and Draw Morphological features of Reversible cell injury
2. Describe and Draw Morphological features of Irreversible cell injury

Overview

- Introduction
- Diagramatic presentation of Reversible injury and Irreversible Injury
- Irreversible Injury – Nuclear Changes
- Differences between Reversible injury and Irreversible Injury

Introduction



Diagrammatic presentation of Reversible injury and Irreversible Injury



Irreversible Injury – Nuclear Changes

- Pyknosis → Nuclear shrinkage and increased basophilia
- Karyorrhexis → Fragmentation of pyknotic nucleus
- Karyolysis → Fading of basophilia of chromatin



Differences between Reversible injury and Irreversible Injury

	Reversible injury	Irreversible Injury
Cell membrane	- Swell - Intact - Blebs	- Do not swell - Disrupts
ER	- Swell	- Lysis
Ribosomes	- Dispersion	- Dispersion
Mitochondria	- Swell - Small densities	- Swells - Large densities
Lysosomes	- Autophagy	- Rupture
Nucleus	- Clumping	- Pyknosis, Karyolysis, Karyorrhexis

(1.04) APOPTOSIS AND EFFEROCYTOSIS

University Exams

Long questions

1. What is apoptosis? Mention the causes and discuss mechanisms of apoptosis.

Short questions

1. Write diagnosis of apoptosis
2. Describe Mechanisms (pathogenesis) of apoptosis with diagrams
3. Describe intrinsic pathway of apoptosis
4. Describe extrinsic pathway of apoptosis
5. Describe Morphological features of apoptosis.
6. Draw Morphology of apoptotic cells.
7. Define Efferocytosis. Write its mechanism
8. Differentiate apoptosis and Necrosis

Very short questions

1. Morphology of apoptotic cells.
2. Define Apoptosis. Write its Types.

Overview

- Introduction
- Apoptosis → Definition
- Types
- Mechanisms →
    - Extrinsic pathway → Initiation phase
    - Intrinsic pathway → Initiation phase
    - Execution phase
    - Efferocytosis
- Morphological changes in apoptosis
- Diagnosis of Apoptosis
- Differences between Apoptosis and necrosis

Introduction

Cell Death →
- Apoptosis → Suicide
- Necrosis → Murder

Apoptosis → Definition

Cell Suicide

- It is a pathway of cell death that is induced by a tightly regulated intracellular program in which cells destined to die activate enzymes (caspase) degrade cells own nuclear DNA and cytoplasmic proteins.

- Cell is phagocytosed
- No leakage outside
- No inflammation

Types



Mechanism



Extrinsic pathway → Initiation phase

Initiated through specific receptors called death receptors

1. FAS protein (CD95)
2. TNF receptors

FAS receptor (Death receptor)
↓
FAS protein (CD95) binds with it
↓
Multiple FAS receptors come together
↓
Cytoplasmic death domain FADD formed
↓
Activate pro- caspase 8 and 10 to active caspase 8 and 10



Intrinsic pathway → Initiation phase

Stimuli→ DNA damage (beyond repair)

Stimuli (DNA damage)
↓
Anti apoptotic molecules Bcl-2, Bcl-x, Mcl -1 are lost
↓
Replaced by pro-apoptotic molecules like Bak, Bax, Bim, Bad
↓
Increased mitochondrial permeability (MPT)
↓
Release to cytochrome C into cytoplasm
↓
Activates Apaf-1 along with procaspase-9
↓
Activated caspase-9



Execution phase

Convergence point for both extrinsic and intrinsic pathways.





Efferocytosis

Immunologically silent clearance of apoptotic cells by professional and non-professional phagocytes (efferocytes)

- Healthy cells avoid being efferocytosed by displaying "don't eat-me" signals → eg. CD47
- Apoptotic cells carry specific 'eat me' signals – e.g. phosphatidylserine (resulting from phospholipid flip-flop)





Phosphatidylserine is a phospholipid
present on inner surface membrane normally
↓
During apoptosis
↓
Flipped to outer surface
↓
Externalization of phosphatidylserine
causes its tagging
↓
Phagocytes ingulf it

Morphological changes in apoptosis



Diagnosis of Apoptosis

1. Apoptosis markers → Annexin-V is a recombinant protein with high affinity for phosphatidylserine.



2. Agarose gel electrophoresis → Fragmented DNA shows Step Ladder Pattern → due to inter-nucleosomal cleavage of DNA by endonuclease



Differences between Apoptosis and necrosis

Apoptosis (Suicide)	Necrosis (Murder)
Single cells or small clusters of cells	Often contiguous cells
Cell shrinkage	Cell swelling
Pyknosis and karyorrhexis	Karyolysis, pyknosis and karyorrhexis
Intact cell membrane	Disrupted cell membrane
Cytoplasm retained in apoptotic bodies	Cytoplasm leaked
Inflammation absent	Inflammation present

(1.05) NECROSIS

University Exams

Long questions

1. Define necrosis. Mention the types, explain the causes and pathology of each type of necrosis.
2. Define necrosis. Enlist different types of necrosis with relevant examples

Short questions

1. Enumerate types of necrosis with appropriate examples and mention their mechanisms.
2. Define Coagulative necrosis. Write its causes and draw microscopic features
3. Define Colliquative necrosis. Write its causes and draw microscopic features
4. Define Caseous necrosis. Write its causes and draw microscopic features
5. Define Fat necrosis. Write its causes and draw microscopic features
6. Define Fibrinoid necrosis. Write its causes and draw microscopic features

Very short questions

1. Coagulative necrosis.
2. Liquifactive necrosis
3. Caseous necrosis.
4. Fat necrosis.
5. Fibrinoid necrosis.

Overview

- Definition
- Types →

    1. Coagulative necrosis
    2. Liquefactive necrosis
    3. Caseous necrosis
    4. Fat necrosis
    5. Fibrinoid necrosis

Definition

- Death of cells and tissues in the living animal
- Necrosis is defined as a localised area of death of tissue followed later by degradation of tissue by hydrolytic enzymes liberated from dead cells
- Accompanied by inflammatory reaction





Types

1. Coagulative necrosis
2. Liquefactive necrosis
3. Caseous necrosis
4. Fat necrosis
5. Fibrinoid necrosis

(1) Coagulative necrosis

Introduction →

- Most common type of necrosis
- Architectural outlines persist but cellular and nuclear details are lost → Ghost or tombstone appearance
- Type of tissue can be recognized

Causes

1. Ischemia due to thrombosis/ embolism in all organs except brain
2. Mild burns (thermal injury)
3. Zenker's degeneration necrosis



Gross

- Pale, firm, and slightly swollen

Microscopy

Hallmark → conversion of normal cells into 'tomb stones'

i.e., outlines of cells retained and cell type can still be recognised but their cytoplasmic and nuclear details are lost.



(2) Liquefactive necrosis

Introduction →

- Architectural, cytoplasmic and nuclear details are lost → semi-fluid material (due to degradation by Hydrolytic enzymes)
- Also known as Colliquative necrosis





Causes

1. Pyogenic bacterial infections
2. Ischemic necrosis of brain

Gross

- Liquefied center containing necrotic debris with a cyst wall



Microscopy

- Necrotic area → cavity containing a mass of necrotic neutrophils, bacteria and tissue debris
- Surrounded by a cyst wall formed by proliferating capillaries and inflammatory cells



(3) Caseous necrosis

Introduction →

- Architectural, cytoplasmic and nuclear details are lost → amorphous debris (resembling cottage cheese)
- Looks Cheese like → Caseous necrosis



Causes

- Mycobacterium tuberculosis
- Syphilis
- Histoplasma
- Coccidiomycosis

Gross

- Soft, granular and yellowish → resemble dry cheese

Microscopy

- Centre → structureless, eosinophilic material having scattered granular debris of disintegrated nuclei
- Surrounding tissue → granulomatous inflammatory reaction



(4) Fat necrosis

Introduction →

- Special form of cell death occurring at fat-rich anatomic locations in body.
- Death of adipose tissue in a living animal

Causes →

- Pancreatic (acute pancreatitis)
- Breast (Traumatic)
- Mesentery (Inflammation)



Gross

- Fat necrosis → yellowish-white and firm deposits.
- Formation of calcium soaps → chalky white appearance

Microscopy

- Necrosed fat cells → cloudy appearance
- Surrounded by inflammatory reaction.



(5) Fibrinoid Necrosis

Introduction →

- Deposition of fibrin-like material

Causes →

- Immunologic injury of vessel wall

Microscopy

- Brightly eosinophilic, hyaline-like deposition in the vessel wall.
- Surrounded by nuclear debris of neutrophils (leucocytoclasis)



	Coagulative	Liquefactive	Caseous	Fat	Fibrinoid
Introduction	- Arch ⊕ - Cyto ⊖ - Nucleus ⊖	- Arch ⊖ - Cyto ⊖ - Nucleus ⊖ } gel	- Arch ⊖ - Cyto ⊖ - Nucleus ⊖ } debris	Fat rich locations	Blood vessels
Causes	- Ischemia of all organs (Except Brain)	- Ischemia Brain - Pyogenic infection	- TB - Syphilis - Histoplasma - Coccidiomycosis	- Breast trauma - Pancreatitis - Mesentery inflammation	- Vasculitis
Gross	- Pale	- Liq. Centre with Cyst Wall	- Dry cheese like	- Yellowish white	-
Microscopy	Tomb- stone appearance	Cavity Cont. debris	Structureless material with granulomatous inflammation	Cloudy appearance	Fibrinoid (ribbon) like

(1.06) NECROTOSIS, PYROPTOSIS, FERROPTOSIS

University Exams

Short questions

1. Define Necroptosis. Write its mechanism.
2. Define Pyroptosis. Write its mechanism.
3. Define Ferroptosis. Write its mechanism.
4. Differentiate Necroptosis from apoptosis
5. Differentiate Pyroptosis from apoptosis

Overview

- Necrotosis
- Pyroptosis
- Ferroptosis

Necroptosis

Introduction →

- Hybrid of necrosis and apoptosis but considered as a part of necrosis.
- Programmed cell death without caspase activation.

Differences from apoptosis

Necroptosis differs from apoptosis by following features →
a) Shows cell swelling
b) Increased lysosomal permeability
c) Cell membrane damage
d) Presence of inflammation

So, they are considered as → Necrosis → caspase independent programmed cell death

Mechanism

Ligation of TNFRI
↓
Recruits receptor-interacting protein kinase 1 and 3 (RIPKI and RIPK3)
↓
RIPK3 phosphorylates a cytoplasmic protein called MLKL
↓
MLKL monomers assemble into oligomers
↓
Translocate from the cytosol to plasma membrane
↓
Cause plasma membrane disruption with inflammation
↓
Necroptosis



Pyroptosis

Introduction →

- Caspase (caspase 1,4,5 and 11) dependent programmed cell death
- Programmed cell death associated with antimicrobial responses during inflammation (innate immunity response).

Differences from apoptosis

Pyroptosis differs from apoptosis by following features →
a) Shows cell swelling
b) Cell membrane damage
c) Presence of inflammation
d) It involves enzyme Caspase-1,4,5, and 11

So, they are considered → caspase dependent programmed cell death

Mechanism

NOD like receptors (NIRs)
↓
Recognize Flagellin components of salmonella and shigella species
↓
Activation of inflammasome
↓
Activation of caspase-1, 4, 5)
↓
Cleaves precursor form of IL-1 to generate biologically active IL-1
↓
Cell Membrane Damage and Inflammation
↓
Pyroptosis



Ferroptosis

Introduction →

- Programmed cell death dependent on iron
- Characterized by the accumulation of lipid peroxides

Differences from apoptosis

- Dependent on iron
- Characterized by accumulation of lipid peroxides
- Master regulator is: Glutathione Peroxidase 4 (Gpx4)

Mechanism

Inactivation of Glutathione Peroxidase 4 (GPx4)
↓
Failure of glutathione-dependent antioxidant defenses
↓
Iron release in the cytosol
↓
Generate Reactive Oxygen Species (ROS) by Fenton Reaction
↓
ROS leads to lipid peroxidation (No repair since GPx4 is inactivated)
↓
Cell death
↓
Ferroptosis



(1.07) GANGRENE

University Exams

Long questions

1. Define gangrene and mention the differences between dry and wet gangrene

Short questions

1. Gangrene and its types.
2. Describe Dry gangrene.
3. Describe Wet gangrene.
4. Describe Gas gangrene.
5. Differences between dry and wet gangrenes.

Overview

- Definition
- Types
- Differences between dry and wet gangrenes
- Gas Gangrene

Definition

Necrosis of tissue associated with superadded putrefaction

GANGRENE = Necrosis + Putrefaction

Types

1. Dry gangrene → less bacterial superinfection → tissue appears dry
2. Wet gangrene → abundant bacterial superinfection→ tissue looks wet and liquefactive
3. Gas gangrene → A variant of wet gangrene

Differences between dry and wet gangrenes

Features	Dry Gangrene	Wet Gangrene
Site	Commonly limbs	Commonly in bowel
Mechanisms	Arterial occlusion	Blockage of both venous drainage and arterial obstruction
Bacteria	Bacteria fail to survive	Numerous present
Putrefaction	Limited due to very little blood supply	Marked due to stuffing of organ with blood
Macroscopy	Organ dry, shrunken and black	Part moist , soft, swollen, rotten and dark
Line of demarcation	Present	Absent
Prognosis	Better due to little septicaemia	Poor due to profound toxaemia
Type of necrosis	Variant of Coagulative necrosis	Liquifactive necrosis is superimposed on coagulative necrosis



Gas Gangrene

- Special form of wet gangrene
- Caused by gas-forming clostridia (gram-positive anaerobic bacteria)
- Swollen, edematous, painful and crepitant due to accumulation of gas bubbles of CO2 within tissues formed by fermentation of sugars by bacterial toxins

(1.08) PATHOLOGICAL CALCIFICATION

University Exams

Short questions

1. Describe Pathological calcification and its types
2. Describe Dystrophic calcification. Write its pathogenesis and sites
3. Describe Metastatic calcification. Write its pathogenesis and sites
4. Difference between Dystrophic & metastatic calcification.

Overview

- Definition
- Types →

    1. Dystrophic calcification
    2. Metastatic calcification

- Difference between Dystrophic & metastatic calcification
- Special stain for calcification

Definition

Deposition of calcium salts in tissues other than osteoid or enamel

Types



Dystrophic calcification

Pathogenesis

Membrane of dead and degenerated cell → damaged
↓
Phospholipid is released
↓
Phosphatases within phospholipid generate phosphate ions
↓
Calcium binds to phosphate ions
↓
Form calcium phosphate

Sites

Dead tissues

- Caseation eg. Tuberculosis
- Dead parasites like trichinosis, Onchocercosis.
- Fat necrosis
- Infarcts
- Thrombi
- Hematoma



Degenerative Tissues

- Atherosclerosis- Monkeberg sclerosis
- Damaged heart valves
- Infected lymph nodes
- Degenerating tumors



Metastatic calcification

Pathogenesis

Those organs that 'lose' acid
↓
Have an underlying alkaline compartment
↓
Alkaline internal component is susceptible to calcification

Sites

- Basement membrane & tubular lamina of kidney
- Pulmonary veins
- Alveolar wall of lung (most common site)
- Cornea & Conjunctiva
- Interstitial tissue of gastric mucosa
- Synovium of the joint
- Systemic arteries
- Tendons

Difference between Dystrophic & metastatic calcification

	Dystrophic Calcification	Metastatic Calcification
Tissue	Calcium deposits in dead and dying tissues	Calcium deposits in normal tissues
Calcium metabolism	Normal calcium metabolism	Deranged calcium metabolism
Calcium in blood	Normal serum calcium levels	Hypercalcemia
Reversibility	Irreversible	Reversible upon correction of metabolic disorder
Pathogenesis	Occurs due to increased binding of phosphates with necrotic and degenerative tissue — binds to calcium forming calcium phosphate precipitates	Increased precipitates of calciumphosphate due to hypercalcemia at certain sites (lungs, stomach, blood vessels, cornea)

Special stain for calcification

1. Von Kossa → black color
2. Alizarin red S → red staining



Von Kossa



Alizarin red S

(1.09) PIGMENTS

University Exams

Short questions

1. Exogenous and endogenous pigments.
2. Hemosiderosis.
3. Brown atrophy of heart.
4. Lipofuscin pigment.
5. Disorders of melanin pigmentation
6. Pathological pigmentation.
7. How is Hemosiderin pigment formed? Give its pathological significance.
8. What is hemosiderin and how it is demonstrated? Enumerate two causes of hemosiderosis.
9. Tattoos
10. Wear and tear pigment

Overview

- Definition
- Types
- Endogenous Pigments →

    1. Melanin
    2. Hemosiderin
    3. Lioifuschin

- Exogenous Pigments →

    1. Carbon
    2. Tatto

Definition

colored substances present in most living beings including humans.

Types



Endogenous Pigments

1. Melanin
2. Hemosiderin
3. Lioifuschin

(1) Melanin

Introduction →

- Brown-black
- Non-hemoglobin derived pigment
- Normally present in hair, skin, mucosa, choroid of eye, meninges and adrenal medulla.
- Synthesized in melanocytes (enzyme tyrosinase) present in basal cells of epidermis



Disorders

1. Generalized hyperpigmentation
2. Focal hyperpigmentation
3. Generalized hypopigmentation
4. Localized hypopigmentation

(1) Generalized hyperpigmentation:

a) Addison's disease → generalized hyper pigmentation of the skin
b) Chloasma observed during pregnancy → hyperpigmentation on skin of face, nipples, and genitalia under influence of estrogen.
c) Chronic arsenical poisoning → raindrop pigmentation of skin.

Addison's Disease

Chloasma

Chronic Arsenical Poisoning

(2) Focal hyperpigmentation:

a) Cäfe-au-lait spots in neurofibromatosis
b) Peutz-Jeghers syndrome → focal peri-oral pigmentation.
c) Melanosis coli → pigmentation of mucosa of the colon.
d) Melanotic tumors, both benign such as pigmented naevi and malignant such as melanoma
e) Dermatopathic lymphadenitis → melanin pigment in macrophages of lymph nodes

Cafe-au-lait spots

Peutz-Jeghers Syndrome-Perioral Pig.



Melanosis Coli



Dermatopathy lymphadenitis



Pigmented naevi

(3) Generalized hypopigmentation

- Albinism

(4) Localized hypopigmentation:

- Vitiligo

(2) Haemosiderin

Introduction →

Iron is stored in the tissues in 2 forms:
1. Ferritin (iron complexed to apoferritin)
2. Haemosiderin (formed by aggregates of ferritin)

Haemosiderin Light microscopy →
- Golden-yellow to brown pigment
- Granular

Prussian blue reaction

Iron can be identified by the Prussian blue histochemical reaction



Golden brown granular pigment Prussian blue reaction

Disorders

HAEMOSIDEROSIS

LOCALISED	GENERALISED (SYSTEMIC)
Local Tissues (Macrophages, fibroblasts, endothelial and alveolar cells)	**Parenchymal Deposits (Liver, Pancreas, Kidney, Heart, Skin) RE Cell (Liver, Spleen, Bone marrow)**
Examples : • Hemorrhage in tissues • Black eye • Brown induration lung • Infarction	**Examples:** • Acquired hemosiderosis (Chronic haemolytic disorders, blood transfusion, parenteral administration of iron) • Hereditary (Idiopathic) haemochromatosis (Increased absorption, genetic defect) • Excessive dietary intake (African iron overload)

(3) Lipofuscin

Introduction →

- Also known as wear and tear pigment = aging pigment = Lipochrome
- Not injurious to the cell but is important as a marker of past free-radical injury → Tell tale sign of free radical injury and lipid peroxidation.
- Composed of polymers of lipids and phospholipids in complex with protein.
- By light microscopy → coarse, golden-brown granular
- Perinuclear brown pigment.





Disorders

- Aging
- Severe malnutrition
- Cancer cachexia
- In heart muscle → change is associated with wasting of muscle → brown atrophy of myocardium

Melanin	Hemosiderin	Lipofuscin
Skin	Iron excess	Wear and tear pigment Sign of cell ageing
Brown	Brown	Brown
Uniform distribution 	Uniform distribution 	Perinuclear
Stain- **Masson Fontana** 	**Pearls Prussian blue** 	

Exogenous Pigments

1. Carbon
2. Tatto

(1) Carbon

Inhaled
↓
Phagocytosed by alveolar macrophages
↓
Transported through lymphatic channels to regional tracheobronchial lymph nodes.
↓
Aggregates of pigment blacken lymph nodes
↓
Anthracosis
↓
Induce emphysema or a fibroblastic reaction
↓
Coal workers' pneumoconiosis



(2) Tatto

Pigments like India ink, cinnabar and carbon
↓
Injected into dermis
↓
Taken up by macrophages
↓
Lies permanently in connective tissue



(1.10) CELLULAR AGEING

University exams

Short questions

1. Role of Telomerase in cellular ageing
2. Role of Sirtuin in cellular ageing
3. Morphological changes during ageing

Overview

- Introduction
- Pathogenesis of Ageing
- Telomere Theory
- Morphological changes during ageing
- Sirtuin

Introduction

Cellular ageing is result of a progressive decline in cellular function and viability caused by → Genetic abnormalities and accumulation of cellular damage because of exposure to exogenous influences

Pathogenesis of Ageing

1. Telomere Theory
2. Genetic control in invertebrates
3. Diseases of accelerated ageing
4. Oxidative stress hypothesis (free radical-mediated injury)
5. Hormonal decline
6. Defective host defenses
7. Failure to renew

Telomere Theory

- After a fixed number of divisions all somatic cells become arrested in a terminally nondividing state → senescence.
- Telomeres are short repeated sequences of DNA (TTAGGG) present at linear ends of chromosomes that are important for protecting DNA
- With every cell division, there is progressive shortening of telomere present at tips of chromosomes
- Telomere length is normally maintained by an enzyme called telomerase



With every cell division
↓
progressive shortening of telomere present at tips of chromosomes
↓
Shortened telomere is repaired by the presence of RNA enzyme → telomerase
↓
After repetitive mitosis (max 60 to 70 times) → Telomerase activity is terminated
↓
Lost telomere is not repaired
↓
DNA cannot be protected
↓
Activation of p53
↓
No mitosis
↓
Cellular senescence (Aging)



Telomerase activity →

- Highest → germ cells
- Lower levels → stem cells
- Very low → somatic tissues



Morphological changes during ageing

A) Cellular changes	• Decrease in cell size and number • Decreased in size and number of mitochondria • Detachment of ribosomes from ER • Increased number of phagolysosomal vacuoles • Defective DNA repair • Non-enzymatic glycosylation of protein
B) Connective tissue changes	• Loss of elasticity (wrinkling of skin) • ↑ BP due to decreased elasticity of artery • Ground glass substance changes (cataract) • Cartilage and bone changes (osteoarthritis osteoporosis)
C) Decreased immunity	• Decreased T cells, B cells, plasma cells, Ab

Sirtuin

- Most effective way to prolong life span is calorie restriction.
- Calorie restriction will induce sirtuin (family of NAO-dependent protein deacetylases).

Mechanism of action →

a) Inhibit metabolic activity.
b) Reduce apoptosis.
c) Stimulate protein folding.
d) Inhibits free radical damage.
e) Increase insulin sensitivity and glucose metabolism

(2.01) VASCULAR EVENTS OF ACUTE INFLAMMATION

University Exams

Long questions

1. Define inflammation. Mention the types. Explain the sequential vascular changes in acute inflammation.
2. Define inflammation. Write types of inflammation. Describe the cardinal signs of acute inflammation.
3. Discuss the vascular phenomenon of inflammation.
4. Describe various mechanisms of vascular leakage (increased vascular permeability) in inflammation.

Short questions

1. Write and explain Cardinal signs of inflammation.
2. Vascular changes in acute inflammation.
3. Differences between acute and chronic inflammation.

Very Short questions

1. Definition of acute inflammation.
2. Types of inflammation

Overview

- Inflammation →
    - Definition
    - Classification
- Acute inflammation
    - Cardinal Signs
    - Steps
    - Vascular events →
        1. Transient vasoconstriction of arterioles
        2. Persistent progressive vasodilatation
        3. Elevate the local hydrostatic pressure
        4. Increased vascular permeability
        5. Slowing or stasis
- Triple Response

Inflammation

Definition →

- Body defense reaction in order to eliminate or limit the spread of injurious agent, followed by removal of the necrosed cells and repair of damaged tissue
- WBC or leukocytes → body's major infection-fighting cells.

Classification

Acute	Chronic
Rapid onset	Late onset
Short duration	Longer duration
Odema	Granuloma formation
Neutrophils	Macrophage, lymphocyte



Acute inflammation

Cardinal Signs

	Latin	English	
1.	Rubor	: redness	Celsus
2.	Calor	: ↑ed local temperature	Celsus
3.	Tumor	: swelling	Celsus
4.	Dolor	: pain	Celsus
5.	Functio laesa	: loss of function	→ Virchow

Steps

Acute inflammation

→ **Vascular events**

→ **Cellular reaction**

Vascular Events

1. Transient vasoconstriction of arterioles
2. Persistent progressive vasodilatation
3. Elevate the local hydrostatic pressure
4. Increased vascular permeability
5. Slowing or stasis

Cellular Events

1. Margination and pavementing
2. Rolling
3. Adhesion
4. Transmigration (diapedeses)
5. Chemotaxis
6. Phagocytosis



Vascular events

(1) Transient vasoconstriction of arterioles

Responsible for blanching seen immediately after injury.

- With mild injury→ 3-5 seconds
- Severe injury→ 5 minutes



(2) Persistent progressive vasodilatation

Vasodilatation results in increased blood volume→ redness(rubor) and warmth (calor)



(3) Elevate the local hydrostatic pressure

Starling's hypothesis





- Transudation of fluid into extracellular space (Oedema→ Transudate) → Tumor

(4) Increased vascular permeability

- Hallmark of acute inflammation
- Leads to escape of protein rich fluid (Oedema → Exudate) → swelling (tumor)
- Most affected vessels are venules





Remember

Initial stage
↓
Elevation in hydrostatic pressure
↓
Escape of fluid
↓
Transudate

Subsequently
↓
Increased vascular permeability
↓
Escape of fluid, protein and cells
↓
Exudate

Altered Vascular Permeability mechanisms

a) Formation of endothelial gaps





b) Direct endothelial injury





c) Leukocyte mediated endothelial cell injury



d) Increased transcytosis





e) Leakage from new blood vessels





Mechanism	Vasculature	Response	Pathogenesis
1. Endothelial gaps	Venules	Immediate transient	Histamine
2. Direct endothelial injury	All levels	Immediate prolong	Direct injury
3. Leukocyte mediated injury	Venules	Delayed prolong	Activated WBC
4. Transcytosis	Venules		Histamine and VEGF
5. Leaky new blood vessels	---		VEGF

(5) Slowing or stasis

Increased concentration of RBCs → raised blood viscosity



Triple Response

- Demonstrated by Lewis experiment
- Lewis induced changes in skin of inner aspect of forearm by firm stroking with a blunt point.

Red line Flare Wheal



1. Red line → due to local vasodilatation of capillaries and venules.
2. Flare → bright reddish surrounding red line and results from vasodilatation of adjacent arterioles
3. Wheal → swelling due to transudation of fluid into extravascular space



(2.02) CELLULAR EVENTS OF ACUTE INFLAMMATION

University Exams

Long questions

1. Define inflammation. Describe the cellular changes that take place in acute inflammation?
2. Define inflammation. Enumerate cellular events in inflammation and discuss in detail about phagocytosis.

Short questions

1. Ennumerate Cellular events in acute inflammation.
2. Explain Leukocyte transmigration.
3. Describe Chemotaxis.
4. Mechanism of Phagocytosis and killing
5. Describe chemotaxis and phagocytosis.
6. Describe Rolling and adhesion during Cellular events in acute inflammation.

Very Short questions

1. Chemoattractants in acute inflammation.
2. Name three steps in phagocytosis

Overview

- Acute inflammation → Steps
- Summary of vascular events

- Cellular Events →

    1. Margination and pavementing
    2. Rolling
    3. Adhesion
    4. Transmigration (diapedeses)
    5. Chemotaxis
    6. Phagocytosis → Steps

Acute inflammation → Steps

Vascular Events

1. Transient vasoconstriction of arterioles
2. Persistent progressive vasodilatation
3. Elevate the local hydrostatic pressure
4. Increased vascular permeability
5. Slowing or stasis

Cellular Events

1. Margination and pavementing
2. Rolling
3. Adhesion
4. Transmigration (diapedeses)
5. Chemotaxis
6. Phagocytosis

Summary of vascular events



Cellular Events

1. Margination and pavementing
2. Rolling
3. Adhesion
4. Transmigration (diapedes)
5. Chemotaxis
6. Phagocytosis





(1) Margination

- Normally → central stream of WBCs and RBCs and peripheral cell-free layer of plasma close to vessel wall → Normal axial flow
- Acute inflammation → WBCs leave enter and comes at periphery of blood vessel → Reversal of axial blood flow



(2) Rolling

Transient Bond between WBCs and endothelial cells → Rolling



(3) Adhesion

Bond between WBCs and endothelial cells becoming firmer → Adhesion



Complementary adhesion molecules (CAM)

Endothelial Molecule	Leukocyte Molecule	Major Role
1. P-selectin	Sialyl-Lewis X	Rolling
2. E-selectin	Sialyl-Lewis X	Rolling and adhesion
3. GlyCam-1 (CD34)	L-selectin	Rolling
4. ICAM-1	B2 integrins	Adhesion
5. VCAM-1	B1 integrins	Adhesion
6. PECAM-1 (CD-31)	PECAM-1 (CD-31)	Diapedesis (trasmigration)





(4) Transmigration (diapedes)

- WBCs Escape out into extravascular space
- Passive phenomenon
- PECAM



(5) Chemotaxis

Unidirectional oriented along a chemical gradient
Potent chemotactic substance →

1. LT-B4
2. C3a
3. C5a → most powerful
4. Interleukines (IL-8)



(6) Phagocytosis

- Engulfment of microbes by the WBC's (cell-eating)
- Appreciated by Metchnikoff (1880)



Steps of Phagocytosis

1. Recognition and Attachment
2. Engulfment
3. Killing and degradation → 3 steps

   a) MPO-dependent
   b) MPO-independent



Step 1

NADPH-oxidase present in cell membrane of phagosome reduces oxygen to superoxide ion (O2-)



Step 2

Superoxide is subsequently converted into H2O2:

$$2O_2 + 2H^+ \longrightarrow H_2O_2$$
(Hydrogen peroxide)

Step 3

MPO-independent killing

- Enzyme MPO acts on H_2O_2 in presence of halides (chloride, iodide or bromide) → form hypohalous acid (HOCl, HOI, HOBr) → Antibacterial action

$$H_2O_2 \xrightarrow[Cl', Br', I']{MPO} HOCL + H_2O$$

(Hypochlorous acid)

MPO-independent killing

- Mature macrophages lack enzyme MPO
- Carry out bactericidal activity by producing OH^- ions from H_2O_2 → Antibacterial action

✓ In presence of O_2 (Haber-Weiss reaction)
✓ In presence of Fe++ (Fenton reaction)





(2.03) CHRONIC INFLAMMATION OR GRANULOMATOUS INFLAMMATION

University Exams

Long questions

1. Define and classify granuloma. Explain the evolution and morphology of tuberculous granuloma.
2. Define granuloma. Explain the pathogenesis of granulomatous diseases
3. Define granuloma. Explain its morphology. List the diseases with granulomatous inflammation.
4. Define Chronic inflammation. How does it differ from acute inflammation. Describe in brief chronic infective granuloma.
5. A 42-year-old male presented with history of fever, cough and weigh loss since two months. X-ray of chest showed cavitary lesion in right apical lobe.
6. What is your diagnosis and why?
7. Describe the etiopathogenesis of this disease.
8. Describe microscopic findings in lung.

Short questions

1. Granulomatous lymphadenitis.
2. Discuss granulomatous inflammation with examples.
3. Microscopic appearance of granuloma.
4. Write short note on Granuloma.
5. Define granuloma. Write its types with examples.
6. Write short note on Giant cells.
7. Chemicals of tuberculous bacilli responsible for producing granuloma.
8. Microscopy of tuberculous granuloma with a diagram

Very Short Answer

1. What are various types of giant cells? Name them and conditions with which they are associated.
2. Define Granuloma.
3. List types of granulomas with example.

Overview

- Introduction
- Granuloma →
    - Pathogenesis
    - Definition
    - Structure or microscopy
- Types of Giant cells
- Types of Granulomas
- Examples of granulomatous inflammation

Introduction

Chronic inflammation is a response of prolonged duration (weeks or months) in which inflammation, tissue injury and attempts at repair (fibrosis) coexist.

Granuloma →

Pathogenesis

Formation of granuloma is a type IV hypersensitivity reaction

3 Steps →

1. Engulfment by macrophages
2. Activation of CD4+ T cells
3. Release of Cytokines



(1) Engulfment by macrophages

Macrophages engulf Ag
↓
Try to destroy it
↓
But Ag is poorly degradable
↓
Macrophages fail to digest and degrade Ag

(2) Activation of CD4+ T cells

Macrophages failed to deal with Ag
↓
Present antigen to CD4+ T lymphocytes
↓
CD4+ T lymphocyte activated
↓
Release lymphokines
↓
a) Interferon-Y
b) IL-1, IL-2
c) TNF-α

(3) Release of Cytokines

- Interferon-γ → activates macrophages and transform it into epitheloid cells
- IL-1 and IL-2 → T cells proliferation
- TNF-α → Fibroblast proliferation

Definition

A focus of chronic inflammation consisting of a microscopic aggregation of →
- Macrophages that are transformed into epithelioid cells
- Epithelioid cells fuse to form → giant cells
- Surrounded by a Collar of mononuclear leukocytes → lymphocytes and plasma cells.

Structure or Microscopy

- Macrophages modified as epithelioid cells in center
- Some epitheloid cells fuse to form multinucleate giant cells.
- Surrounded peripherally by T lymphocytes
- Healing by fibroblasts or collagen
- Caseous necrosis in some granulomatous conditions



Types of Giant cells

1. Foreign body giant cells
2. Langhans' giant cells
3. Trouton giant cells
4. Giant cells in tumors

INFLAMMATORY GIANT CELLS			TUMOUR GIANT CELLS		
					
Foreign Body Type	**Langhans' Type**	**Touton Type**	**Anaplastic Tumour Giant Cell**	**Reed-Sternberg Cells**	**Osteoclastic Tumour Giant Cell**

Foreign body giant cells →

Nuclei are scattered throughout cytoplasm.
E.g. chronic infective granulomas, leprosy and tuberculosis

Langhans' giant cells →

Nuclei are arranged either around periphery in form of horseshoe ring or clustered at two poles of giant cell.
E.g. tuberculosis and sarcoidosis.

Touton giant cells→

Vacuolated cytoplasm due to lipid content
E.g. Xanthoma

Types of Granulomas

Caseating Granuloma	Non-caseating Granuloma
1. Tuberculosis 2. Histoplasmosis 3. Syphilis 4. Coccidiomycoses 	1. Tuberculosis 2. Sarcoidosis 3. Hodgkin's lymphoma 

Naked Granulomas →

Not have surrounding rim of mononuclear cells
- Composed of epithelioid histiocytes and multinucleated giant cells
- Example → Sarcoidosis

Stellate granuloma →

Star shaped Granuloma
- Granuloma with central neutrophilic infiltrates.
- Examples →
    - Cat scratch disease
    - LGV (lymphogranuloma venereum)

Examples of Granulomatous Inflammation

Bacterial

1) Tuberculosis	Mycobacterium tuberculosis
2) Leprosy	Mycobacterium leprae
3) Syphilis	Treponema pallidum
4) Granuloma inguinale Donovanosis)	C. donovani
5) Brucellosis (Mediterranean fever)	Brucella abortus
6) Cat scratch disease	Coccobacillus
7) Tularaemia	Francisella tularensis
8) Glanders	Actinobacillus mallei

Fungal

1. Actinomycosis	Actinomycetes israelii
2. Blastomycosis	Blastomyces demtitidis
3. Cryptococcosis	Cryptococcus neoformans
4. Coccidioidomycosis	Coccidioidesimmitis

Parasitic

1. Schistosomiasis (Bilharziasis)	Schistosoma mansoni, haematobiumjaponicum

Miscellaneous

1.Sarcoidosis	Unknown
2.Crohn's disease	Unknown
3. Silicosis	Silica dust
4.Berylliosis	Metallic beryllium
5.foreign body granulomas	Talc, suture, oils, wood splinter etc.

(2.04) CHEMICAL MEDIATORS OF INFLAMMATION

University Exams

Long questions

1. Define inflammation. Discuss in detail about chemical mediators of inflammation.
2. Define chemical mediators of inflammation. Classify them. Describe functions of plasma derived chemical mediator.
3. Classify chemical mediators of inflammation. Describe the functions of Arachidonic acid derivatives.

Short questions

1. Chemical mediators of inflammation.
2. Cell-derived mediator of Inflammation.
3. Plasma-derived mediator of Inflammation.
4. Role of arachidonic acid metabolites in acute inflammation.
5. Cytokines.
6. Chemokines.
7. Classify chemical mediators of inflammation.

Very short questions

1. Name the cell-derived mediator of Inflammation.
2. Role of prostaglandins in acute inflammation.
3. Thromboxane A2.
4. Platelet activation factor.
5. Histamine and Serotonin.

Overview

- Classification
- Cellular Inflammatory mediators

Performed

a) Histamine
b) Serotonin
c) Lysosomal enzymes
d) Neuropeptides

- Cellular Inflammatory mediators

Newly Synthesized

a) Nitric Oxide
b) Platelet activating factor
c) Cytokines
d) Arachidonic acid derivative

- Plasma Inflammatory mediators

a) Kinin and clotting system
b) Complements

Classification

2 types →

a) Cellular (mediators present within cells)
b) Plasma (synthesized by liver and secreted in plasma)



Cellular Inflammatory mediators

Performed

1. Histamine
2. Serotonin
3. Lysosomal enzymes
4. Neuropeptides

(1) Histamine

- Synthesized from histidine amino acid.
- Richest source are mast cells.

Functions

a) Vasodilation mainly (but induce vasoconstriction of large arteries)
b) Increased vascular permeability.
c) Increased smooth muscle contraction (bronchoconstriction).

(2) Serotonin

- 5-hydroxytryptamine, 5-HT
- Richest source are platelets
- Also present in enterochromaffin cells.
- Function is same as of histamine.

(3) Lysosomal Enzymes

- Present in Lysosomes of neutrophil and monocytes.
- Lysosomal granules are of three types:

  a) Primary (Azurophilic)
  b) Secondary (Specific)
  c) Tertiary granules

Primary granules	Secondary granules	Tertiary granules
• Azurophilic granules • Develop at promyelocytic stage • More destructive **It contains** • MPO (myeloperoxidase) • Elastase • Defensin • Cathepsin G • Bacterial permeability proteins • Acid hydrolase • Phospholipase- A_2 • Non-specific collagenase	• Specific granules • Develop at myelocytic stage • Less destructive **It contains** • Lactoferrin • Lysozyme • Type IV collagenase • B1 microglobulin • Gelatinase • Vitamin B_{12} binding proteins • Cytochrome B	• c-particle • Mainly involved in chemotaxis diapedesis and matrix degradation **It contains** • Gelatinase

(4) Neuropeptides

- Examples are substance p & neurokinin A.
- Secreted by sensory nerves and leukocyte.
- Nerve fibers containing substance p is rich in lung and gastrointestinal tract.

Actions are:

a) Mediator of vascular permeability.
b) Pain Signal transmission.
c) Blood pressure' regulation.
d) Stimulates immune and endocrine cell secretion.

Cellular Inflammatory mediators

Newly Synthesized

Formed during process of inflammation.
a) Nitric Oxide
b) Platelet activating factor
c) Cytokines
d) Arachidonic acid derivative

(a) Nitric oxide (NO)

Synthesized from arginine amino acid by enzyme nitric oxide synthase (NOS).

Actions are →

a) Vasodilation
b) Platelet anti-aggregators
c) Microbicidal action

(b) Platelet-activating factor (PAF)

- Phospholipid derived mediator.
- Most potent inflammatory mediators

Actions are:

a) Platelet aggregation
b) Vasoconstriction
c) Bronchoconstriction
d) Increased vascular permeability
e) Transmits signals" between cells.

(c) Cytokines

- Soluble polypeptides.
- Secreted by both hematopoietic and nonhematopoietic cells.
- Highly specific
- Involved in inflammation, immunological reactions and wound healing.

Chemokines

- Family of small (8 to 10 KO) proteins.
- Stimulate leukocyte movement (chemotaxis)
- Perform their action by binding to seven-transmembrane G protein-coupled receptors.

	C-X-C (α-chemokines)	C-C (b-chemokines)	C (g chemokines)	CX_3C
FUNCTION	• Neutrophil recruitment	• Recruit monocytes, eosinophils, basophils and lymphocytes but not neutrophil.	• Recruit lymphocytes	
MEMBERS	• **IL-8** (produced by macrophages and endothelial cells) activation by IL-1 and TNF-a	• **Eotaxin** (for eosinophil) • Monocyte chemo attractant protein-1 • **RANTES** (Regulated and Normal T-cells Expressed and Secreted) • **MIP 1a** (macrophage Inflammatory protein 1a)	• **Lymphotactin**	• **Fractalkine** It exists in two forms: • Endothelial surface bound protein Promotes, mononuclear cell adhesion • Soluble form Chemoattractant for mononuclear

(d) Arachidonic acid (AA) metabolites

- Arachidonic acid is a 20-carbon polyunsaturated fatty acid derived from essential fatty acid Linoleic acid.
- Activated cells release membrane-bound arachidonic acid (AA) through enzymatic activity of phospholipase A2.
- AA is then catabolized to generated short-range Lipid mediators (eicosanoids) through enzymatic activities.
- Eicosanoids bind to membrane G-protein- coupled receptors and involve in almost every aspect of inflammation.

Arachidonic acid Metabolite

Phospholipids
↓
Phospholipase A
↓
Arachidonic acid

- **Cycloxygenase pathway**
  1. Prostaglandins
  2. Prostacyclins
  3. Thromboxane A_2(TxA_2)
- **Lipoxygenase pathway**
  1. Leukotrienes
  2. Lipoxin A and B
- **Cytochrome p450 epoxygenase pathway**
  1. EET (epoxyeicosatetraenoic acid)
  2. HETE (hydroryelcosatetraenoic acid)
- **Iso-eicosanoid pathway**
  1. Iso-prostanes

Cyclo-oxygenase pathway

AA
↓
Cyclo-oxygenase pathway (cox)
↓
PGG2 and PGH2

- **PGI2 (prostacyclin)**
  - **Synthesised In endothelium**
    - Inhibits platelet aggregation
    - Vasodilatation
- **TXA,(thromboxane A_2)**
  - **Formed by platelets**
    - Platelet aggregation
    - Vasoconstriction
- **PGD_{2a} PGE_{2a} PGF_{2a}**
  - ↑ Vasodilation
  - ↑ Vascular permeability
  - PGE_{2a} →↑is for pain and fever
  - PGD_{2a} &PGF_{2a} for neutrophil chemotaxis

LOX (Lipoxygenase) pathway

Arachidonic acid
↓
Lipoxygenase pathway (LOX)

- **5-LOX**
  ↓
  LTA_4
  - **LTB_4**
    Chemotaxis
  - **LTC_4 LTD_4 LTE_4**
    Slow-reacting (SRS-A) substance of anaphylaxis.
    ---
    1. Vasoconstriction
    2. Increase vascular permeability
    3. Bronchoconstriction
- **12-LOX**
  ↓
  Lipoxin (Endogenous negative regulator of leukotrienes)
  - Anti inflammatory

  a. Inhibit neutrophil chemotaxis
  b. Inhibits NK-cell cytotoxicity
  c. Activates monocytes and macrophage for phagocytosis of neutrophils

Epoxygenase pathway



Iso-eicosanoid pathway



Plasma Inflammatory mediators

a) Kinin and clotting system
b) Complement

(a) Kinin and clotting system

- Inflammation promotes clotting by increasing production of several coagulation factors.
- Thrombin promotes inflammation by binding to protease-activated receptors (PARs) on platelets, endothelium, and smooth muscle cells.
- This mobilizes P-selectin to induce cytokine and chemokine production and increases synthesis of PAF, NO, and prostaglandins.

Kinin system



(b) Complement

3 major pathways →

1. Classic pathway
2. Mannose Binding Lectin (MBL) pathway
3. Alternative pathway

Complement Pathways



(3.01) WOUND HEALING (REGENERATION AND REPAIR)

University Exams

Long questions

1. What do you understand by the terms-"healing", "regeneration" and "repair"? Describe steps of Repair.
2. Define repair. Describe its steps in detail

Short questions

1. Differentiate Regeneration and Repair
2. Granulation tissue.
3. Scar
4. Angiogenesis (neovascularizalion) in repair.
5. Steps of wound healing.

Overview

- Healing →
  - Definition
  - Mechanisms
- Repair → 3 Phases →
  1. Inflammatory phase
  2. Proliferative phase
     a) Epithelialization
     b) Fibroplasia
     c) Angiogenesis → Steps
  3. Maturation phase

Healing

Definition →

Body's response to injury in an attempt to restore normal structure and function

Mechanisms

Regeneration

- Parenchyma replaced by parenchymal cells of the same type
- Restoration of normal structure and function
- Result - Same tissue



Repair

- Parenchyma replaced by fibrous tissue
- Restoration of normal shape and function is impaired
- Result - scar

Repair → 3 Phases

1. Inflammatory phase
2. Proliferative phase

   a) Epithelialization
   b) Fibroplasia
   c) Angiogenesis

   } Granulation tissue

3. Maturation phase: Scar

(1) Inflammatory Phase

Within seconds of the injury
↓
Release of epinephrine, norepinephrine, prostaglandins, serotonin, and thromboxane
↓
Vasoconstriction
↓
Endothelial cells retract
↓
Expose the subendothelial collagen surfaces
↓
Platelets adhesion
↓
Platelet Activation
↓
Platelet Aggregation → Release Neutrophils and macrophages ↓ Inflammation
↓
Primary platelet plug formation
↓
Coagulation cascade
↓
Secondary plug (fibrin product)
↓
Hemostasis

Inflammatory phase



(2) Proliferative Phase



(a) Epithelialization

- Cell travel about 3 cm from point of origin in all directions
- Fibrogenesis

(b) Fibroblasts lay bed of collagen

- Fills defect

(c) Angiogenesis

Steps of angiogenesis

1. Vasodilation → in response to NO
2. Increased permeability → induced by VEGF
3. Separation of pericytes → from abluminal surface
4. Breakdown of basement membrane → to allow formation of a vessel sprout
5. Migration of endothelial cells → toward area of tissue injury
6. Proliferation of endothelial cells → just behind leading front tip of migrating cells
7. Recruitment of periendothelial cells → to form mature vessel



- End product of Proliferation Phase → Granulation tissue

(3) Maturation Phase

New collagen forms
↓
Collagen is remodelled into a more organized matrix
↓
Increases tensile strength to wounds
↓
Fibroblasts leave wound
↓
Final product → scar



- Tensile strength never reaches the normal strength of unwounded tissue (i.e., 100%).
- Maximum strength is 70-80% of normal tissue. (at end of 3 months)



Inflammation **Granulation tissue** **Scar**

(3.02) HEALING OF SKIN WOUNDS (PRIMARY AND SECONDARY INTENTION)

University Exams

Long questions

1. Describe and compare with help of suitable diagrams → wound healing by primary and secondary intention.
2. What is repair? Describe healing of wound by first intention.

Short questions

1. Describe healing. Write down the differences between primary and secondary healing.
2. Mechanism of wound healing by first intention (primary union).
3. Mechanism of wound healing by secondary intention (secondary union).

Overview

- Healing of Skin Wounds →

    1. Primary Intention (primary union)
    2. Secondary Intention (Secondary union)

- Differences between primary union and Secondary union

Healing of Skin Wounds

(1) Primary Intention (primary union):

- No separation of wound edges
- All Layers are closed.
- In a minimum amount of time
- Minimal scar formation.

(2) Secondary Intention (Secondary union):

- Wound edges separated
- Deep layers are closed but superficial layers are left to heal from the inside out.
- More time required
- Abundant scar formation.

Healing by Primary Intention (primary union)



Healing by Secondary Intention (Secondary union)



Differences between primary union and Secondary union

Features		Primary Union	Secondary Union
1.	Cleanliness of wound	Clean	Unclean
2.	Infection	Uninfected	May be infected
3.	Margins	Surgical, clean	Irregular
4.	Sutures	Used	Not used
5.	Healing	Scanty granulation tissue	Exuberant granulation tissue
6.	Outcome	Neat linear scar	Contracted irregular scar
7.	Complications	Infrequent	Frequent

(3.03) FACTORS AFFECTING WOUND HEALING AND COMPLICATIONS OF WOUND HEALING

University Exams

Short questions

1. Enumerate factors affecting wound healing.
2. Differentiate Hypertrophic scar and Keloid
3. Describe complications of wound healing.

Very Short questions

1. Enumerate Local factors which delay wound healing.
2. Enumerate Systemic factors influencing wound healing.

Overview

- Factors Affecting Wound Healing
- Complications of Wound Healing →

  1. Hypertrophic scar
  2. Keloid

  } - Introduction
  - Gross
  - Microscopy

- Differences between Hypertrophic scar and Keloid

Factors Affecting Wound Healing

Local	Systemic
1. Infections 2. Decreased blood supply 3. Denervation 4. Foreign body 5. Poor surgical technique 6. Hematoma 7. Mechanical stress	1. Increased age 2. Anemia (Hypoxia) 3. Steroidal or cytotoxic drugs 4. Diabetes 5. Obesity 6. Malignancy 7. Malnutrition • Vitamins A, Cand E deficiency • Deficiency of polyunsaturated long fatty acids • Zinc deficiency 8. Uremia

Complications of Wound Healing

Insufficient fibrosis:

- Wound dehiscence
- Hernia
- Ulceration

Excessive fibrosis:

- Hypertrophic scars
- Keloid

(1) Hypertrophic scar

Introduction →

- Excessive collagen accumulation forms a raised scar within wound boundaries.
- Onset → less than 3 months.
- Spontaneous regression → within 12 to 18 months.

Gross →

Scar tissue do not grow beyond boundaries of original wound and may regress spontaneously.





Microscopy →

Scar tissue with thin collagen bundles that is oriented parallel to skin surface.

(2) Keloid

Introduction →

- Can progression beyond original area of injury without subsequent regression.
- Onset → within 1 months to 1 year.
- May be genetically associated.
- Most common site affected → sternum.
- No spontaneous regression.

Gross →

Scar Tissue extends beyond borders of original wound and it does not usually regress spontaneously.



Microscopy →

well demarcated haphazardly arranged and thicker dense eosinophilic collagen fibers involving dermis.

Differences between Hypertrophic scar and Keloid

	Hypertrophic scar	Keloid
Gross	▪ Remains within boundaries of wound ▪ Spontaneously regression ▪ Tend not to reoccur if excised ▪ Most often affect the skin overlying joints	▪ Outgrows wound area ▪ No spontaneously regression ▪ Tend to reoccur if excised ▪ Affect variety of sites- ear lobes, shoulders, anterior chest wall, upper back
Microscopy	▪ Larger number of myofibroblasts ▪ Extracellular matrix composed of type III collagen in linear fibrils running parallel with the skin ▪ Highly vascular	▪ Fewer myofibroblasts ▪ Extracellular matrix composed of type I collagen arranged chaotically in thick bundles that project under normal epidermis ▪ Less vascular
Epidemiology	▪ Non-inheritable	▪ Inheritable
Etiology	▪ Only occur following injury	▪ May occur without injury
Symptoms	▪ Mild pruritis, not painful	▪ Pruritic and painful

(4.01) EDEMA AND EFFUSION

University Exams

Long questions

1. Discuss causes and pathogenesis of edema.
2. Classify and list different pathophysiologic categories of edema and describe pathogenesis of cardiac edema.
3. Define Edema. Discuss in detail etiopathogenesis of edema. Describe different types of edema with examples.
4. Define oedema. Discuss pathogenesis of oedema in congestive cardiac failure and renal disease.

Short questions

1. Differences between transudates and exudates.
2. Define oedema. Give classification of oedema.
3. Define edema. Mention the types and write pathogenesis in brief.
4. Cardiac edema → pathogenesis
5. Renal edema → pathogenesis
6. Draw a figure showing pathways leading to systemic edema from primary heart failure.

Very short questions

1. Transudate.
2. Exudate
3. Define oedema and effusion

Overview

- Oedema → Definition
- Effusion → Definition
- Normal fluid exchange (Starlings law)
- Pathogenesis of oedema
- Types of oedema

Oedema→ Definition

Abnormal and excessive accumulation of fluid in interstitial tissue space



Effusion→ Definition

Abnormal and excessive accumulation of fluid in body cavities





Normal fluid exchange



At arterial end→

- Hydrostatic pressure (outward force) > osmotic pressure (inward force)
- **Net Outward driving force = 38 - 25 = 13 mmHg**
- Fluid **comes out** from capillary into interstitial space.

At venous end →

- Osmotic pressure (inward pressure) > hydrostatic pressure (outward pressure)
- **Net Inward-driving force = 25 -12 = 13 mmHg**
- Fluid **comes back in** capillary from interstitial space.



Pathogenesis of edema

1. Increased capillary hydrostatic pressure
2. Decreased plasma oncotic pressure
3. Lymphatic obstruction
4. Sodium and water retention
5. Increased capillary permeability

1) Increased capillary Hydrostatic pressure

- **Oedema of cardiac disease** e.g. in congestive cardiac failure
- **Postural oedema** e.g. transient oedema of feet and ankles due to increased venous pressure seen in individuals Whose job involves standing for long hours such as traffic constables



2) Decreased plasma oncotic pressure

- Oncotic pressure → pressure exerted by total plasma proteins.
- Albumin has four times higher plasma oncotic pressure than globulin
- Hypoalbuminemia → Oedema
- Edema takes place when →
    - Total plasma protein is below 5 gm/dl (normal 6-8 gm/dl)
    - Albumin is below 2.5 gm/dl (normal 3.5- 5gm/dl)



Examples →

Inadequate synthesis of albumin → end-stage cirrhosis
Protein malnutrition
Increased loss of albumin → Nephrotic syndrome

3) Lymphatic Obstruction

- Normally interstitial fluid in tissue spaces escapes lymphatics.
- Obstruction to outflow of lymphatics → L**ymphoedema**

Examples →

- **Removal of axillary lymph nodes** in radical mastectomy for carcinoma of breast → lymphoedema of affected arm.



- **Inflammation of the lymphatics as seen in filariasis (elephantiasis)**→ lymphoedema of scrotum and legs → elephantiasis



- **Milroy's disease or hereditary lymphoedema** → due to abnormal development of lymphatic channels



4) Sodium And Water Retention

- **Oedema of cardiac disease** e.g. in congestive cardiac failure.
- **Oedema of renal disease** e.g. in nephrotic and nephritic syndrome.
- Increased capillary permeability



5) Increased capillary permeability

Inflammation
↓
Capillary endothelium is injured
↓
Gaps between endothelial cells
↓
Leakage of plasma proteins into interstitial fluid
↓
Reduced plasma oncotic pressure and elevated oncotic pressure of interstitial fluid
↓
Oedema



Examples

- **Generalized oedema →** in systemic infections, poisonings, certain drugs and chemicals, anaphylactic reactions and anoxia.
- **Localised oedema →** allergic reactions, insect-bite, irritant drugs and chemicals.

Generalized oedema

Localised oedema

Pathogenesis of edema

Hydrostatic pressure	Plasma osmotic pressure	Lymphatic obstruction (Lymphedema)	Sodium retention	Capillary permeability
• Cardiac failure	• Liver cirrhosis	• After breast surgery	• Cardiac failure	• Acute and chronic inflammation
• Postural oedema	• Malnutrition	• Filariasis	• Renal failure	
	• Renal failure • (Nephrotic syn)	• Milroy disease		

Types of oedema

Features	Transudate	Exudate
Definition	Protein-poor, cell-poor fluid	Protein-rich, cell-rich fluid
Protein content	Low (less than 1 gm/dl)	High (2.5-3.5 gm/dl)
Cells	Few cells	Many cells
LDH	Low	High
Specific gravity	Low (less than 1.015)	High (more than 1.018)
pH	> 7.3	< 7.3
Glucose content	Same as in plasma	Low (less than 60 mg/di)
Character	Non-inflammatory oedema	Inflammatory oedema
Examples	Oedema in congestive cardiac failure	Purulent exudate such as pus


Hydrostatic pressure
Colloid osmotic pressure
Normal
Plasma proteins
Fluid leakage
Increased hydrostatic pressure
• venous outflow obstruction,
[E.g. congestive heart failure]
Decreased colloid osmotic pressure
• decreased protein Synthesis [E.g. liver disease]
• increased protein loss [E.g. kidney disease]
Transudate
(Low protein content, few cells)
Fluid & protein leakage
Exudate
(High protein content & may contain some white & red cells)
Vasodilation & stasis
Increased interndothelial spaces
Inflammation

(4.02) HYPEREMIA AND CONGESTION (CVC LUNG, CVC LIVER, CVC SPLEEN)

University Exams

Long questions

1. Define hyperemia and Congestion. Describe types of congestion. Describe gross & microscopic features of lung & liver in chronic passive venous congestion.

Short questions

1. Differences between hyperemia and Congestion
2. Chronic venous congestion (CVC) liver → Causes, gross and microscopic features.
3. Chronic venous congestion (CVC) lungs → Causes, gross and microscopic features.
4. Chronic venous congestion-spleen → Causes, gross and microscopic features.
5. Heart failure cells
6. Nutmeg liver.
7. Gamma-Gandy bodies.

Very short questions

1. Microscopy of CVC lung with a diagram.
2. Microscopy of CVC liver with a diagram
3. Microscopy of CVC spleen with a diagram
4. Brown induration of lungs

Overview

- Hyperaemia And Congestion →
  - Definition
  - Differences
- Congestion →
  - In Left heart failure
  - In Right heart failure
- CVC of lung
- CVC of liver
- CVC of spleen

} Gross, Microscopy

Hyperaemia And Congestion

Definition →

Localised increase in volume of blood within dilated vessels of an organ or tissue.



Differences

	Hyperemia	Congestion
1.	Active process	Passive process
2.	Vasodilatation of artery	Impaired venous blood flow
3.	During exercise & in inflammation	Venous obstruction & cardiac failure
4.	Oxygenated blood (Red, Erythema)	Deoxygenated blood (Blue, Cyanosis)

Congestion

In left heart failure →

- Pulmonary congestion (CVC lungs)

In right heart failure →

- Systemic venous congestion (CVC of systemic organs →Liver, spleen)



CVC Lung

Gross

- C/S → dark and rusty brown → brown induration of lungs

Microscopy

- Alveolar capillaries→ congested.
- Initially→ excess fluid collects in interstitial lung spaces → Interstitial oedema
- Later → fluid fills alveolar spaces → Alveolar oedema
- Hemosiderin-containing macrophages in lung alveoli → Heart failure Cells





CVC Liver



Gross

- C/S → Nutmeg appearance → due to red (congested centre of lobules) and yellow (Fatty peripheral zone) mottled appearance

Microscopy

- Red → Centrilobular haemorrhagic necrosis
- Yellow → fatty change in peripheral zone of lobule
- So nutmeg appearance





CVC Spleen



Gross

- Congested, tense and cyanotic
- C/S → gray tan

Microscopy

- **Red pulp** → enlarged due to marked sinusoidal dilatation
- **Capillarization of sinusoids** → Sinusoids may get converted into capillaries
- Fibrous tissue get deposits of hemosiderin pigment → Gamna-Gandy bodies



Remember

- CVC of lung → presence of **heart failure cells** (hemosiderin laden macrophages)
- CVC of liver → produces **nut-meg liver**
- CVC of spleen → presence of **Gamna-Gandy bodies**

	CVC Lung	CVC Liver	CVC Spleen
Gross	**Brown induration**	**Nutmeg liver**	**Cyanotic, grey-tan**
Microscopy	– **Interstitial oedema** ↓ – **Alveolar oedema** – **Heart Failure Cells**	– **Centri lobular→ Hemorrhagic necrosis** – **Peripheral→ Fatty Change**	– **Red pulp enlargement** – **Gamna-gandy body**

(4.03) THROMBOSIS

University Exams

Long questions

1. Define and describe pathogenesis of thrombus. Add a note on fate of thrombus.
2. Define and classify thrombosis. Explain the etiopathogenesis and pathology of thrombosis. Add a note on fate of thrombus.
3. Discuss causes, pathogenesis and effects of venous thrombosis.
4. Discuss the pathogenesis, types, morphology & fates of a thrombus.

Short questions

1. Fate of a thrombus.
2. What is Virchow triad? Discuss its role in thrombus formation.
3. Define and classify thrombosis
4. Types of thrombus.
5. Mural thrombus.
6. Difference between arterial and venous thrombus.

Very short questions

1. 3 primary conditions associated with increased risk for thrombosis.
2. Fate of thrombus.

Overview

- Introduction
- Pathophysiology →

    1. Endothelial injury
    2. Altered blood flow
    3. Hypercoagulable States

- Types of thrombus
- Gross
- Differentiate Arterial vs Venous thrombus
- Fate of thrombus

Introduction

- **Haemostasis** → occurs **after injury to CVS** and is **useful** as it stop escape of blood and plasma
- **Thrombosis** → occurs in unruptured **CVS (without injury)** and has **harmful effects** of ischemia
- Thrombosis is formation of a blood clot (solid mass) in unruptured CVS from constituents of flowing blood → obstructin flow of blood



Pathophysiology



Virchow's triad →
1. Endothelial injury
2. Altered blood flow (Stasis or turbulence)
3. Hypercoagulable States (Thrombophilia)

1) Endothelial Injury

Intact endothelium has both **antithrombotic property and prothrombotic property** → So no thrombosis

Prothrombotic

Thromboplastin or tissue factor→ released from endothelial cells.
↓
Von Willebrand factor → causes adherence of platelets to subendothelium.
↓
Platelet activating factor → activator and aggregator of platelets.
↓
Inhibitor of plasminogen activator→ suppresses fibrinolysis.

Antithrombotic

Heparin-like substance → accelerates action of antithrombin III
↓
Thrombomodulin → converts thrombin into activator of protein C
↓
Tissue plasminogen activator → accelerates fibrinolytic activity.
↓
Inhibitors of platelet aggregation → s ADPase, PGI2 (or prostacyclin).

Endothelial injury
↓
Exposes vWF and tissue factor (prothrombotic)
↓
Thrombosis

Steps →
a) Arteriolar vasoconstriction
b) Primary hemostasis
c) Secondary hemostasis

Arteriolar vasoconstriction

Occurs immediately and is transient



Primary hemostasis

Formation of the platelet plug





Injury
↓
Disruption of the endothelium
↓
Exposes subendothelial collagen
↓
1. Platelet adhesion
↓
2. Platelet activation
↓
3. Platelet aggregation/recruitment
↓
Primary hemostatic plug



1. Platelet adhesion → Platelets adhere to collagen in subendothelium due to presence of receptor (Gp) Ia-IIa and Gp Ib-IX
    - Vwf forms a bridge between collagen of endothelium and platelet

2. Platelet activation → 2 types of granules
    a) Alpha -granules
    b) Dense granules

3. Platelet aggregation → Adherence of platelets to one another due to presence of receptor Gp IIb-IIIa

Secondary hemostasis

Deposition of fibrin





Injury
↓
Tissue factor binds and activates factor VII or facror XII
↓
A cascade starts (Intrinsic,extrinsic, common)
↓
Fibrinogen into insoluble fibrin
↓
Fibrin meshwork is formed
↓
Secondary hemostasis

2) Alterated Blood Flow

Normally
↓
Platelets separated from endothelium by a slower moving layer of plasma
↓
Axial or Laminar flow of blood

Turbulence and stasis
↓
Bring platelets into contact with the Endothelium
Prevent washout of activated clotting factors by fresh flowing blood
Prevents inflow of clotting factor inhibitors
↓
Normal axial flow of blood is disturbed
↓
Thrombosis



A, NORMAL AXIAL FLOW



B, MARGINATION AND PAVEMENTING

Turbulence → Arterial and cardiac thrombi

Stasis → Venous thrombi

3) Hypercoagulable States (Thrombophilia)

Hypercoagulability (also called thrombophilia) → any disorder of blood that predisposes to thrombosis.

2 Types→

1. Hereditary or primary
2. Acquired or secondary

INHERITED (PRIMARY) FACTORS

i. Deficiency of antithrombin III
ii. Deficiency of protein C
iii. Deficiency of protein S
iv. Mutation in factor V Leiden
v. Defects in fibrinolysis (dysfibrinogenaemia, plasminogen disorders)
vi. Increased levels of coagulations factors (II and VIII)

ACQUIRED (SECONDARY) FACTORS

a) Risk factors: i) Advancing age, ii) prolonged bed-rest, iii) prolonged immobilisation (e.g. in plaster cast, long distance travel), iv) cigarette smoking, v) obesity
b) Predisposing clinical conditions:

   i. Heart diseases (e.g. myocardial infarction, CHF, rheumatic mitral stenosis, cardiomyopathy)
   ii. Vascular diseases (e.g. atherosclerosis, aneurysms of the aorta and other vessels, varicosities of leg veins)
   iii. Hypercoagulable conditions (e.g. polycythaemia, myeloproliferative disorders, dehydration, nephrotic syndrome, disseminated cancers)
   iv. Shock
   v. Tissue damage e.g. trauma, fractures, burns, major surgery on bones, abdomen or brain.
   vi. Late pregnancy and puerperium
   vii. Certain drugs (e.g. anaesthetic agents, oral contraceptives, hormonal replacement therapy).

c) Antiphospholipid antibody (APLA) syndrome:

i. Lupus anticoagulant antibody
ii. Anti-cardiolipin antibody

- Most common inherited cause of thrombosis → mutation in **factor V gene called Leiden mutation** → formation of defective (mutated) factor V called factor V Leiden → causes unchecked coagulation.

Types of Thrombi

3 types depending on site of origin→

1. Cardiac thrombi (vegetations)
2. Arterial thrombi
3. Venous thrombi



Gross

- Arterial thrombus → white thrombi → contains more platelets and relatively less fibrin
- Venous thrombi → red thrombi → contain more enmeshed red cells and relatively few platelets.



Differentiate Arterial vs Venous thrombus

Features	Arterial thrombus	Venous thrombus
Pathogenesis	Endothelial injury or turbulence	Stasis
Blood flow	Active	Sluggish
Sites	Coronary, cerebral, femoral arteries	Superficial and deep leg veins
Propagation	Retrograde manner	Antegrade manner
Lines of Zahn	Present	Absent
Microscopic	Pale platelet layer alternating with dark red cell layer so also called as white thrombi	RBC mixed with relatively less platelets, so also called as red thrombi
Occlusion	Incomplete	Complete
Complications	Ischemia and infarction	Edema and ulceration





Fate of Thrombus



(4.04) EMBOLISM

University Exams

Long questions

1. Define embolism. Discuss different types of embolisms.
2. What is embolism? Mention different types of embolism with examples. Add a note on air embolism and fat embolism.
3. A 20-year-old male suffered fracture femur in road traffic accident and was treated. 2 weeks later he suddenly collapsed and died. What is the diagnosis? What are the postmortem findings relevant to diagnosis?
4. What is embolism? Describe local and distant effect on organ due to different types of emboli.
5. Define embolism. Discuss different types of embolisms. Write in brief about thromboembolism.
6. Define embolus. Enumerate different types of embolism. Write in brief about pulmonary embolism.

Short questions

1. Embolism-types.
2. Thromboembolism
3. Fat embolism.
4. Air embolism.
5. Amniotic fluid embolism.
6. Pulmonary thromboembolism.

Very short questions

1. Define and list different types of emboli.
2. Paradoxical embolism.
3. Decompression sickness
4. Arterial and venous effects of embolism

Overview

- Definition
- Classification →

    a) Depending upon matter in emboli
    b) Depending upon whether infected or not
    c) Depending upon source of emboli
    d) Depending upon flow of blood

- Effects of Arterial embolism
- Effects of Venous embolism
- Important embolism →

    1. Thromboembolism
    2. Fat Embolism
    3. Amniotic Fluid Embolism
    4. Gas Embolism → Decompression Sickness



Definition

An embolus is a detached intravascular solid, liquid, or gaseous mass that is carried by blood from its point of origin to a distant site, where it often causes tissue dysfunction or infarction.

Classification

a) Depending upon matter in emboli
b) Depending upon whether infected or not
c) Depending upon source of emboli
d) Depending upon flow of blood

a) Depending upon matter in emboli

1. **Solid** e.g. **detached thrombi (thrombo-emboli),** atheromatous material, tumor cell clumps, parasites, bacterial clumps, foreign bodies.
2. **Liquid** e.g. **fat globules, amniotic fluid**
3. **Gaseous** e.g. **air**

b) Depending upon whether infected or not:

1. Bland → when sterile
2. Septic → when infected

c) Depending upon source of emboli

1. Arterial emboli
2. Venous emboli

d) Depending upon flow of blood

1. **Paradoxical embolus** → An embolus which is carried from venous side of circulation to arterial side or vice versa e.g. through patent foramen ovale
2. **Retrograde embolus** → An embolus which travels against flow of blood



Effects of arterial emboli

Depend upon their size, site of lodgement, and adequacy of collateral circulation

1. Infarction of organ
2. Gangrene in lower limbs
3. Myocardial infarction → following coronary embolism.
4. Sudden death → from middle cerebral artery embolism

Effect of venous embolism

Embolus in vein
↓
Flows through venous drainage into larger veins (SVC and IVC)
↓
Right side of heart (RA → RV)
↓
Pulmonary artery
↓
Pulmonary embolism



Important embolism

1) Thromboembolism

- A detached thrombus or part of thrombus
- Most common type of embolism (90%)



2) Fat Embolism

Fat globules



Etiology

Traumatic

Trauma to bones and soft tissue

Non traumatic

Extensive burns
Diabetes mellitus
Fatty liver
Pancreatitis
Sickle cell anaemia
Decompression sickness
Inflammation of bones and soft tissues
Extrinsic fat or oils introduced into the body
Hyperlipidaemia
Cardiopulmonary bypass surgery

Pulmonary fat embolism
↓
In patients **fractures of bones**
↓
Sudden Death
↓
On Autopsy
↓
Presence of numerous fat emboli in capillaries lung

Special stains for fat



Sudan black



Oil red O



3) Amniotic Fluid Embolism

Most serious, unpredictable and unpreventable cause of maternal mortality.

During labor and in immediate postpartum period
↓
Contents of amniotic fluid may enter uterine veins
↓
Reach right side of heart
↓
Pulmonary embolism

Composition

- Amniotic fluid components:
- Epithelial squames
- vernix caseosa
- Lanugo hair
- Bile from meconium
- Mucus



4) Gas Embolism

Air, nitrogen and other gases can produce bubbles within circulation → obstruct blood vessels → damage to tissue.

2 forms →

1. Air embolism
2. Decompression sickness



Decompression Sickness
Specialized form of gas embolism

Also known as
- Caisson's disease
- Divers' palsy
- Aeroembolism

Pathogenesis

Divers or workers in caissons (diving-bells)
↓
Descend to high atmospheric pressure
↓
Increased amount of atmospheric gases (N_2, O_2, CO_2) dissolved in blood
↓
When such an individual ascends too rapidly
↓
Dissolved gases comes out as minute bubbles
↓
Decompression sickness



Effects

- **Bends**→ pain in joints, ligaments and tendons.
- **Chokes**→ occur due to accumulation of bubbles in Lungs → acute respiratory distress.
- **Cerebral effects** → vertigo, coma, and sometimes death.

(4.05) ISCHEMIA

University Exams

Long questions

1. Define Ischemia. Write its etiology. Ennumerate Factors determining severity of Ischaemic injury.
2. Define Ischemia. What are its adverse effects.

Short questions

1. Define ischaemia.
2. Adverse effects of ischaemia.
3. Factors determining severity of Ischaemic injury.
4. Types of hypoxia.

Overview

- Definition
- Types
- Etiology
- Adverse effects of ischaemia → Hypoxia
- Factors determining severity of Ischaemic injury

Definition →

Deficient blood supply to part of a tissue relative to its metabolic needs

Types →

1. Complete (complete ischaemia)
2. Partial (partial ischaemia)

Etiology

1. Causes in heart
2. Causes in arteries
3. Causes in veins

1. Causes in heart

- **Inadequate cardiac output** resulting from heart block, ventricular arrest and fibrillation

2. Causes in arteries

- Luminal occlusion of artery (intraluminal):

  a) Thrombosis
  b) Embolism



- Outside pressure on an artery (extramural):

  a) Ligature
  b) Tourniquet
  c) Tight plaster, bandages
  d) Torsion

- Causes in arterial walls (intramural):

  a) Vasospasm
  b) Arteriosclerosis
  c) Polyarteritis nodosa
  d) Thromboangitis obliterans
  e) Severed vessel wall

3. Causes in veins

- Luminal occlusion of vein (intraluminal):

  a) Thrombosis of mesenteric veins
  b) Cavernous sinus thrombosis

- Causes in the vessel wall of vein (intramural):

  a) Varicose veins of the legs

- Outside pressure on vein (extramural):

  a) Strangulated hernia
  b) Intussusception
  c) Volvulus

Adverse effects of ischaemia → Hypoxia

- Hypoxic hypoxia: due to low oxygen in arterial blood.
- Anaemic hypoxia: due to low level of haemoglobin in blood.
- Stagnant hypoxia: due to inadequate blood supply
- Histotoxic hypoxia: low oxygen uptake due to cellular toxicity.



Factors Determining Severity of Ischaemic Injury

1. Anatomic pattern of arterial blood supply
2. Type of tissue affected
3. Rapidity of development
4. Degree of vascular occlusion

1) Anatomic pattern of arterial blood supply→

4 different patterns

a) Single arterial supply without anastomosis
b) Single arterial supply with rich anastomosis
c) Parallel arterial supply
d) Double blood supply

a) Single arterial supply without anastomosis—>

- Results in ischaemic necrosis

Example:
  a) Central artery of retina
  b) Interlobular arteries of kidneys

b) Single arterial supply with rich anastomosis

- Blockage of one vessel → **re-establish blood supply** bypassing blocked arterial branch
- **Infarction is less common**

Example:
1. **Superior mesenteric artery** supplying blood to small intestine.
2. **Inferior mesenteric artery** supplying blood to distal colon.

c) Parallel arterial supply

- Vitality of tissue is maintained by **alternative blood supply** in case of occlusion of one

Example:
a) Blood supply to the brain in region of **circle of Willis.**
b) Arterial supply to forearm by **radial and ulnar arteries.**

d) Double blood supply

- Dual blood supply

Example:
a) Lungs are perfused by bronchial circulation as well as by pulmonary arterial branches.
b) Liver is supplied by both portal circulation and hepatic arterial flow.

2) Type of tissue affected

- Vulnerability of tissue to ischaemia is variable
- More vulnerable tissues →
    - **Brain** (cerebral cortical neurons).
    - **Heart** (myocardial cells).
    - **Kidney** (epithelial cells of PCT).

3) Rapidity of development

- **Sudden** vascular obstruction → more severe effects (there is less time for collaterals to develop)
- **Gradual** vascular obstruction → less severe effects

4) Degree of vascular occlusion

- Complete obstruction → more severe ischaemic injury
- Partial occlusion → less severe ischaemic injury

(4.06) INFARCTION

University Exams

Long questions

1. Define and classify Infarction. Types of infarction.
2. Define and classify Infarction. Write its gross and microscopic features

Short questions

1. Infarct → types with examples.
2. Differentiate between red and white infarcts.

Very short questions

1. Define infarction. What are the different types of infarcts with common sites of occurrence.
2. Organs involved in pale and red infarction.
3. Kidney infarct.

Overview

- Definition
- Etiology
- Types of infarcts
- Gross
- Microscopy



Definition

An infarct is an area of ischemic necrosis caused by occlusion of either arterial supply or venous drainage.

Etiology

1. Interruption in arterial blood supply
2. Venous obstruction

Types of Infarcts

1. According to their age
2. According to presence or absence of infection
3. According to their colour

1) According to their age:

- Recent or fresh
- Old or healed

2) According to presence or absence of infection:

- Bland → when free of bacterial contamination
- Septic → when infected

3) According to their colour:

Pale or anaemic
- Caused by arterial occlusion
- Seen in compact organs e.g. in the kidneys, heart, spleen.

Red or haemorrhagic
- Caused either by pulmonary arterial obstruction (e.g. in the lungs) or by venous occlusion
- Seen in soft loose tissues

Feature	Red infarcts	White infarcts
Cause	• Arterial occlusion in loose tissues or organs having dual blood supply • Venous occlusion (in ovarian torsion)	• Arterial occlusion in solid organs with end arterial circulation
Affected organs	Lung and small intestine	Solid organs (heart, spleen, kidney)
Properties	Ill define hemorrhagic margins which change in color to brown	Well defined margins and progressively paler with time
Edema	Usually present	Usually absent

Red infarct in lung

White infarct in spleen

Gross

Wedge-shaped→
- Apex pointing towards occluded artery
- Wide base on surface of organ

Wedge shaped infarct

Microscopy

- All infacts → coagulative necrosis
- Exception → Cerebral infarcts → liquefactive necrosis



All organ infarct



Cerebral infarct

(4.07) SHOCK

University Exams

Long questions

1. Define shock. Enumerate types of shock and discuss pathogenesis and morphological changes in shock.
2. Discuss pathogenesis of septic shock. Describe morphological changes in various organs in shock.
3. Define shock. What are different types of shock? Describe pathogenesis of septic shock.
4. Define shock and describe pathogenesis of cardiogenic shock.
5. Define shock and describe pathogenesis and clinical features of hypovolemic shock.

Short questions

1. Define shock. Discuss endotoxic (septic) shock.
2. Hypovolumic / Hemorrhagic shock.
3. Discuss pathogenesis of cardiogenic shock.
4. Morphological features of shock.

Very Short questions

1. Definition and causes of shock.
2. Enumerate 3 major types of shock with suitable examples.

Overview

- Introduction
- Classification –

1. Hypovolemic shock 2. Cardiogenic shock 3. Septic shock	Introduction Causes Pathogenesis Symptoms

- Morphological features of Shock

Introduction

- Shock is clinical syndrome that results from poor tissue perfusion

Poor tissue perfusion
↓
Body do not receive enough oxygen and nutrients to allow cells to function
↓
Cellular death
↓
Organ failure
↓
Whole body failure
↓
Death

Classification

1. Hypovolemic shock
2. Cardiogenic shock
3. Septic shock

1) Hypovolemic shock

Introduction →

- Most common form of shock
- Occurs due to decreased blood volume

Causes →

1. Acute haemorrhage
2. Dehydration from vomiting's, diarrhoea
3. Burns
4. Excessive use of diuretics
5. Acute pancreatitis

Pathogenesis



Symptoms

1. Increased heart rate (tachycardia)
2. Low blood pressure (hypotension)
3. Low urinary output (oliguria to anuria)
4. Alteration in mental state (agitated to confused to lethargic)

3 stages→

- Mild hypovolemia or stage I → < 20% volume loss → Only mild tachycardia is there with normal BP.
- Moderate hypovolemia or stage II → 20-40% volume loss → Tachycardia with normal BP in supine position but hypotension in erect posture.

- Severe hypovolemia or stage III → > 40% volume loss → Classical signs of shock appear e.g. tachycardia, hypotension, disorientation etc.

2) Cardiogenic shock

Introduction →

- Caused by inadequate pumping action of heart

Causes

1. Deficient emptying e.g.

    a) Myocardial infarction
    b) Cardiomyopathies
    c) Rupture of the heart, ventricle or papillary muscle
    d) cardiac arrhythmias

2. Deficient filling e.g.

    a) Cardiac tamponade from haemopericardium

3. Obstruction to the outflow e.g.

    a) Pulmonary embolism
    b) Ball valve thrombus
    c) Tension pneumothorax
    d) Dissecting aortic aneurysm



Pathogenesis

- Acute Circulatory Failure
- Cardiogenic shock occurs when > 40% of left ventricle is destroyed.

Symptoms

- Decreased tissue perfusion (tissue hypoxia) → Forward failure
- Pulmonary oedema → Backward failure

3) Septic (Toxaemic) shock

Introduction →

- Severe bacterial infections or septicaemia

Causes →

1. Gram-negative septicaemia **(endotoxic shock)** e.g. E.coli, Proteus, Klebsiella, Pseudomonas and Bacteroides
2. Gram-positive septicaemia **(exotoxic shock)** e.g. streptococci, pneumococci

Pathogenesis





Symptoms

2 Stages →

1. **Early hyperdynamic stage** → vasodilatation → Tachycardia and warm extremities.
2. **Late hypodynamic stage** → Increased vascular permeability → decreased output, and oliguria (renal failure).

	Hypovolemic Shock	Cardiogenic Shock	Septic Shock
Causes	• Hemorrhage • Vomiting • Diarrhea • Burns • Diuretics	• MI • Arrhythmias • Cardiomyopathies • Tamponade	• Gram Positive Septicaemia • Gram Negative Septicaemia
Pathogenesis	↓ Circulatory Blood Volume	Acute Circulatory Failure	Bacteria→Macrophage activation ↓ Cytokines VD ↑ Permeability
Symptoms	• ↑ HR • ↓ BP • ↓ Urine Output • Altered Mental Status	• Tissue Hypoxia • Pulmonary Oedema	• Hyperdynamic circulation (warm extremities) • Inflammatory Oedema

Morphological changes in shock

Brain	Ischemic encephalopathy
Heart	Coagulative necrosis
Liver	Fatty change with hemorrhagic central necrosis
Kidneys	Acute tubular necrosis → oliguria and electrolyte disturbances
Adrenal	Cortical cell lipid depletion
GIT	Hemorrhagic enteropathy
Lungs are uncommonly affected but may show features of diffuse alveolar damage or shock lung.	

(5.01) GENETIC INHERITANCE OF SINGLE GENE DISORDERS (AD, AR, XD, XR)

University Exams

Short questions

1. Describe pedigree and important features of Autosomal Dominant (AD) Inheritance Diseases with examples.
2. Describe pedigree and important features of X - linked recessive disorders (XR) with examples.

Very Short questions

1. Name 4 autosomal dominant disorders.
2. Ennumerate autosomal recessive disorders.
3. Name sex-linked inherited diseases.
4. X-linked disorders - give 4 examples.
5. Ennumerate X - linked recessive disorders.
6. Ennumerate X - linked dominant disorders.

Overview

- Normal human Karyotype
- Genetic inheritance of Single gene disorders →

1. Autosomal Dominant (AD) Diseases
2. Autosomal Recessive (AR) Diseases
3. X - Linked Dominant (XD) Disorders
4. X - Linked Recessive (XR) Disorder

} - Features
- Pedigree
- Examples

Normal human Karyotype

Out of a total number of 46 chromosome→

a) **22 pairs** → Autosomes
b) **23rd pair** → sex chromosomes.

- XX → Females
- XY → Male



Genetic inheritance of Single gene disorders

1. Autosomal Dominant (AD) Diseases
2. Autosomal Recessive (AR) Diseases
3. X - Linked Dominant (XD) Disorders
4. X - Linked Recessive (XR) Disorder



Pedigree Symbols

(1) Autosomal Dominant (AD) Diseases

Features →

- Mutated genes can express themselves in heterozygous state.
- Onset → Variable
- Generation skip → Not seen
- Gender bias → Do not show

Pedigree →



Examples

Vo Familial Hypercholesterolemia Autosomal DOMINANT Hai

Vo	**V**on Willebrand Disease
Familial	**F**amilial Adenomatous Polyposis
Hypercholesterolemia	**H**ypercholesterolemia (Familial)
Autosomal	**A**dult polycystic kidney
D	**D**ystrophia myotonica
O	**O**steogenesis imperfecta
M	**M**arfan syndrome
I	**I**ntermittent porphyria
N	**N**eurofibromatosis-1
A	**A**chondroplasia
N	**N**eurofibromatosis – 2
T	**T**uberous sclerosis
Hai	**H**untington's disease; **H**ereditary pherocytosis

(2) Autosomal Recessive (AR) Diseases

Features →

- Mutant genes express themselves only in Homozygous state.
- Onset → Early uniform (usually in childhood).
- Generation skip → seen
- Gender bias → Do not show

Pedigree →



Examples

ABCDEFGH SPW

A	-	**A**lbinism, Alpha 1 Antitrypsin Deficiency
B	-	**B**eta Thalassemia
C	-	**C**ystic Fibrosis, **C**GD, **C**AH
D	-	**D**eafness (SNHL), **D**ubin Johnson
E	-	**E**nzyme Deficiencies (Glycogen Storage and Lysosomal Storage)
F	-	**F**riedrich's Ataxia, **F**anconi Anemia
G	-	**G**alactosemia
H	-	**H**emochromatosis, **H**urler syndrome
S	-	**S**ickle Cell Disease
P	-	**P**henylketonuria
W	-	**W**ilson's disease

(3) X - Linked Dominant (XD) Disorders

Features →

- Affected male → transmit it to 100% of their daughters and none of their sons.
- Affected female → transmit it to 50% of their sons and daughters.

Pedigree →



Examples

FAIR

F - **F**acial **(oro)** syndrome
A - **A**lports
I - **I**ncontinenta pigmento
R - **R**esistant rickets

(4) X - Linked Recessive (XR) Disorder

Features →

- Mutant recessive gene **expresses itself in a male child** because it is not suppressed by a normal allele → So males are Cases
- Mutant recessive gene **do NOT expresses itself in a female child** because in female → presence of a normal allele on other X-chromosome prevents expression of disease → So females only act as carriers.

Pedigree →



Examples

GRAHAM BELL

G - **G**6 PD deficency
R - **R**etinitis pigmentosa
A - **A**ndrogen insensitivity
H - **H**emophilia A & B, Hydrocephalus
A - **A**drenoleukodystrophy
M - **M**enkes disease
B - **B**eckers & Duchenne's muscular dystrophy, **B**lindness (color blindness)
E - **E**mery- Dreifuss dystrophy
L - **L**esch nyhan syndrome
L - **L**owe disease

	AD	AR	XD	XR
Features				
Pedigree				
Examples				

(5.02) CYTOGENETIC ABNORMALITIES

University Exams

Short questions

1. Barr body
2. Aneuploidy
3. Down's syndrome
4. Klinefelter syndrome
5. Turner's syndrome
6. Patau syndrome
7. Lyon hypothesis
8. Trisomy 18 and Trisomy 13
9. Cytogenetic disorders involving Sex chromosomes

Overview

- Introduction
- Cytogenetic disorders involving Autosomes →

  1. Down Syndrome
  2. Patau syndrome
  3. Edwards syndrome

  } Introduction, Karyotype, Clinical Features

- Lyon hypothesis
- Cytogenetic Disorders Involving Sex Chromosomes →

  1. Klinefelter Syndrome
  2. Turner Syndrome

  } Introduction, Karyotype, Clinical Features

Introduction

- Gametes contain half the number of chromosomes (haploid) and are represented as **(23, X) or (23, Y).**
- An exact multiple of haploid chromosomes is called **Euploidy** (2n, 3n, 4n etc).
- When exact multiple of haploid chromosomes is not present, it is called **Aneuploidy.**
- Genome mutations involve loss or gain of whole chromosomes, giving rise to **monosomy or trisomy.**





Cytogenetic disorders involving Autosomes

1. Down Syndrome
2. Patau syndrome
3. Edwards syndrome

(1) Down Syndrome

Introduction →

- Trisomy 21
- Most common chromosomal disorder

3 mechanism →

- Meiotic nondisjunction (maternal non disjunction)
- Robertsonian translocation
- Mosaicism

Karyotype



Clinical features

- Flat facial profile
- Mental retardation
- Microgenia (abnormally small chin)
- Oblique palpebral fissures with epicanthic skin folds (mongoloid slant)
- Muscle hypotonia (poor muscle tone)
- Single palmar fold (Simian crease)
- Curvature of little finger towards other four fingers (Clinodactyly)
- Protruding tongue or macroglossia
- White spots on the iris known as Brushfield spots
- Excessive joint laxity including atlanto-axial instability
- Excessive space between large toe and second toe (Sandle toe)
- A single flexion furrow of the fifth finger



Complications

- Congenital cardiac defects → Most common is **ventricular septal defect**
- Increased risk of leukemia → More commonly **ALL**
- **Hypothyroidism**
- Males are totally **infertile and** Reduced fertility in females
- Virtually all patients with Down syndrome develops **Alzheimer disease** after 4th decade.

(2) Patau syndrome

Introduction →

Trisomy 13

Karyotype



Clinical features →

- Cleft Lip
- Flexed fingers with Polydactyly
- Low-set malformed ears
- Ocular hypotelorism
- Microphthalmia
- Cardiac malformations
- Scalp defects
- Hypoplastic or absent ribs

(3) Edwards syndrome

Introduction →

Trisomy 18

Karyotype



Clinical features →

- Closed fists with index finger overlapping the 3rd digit and the 5th digit overlapping the 4th
- Narrow hips with limited abduction
- Short sternum
- Rocker-bottom feet
- Microcephaly and prominent occiput
- Micrognathia
- Mental retardation



Lyon hypothesis (X chromosome inactivation)

- Only one of the X chromosomes is genetically active
- Other X undergoes heteropyknosis and is rendered inactive
- **Inactive X is seen in interphase nucleus** as a darkly staining small mass → Barr body
- Used to test **genetic femaleness.**
- **Number of Barr bodies = (N-l) (**where N number of X chromosomes)



Examples

	Genotype	Number of Barr bodies
In normal male	(XY)	(1 – 1 = O Barr body)
In normal female	(XX)	(2 – 1 = 1 Barr body)
In turner syndrome	(XO)	(1 – 1 = 0 Barr body)
In Klinefelter syndrome	(XXY)	(2 – 1 = 1 Barr body)
Superfemale	(XXX)	(3 – 1 = 2 Barr bodies)

Cytogenetic Disorders Involving Sex Chromosomes

1. Klinefelter Syndrome
2. Turner Syndrome

(1) Klinefelter Syndrome

Introduction →

- Hypogonadism in phenotypic males resulting from **trisomy of sex chromosomes**
- Karyotype → **47, XXY**
- Most common causes of **hypogonadism in male.**

Karyotype



Clinical features

- Male infertility
- Eunuchoid body habitus
- Minimal or no mental retardation
- Failure of male secondary sexual characteristics
- Gynecomastia
- Atrophic testes
- Elevated FSH and estrogen.
- Low testosterone levels.
- Increased incidence of type 2 diabetes
- Most common cardiac abnormality is mitral valve prolapse.



(2) Turner Syndrome

Introduction →

- Hypogonadism in phenotypic females resulting from **monosomy of X chromosome.**
- Karyotype → (45, X)

Karyotype



Clinical features

- Lymph edema of neck, hands, and feet
- Short fourth metacarpal
- Webbing of neck

- Short stature
- Cubitus valgus
- Broad chest and widely spaced nipples.
- Primary amenorrhea
- Failure of the development of normal secondary sex characteristics
- Severely atrophic and fibrous ovaries
- Congenital heart diseases: Bicuspid aortic valve (most common cardiac defect); preductal coarctation of aorta



(6.01) HYPERSENSITIVITY REACTIONS

University Exams

Long questions

1. Define Hypersensitivity reactions. Write pathogenesis and examples of various types of Hypersensitivity reactions

Short questions

1. Type I hypersensitivity reaction (anaphylaxis) → pathogenesis and examples
2. Type II hypersensitivity reaction (Cytotoxic) → pathogenesis and examples
3. Type III hypersensitivity reaction (Immune complex) → pathogenesis and examples
4. Type IV hypersensitivity reaction (delayed type) → pathogenesis and examples
5. Type V hypersensitivity reaction → examples

Very Short questions

1. Types of hypersensitivity.
2. Atopy and Anaphylaxis.

Overview

- Introduction
- Classification

1. Type I (Allergy)
2. Type II (Cytotoxic)
3. Type V (Stimulatory Type)
4. Type III (Immune complex Mediated)
5. Type IV (T Cell-mediated)

} Pathogenesis, Examples

Introduction

- Immune response is a protective process but it may sometimes be injurious to the host.
- Hypersensitivity → Exaggerated state of normal immune response which results in adverse effects on body.

Classification

Coomb and Gel classification

1. Type I (Allergy)
2. Type II (Cytotoxic)
3. Type V (Stimulatory Type)
4. Type III (Immune complex Mediated)

} Immediate, Humoral immunity

5. Type IV (T Cell- mediated) → Delayed, Cell mediated immunity

1) Type I (Allergy) Reaction

Pathogenesis →

During first contact of host with the Ag
↓
Ag is captured by APC
↓
Presented to T cell which differentiates into TH2 cell
↓
TH2 cell releases IL-3, IL-4 and IL-5
↓
IL-4 causes activation of B cell
↓
Activated B cells differentiate to form IgE-secreting plasma cells
↓
IgE antibodies formed
↓
Bind to the Fc receptors present on mast cells
↓
Mast cells are now fully sensitized for next event
↓
During second contact with same Ag (shocking dose)
↓
IgE antibodies on surface of mast cells-basophils are activated
↓
Cause cell damage—membrane lysis
↓
Degranulation of mast cells-basophils
↓
Histamine, Serotonin, Vasoactive intestinal peptide (VIP), Leukotrienes B4 and D4, Prostaglandins, Platelet activating factor



Effects of granules:

1. Increased vascular permeability
2. Smooth muscle contraction
3. Early vasoconstriction followed by vasodilatation
4. Shock
5. Increased gastric secretion
6. Increased nasal and lacrimal secretions

2 forms →
- Acute, potentially fatal, systemic form → **Anaphylaxis**
- Recurrent non-fatal localised form → **Atopy**

Examples

All Allergies →
- Eczema
- Hay fever
- Asthma (atopy)
- Urticaria
- Anaphylactic shock
- Acute dermatitis

PCTS (proximal convulated tubuleS)
P - **P**K rxn
C - **C**asoni test
T - **T**heoblad smith phenomena
S - **S**chultz Dale phenomenon

2) Type II (Cytotoxic) Reaction

Pathogenesis →

Antigen on the surface of target cells
↓
Circulating B lymphocytes get activated
↓
Differentiate to form plasma cells
↓
Antibodies formed (IgG or IgM)
↓
Ag - Ab complex formed
↓
Activation of classical pathway of serum complement
↓
Generates activated complement componentC3b
↓
Activated C3b bound to the target cell acts as an opsonin
↓
Phagocytosis



Examples

My	**My**asthenia gravis
Blood	**Blood** transfusion reactions
Group	**G**oodpasture syndrome and **G**raves' disease
Is	**I**nsulin resistant diabetes, **I**TP
R	**R**heumatic fever
H	**H**yperacute graft rejection
Positive	**P**ernicious anemia and **P**emphigus vulgaris

3) Type V (Stimulatory Type) Reaction

Pathogenesis →

Modification of Type II hypersensitivity

Ag on target cell surface
↓
Ab interact with Ag
↓
Ag - Ab reaction enhances the activity of affected cell
↓
Cell proliferation and differentiation (instead of inhibition or killing)

Examples →

a) Grave's disease
b) Myasthenia gravis

4) Type III (Immune Complex Mediated) Reaction

Pathogenesis →

Soluble Antigen
↓
Circulating B lymphocytes get activated
↓
Differentiate to form plasma cells
↓
Antibodies formed
↓
Immune complexes are formed by interaction of soluble antibody and antigen
↓
Immune complexes get deposited into tissues
↓
Fc component of antibody links with complement and activates complement
↓
Formation of C3a, C5a and membrane attack complex
↓
Accumulated neutrophils and macrophages in the tissue release cytokines
↓
Inflammation and Tissue destruction



2 forms
- Arthus reaction (localised) →due to relative antibody excess
- Serum sickness (generalised) → due to relative antigen excess.

Examples

S	**S**erum sickness, **S**chick test and **S**LE
H	**H**enoch-Schonlein Purpura
A	**A**rthus reaction
R	**R**eactive arthritis, **R**aji assay
P	**P**olyarteritis nodosa (**P**AN) and **P**ost Streptococcal glomerulonephritis (**P**SGN)

5) Type IV (T Cell-mediated) Reaction

Pathogenesis →

Ag is recognised and processed by APC
↓
Presented to helper T cells (TH1 cells)
↓
TH1 cells release cytokines (TNF-α, IFN-γ, IL-2 and IL-12)
↓
Granuloma formation



Release of Cytokines

- IL-1 and IL-2 → Proliferation of more T cells.
- Interferon-γ → Activates macrophages and transform it into epitheloid cells
- TNF-α → Fibroblast proliferation

Examples

John gave TU LIP and Pop Corn to Hashi

John mote reaction
Tuberculin reaction
LePromin reaction
Pernicious anaemia
Contact dermatitis
Hashimoto thyroiditis

Remember

- Types I, II and III → Immediate → Antibody-mediated → Humoral immunity
- Type IV → Delayed → Cell mediated



	Type I (Allergy)	Type II (antibody-mediated, cytotoxic)	Type III (immune-complex)	Type IV (delayed, T cell-mediated)
Definition	Rapidly developing immune response in a previously sensitized person	Reaction of humoral antibodies that attack cell surface antigens & cause cell lysis	Results from deposition of antigen-antibody complexes on tissues	Cell-mediated, slow & prolonged response
Peak action time	15-30 minutes	15-30 minutes	Within 6 hours	After 24 hours
Mediated by	IgE antibodies	IgG or IgM antibodies	IgG, IgM antibodies	Cell-mediated

(6.02) TRANSPLANT REJECTION

University Exams

Long questions

1. Describe mechanisms of transplant rejection reactions.
2. Describe mechanisms of Hyperacute, Acute and chronic transplant rejection

Short questions

1. Types of grafts
2. Discuss Graft Versus Host Disease (GVHD).
3. Homograft rejection.
4. Transplant rejection reactions.
5. Direct and Indirect Transplant rejection reactions.

Very Short questions

1. Autograft
2. Isograft
3. Allograft
4. Xenograft

Overview

- Introduction
- Classification of Transplants →

    1. Based on genetic relationship between donor and recipient
    2. Based on anatomical site of graft

- Histocompatibility
- Types of Graft Rejection → Hyperacute, Acute, Chronic
- Mechanism of Graft Rejection → Direct pathway, Indirect pathway
- Graft versus host (GVH) reaction

Introduction

Transplantation → transfer of a graft or transplant (cells, tissues, or organs) from one site to another.

- Individual from whom transplant is taken → Donor
- Individual to whom it is transplanted → Recipient



Classification Of Transplants

1. Based on genetic relationship between donor and recipient
2. Based on anatomical site of graft

1) Based on genetic relationship between donor and recipient

- Autograft (autogenic graft): Graft from self
- Isograft (syngraft): Graft from genetically identical person, e.g. identical twin
- Allograft (homograft): Graft from genetically unrelated member of same species
- Xenograft (heterograft): Graft from different species



2) Based on anatomical site of graft

- Orthotopic grafts → When tissue or organ grafts are transplanted to their anatomically normal sites in recipient e.g. as in skin grafts
- Heterotopic grafts → Placed in anatomically abnormal sites e.g. thyroid tissue is transplanted in a subcutaneous pocket.

Histocompatibility

Histocompatible →

- If a graft and recipient tissues are histocompatible to each other (i.e. antigenically similar) → graft is accepted
- Autografts and isografts are histocompatible

Histoincompatible →

- If a graft and recipient tissues are histoincompatible (i.e. antigenically dissimilar) → graft is rejected
- Allografts and xenografts are usually histoincompatible

Types of Graft rejection

Graft rejection	Time taken for rejection	Immune mechanisms involved
Hyperacute	Minutes to hours	Preformed antibodies (Anti ABO and/or anti-HLA)
Acute	Weeks to months	Cytotoxic T cell mediated Antibody mediated
Chronic	Months to years	Chronic DTH mediated Antibody mediated

Mechanism of Graft rejection

T cells in recipient may recognize donor alloantigens in the graft in 2 different ways depending on what cells in graft are displaying these alloantigens to recipient T cells →

- Direct pathway → Allogeneic MHC molecules on graft's APCs are directly presented to recipient's helper T cells → responsible for acute graft rejections

- Indirect pathway → Donor alloantigens are processed and presented by MHC molecules present on recipient APCs to recipient's helper T cells→ responsible for chronic rejection



Graft versus host (GVH) reaction

- Graft mounts an immune response against the host (i.e. recipient) and rejects the host

(In contrary to usual situation of graft rejections in which → recipient mounts an immune response against graft antigens)

- Runt disease in animals



Conditions for GVHD

- Graft must contain immunocompetent T cells (e.g. stem cells or bone marrow or thymus transplants)
- Recipient may be immunologically suppressed and therefore cannot mount immune response against the graft.

(6.03) AUTOIMMUNITY

University Exams

Long questions

1. Define Autommunity. Describe mechanisms of Autoimmunity with examples

Short questions →

1. Immunological Tolerance
2. Mechanisms of Central tolerance and Peripheral tolerance
3. Ennumerate mechanisms of Autoimmunity
4. Sequestration of self-antigen
5. Anergy

Overview

- Introduction
- Immunological Tolerance →

    a) Central tolerance
    b) Peripheral tolerance →

        1. Anergy
        2. Phenotypic Skewing
        3. Apoptosis By AICD
        4. Regulatory T Cells (Treg Cells)
        5. Sequestration of self-antigen

- Mechanisms of Autoimmunity →

    1. Breakdown of T Cell Anergy
    2. Failure of AICD
    3. Loss of Treg cells
    4. Providing T Cell help to Stimulate Self-reacting B Cells
    5. Release of Sequestered Antigens
    6. Molecular Mimicry

- Examples of Autoimmune Diseases

Introduction

- Group of disorders in which tissue injury is caused by humoral (by auto-antibodies) or cell mediated immune response (by auto-reactive T cells) to self-antigens
- Normally → a protective mechanism called tolerance→ immune system does not attack self-antigens
- Autoimmune diseases → Any breach in tolerance mechanisms → immune system attack self-cells



Immunological Tolerance

Central Tolerance

1. In the thymus for T cells
2. In the bone marrow for B cells

Peripheral Tolerance

1. Anergy
2. Phenotypic Skewing
3. Apoptosis By AICD
4. Regulatory T Cells (T_{reg} Cells)
5. Sequestration of self-antigen



a) Central Tolerance

- Deletion of self-reactive T and B lymphocytes during their maturation in central lymphoid organs

1. In thymus → for T cells
2. In bone marrow → for B cells

b) Peripheral Tolerance

- Back-up mechanisms that occur in peripheral tissues to counteract self-reactive T cells that escape central tolerance.

1. Anergy
2. Phenotypic Skewing
3. Apoptosis By AICD
4. Regulatory T Cells (Treg Cells)
5. Sequestration of self-antigen

1. Anergy

Unresponsiveness to antigenic stimulus.

Self-reactive T cells interact
with APCs presenting self-antigen
↓
B7 molecules on APC bind to CTLA-4 molecules on
T cells instead of CD28 molecules
↓
Co-stimulatory signal is blocked
↓
No activation



2. Phenotypic Skewing

Self-reactive T cells interacting
with APCs presented with self-antigens
↓
Activation of T cells
↓
Secrete non-pathogenic cytokines and chemokine

3. Apoptosis By AICD

- Activation-induced cell death

Self-reactive T cells interacting
with APCs presented with self-antigens
↓
Activation of T cells
↓
Induces upregulation of Fas ligand
↓
Interacts with death receptor Fas
↓
Apoptosis



Apoptosis By AICD



4. Regulatory T Cells (T_{reg} Cells)

T_{reg} cells
↓
Secrete cytokines (e.g., IL-10 and TGF-β])
or killing by direct cell to cell contact
↓
Down regulate self-reactive T cells

5. Sequestration of self-antigen

- Certain self-antigens can evade immune recognition by sequestration in immunologically privileged sites
- e.g. corneal and lens proteins, testicular antigens and antigens from brain.

Mechanisms of Autoimmunity

Autoimmunity results due to breakdown of one or more of the mechanisms of immunological tolerance.

1. Breakdown of T Cell Anergy
2. Failure of AICD
3. Loss of Treg cells
4. Providing T Cell help to Stimulate Self-reacting B Cells
5. Release of Sequestered Antigens
6. Molecular Mimicry



1. Breakdown of T Cell Anergy

- In presence of tissue necrosis and local inflammation
- Eg. → Multiple Sclerosis, Rheumatoid Arthritis and Psoriasis

2. Failure of AICD

- Failure of auto reactive activated T cells to undergo activation induced cell death (AICD)
- Eg. → SLE (Systemic Lupus Erythematosus)

3. Loss of T_{reg} Cells

- Loss of regulatory T cell-mediated suppression of self-reactive lymphocytes.

4. Providing T Cell Help to Stimulate Self-Reacting B Cells

- Antibody response to self-antigens occurs only when potentially self-reactive B cells receive help from T cells.



5. Release of Sequestered Antigens

Sequestered antigens -never exposed to
tolerance mechanisms during development of immune system
↓
Injury to these organs
↓
Release of such sequestered antigens
↓
Mount an immune response

E.g. →
- Post vasectomy orchitis→ Spermatozoa release
- Post-traumatic uveitis → ocular antigens release

6. Molecular Mimicry

Microorganisms which share antigenic determinants (epitopes) with self-antigens entre human body →
Molecular Mimicry
↓
Antibodies are produced for micro-organism
↓
Antibodies cross-react with self-antigen



E.g. →
a) Shigella flexneri and HLA B27
b) Mycobacterium tuberculosis and joint membranes
c) Coxsackie B virus and myocardium
d) Streptococcus bacteria and RHD

Examples of Autoimmune Diseases

1. Graves' disease
2. Hashimoto's disease
3. Diabetes mellitus type 1
4. Goodpasture's syndrome
5. Rheumatoid fever
6. Addison's disease Ankylosing spondylitis
7. Antiphospholipid antibody syndrome (APS)
8. Aplastic anemia
9. Guillain-Barré syndrome (GBS)
10. Idiopathic thrombocytopenic purpura
11. Multiple sclerosis
12. Sjögren's syndrome

(6.04) SYSTEMIC LUPUS ERYTHEMATOSUS (SLE)

University Exams

Long Questions

1. Define SLE. Describe the etiopathogenesis, types and clinical features of systemic lupus erythematosus (SLE).
2. Define SLE. Discuss Renal manifestations in SLE

Short Questions

1. Define SLE. Discuss autoimmunity in relation to SLE.
2. Etiology and pathogenesis of systemic lupus erythemalosus.
3. Lupus erythematosus (LE) cell.
4. Clinical criteria of SLE
5. Antibodies in SLE.
6. Immunofluorescence Patterns in SLE.
7. LE cell and associated conditions.
8. Criteria for diagnosis of SLE

Overview

- Introduction
- Types
- Spectrum of autoantibodies
- Immunofluorescence Patterns
- Etio-Pathogenesis
- LE cell Phenomenon
- Clinical and pathologic manifestations
- Criteria for diagnosis of SLE

Introduction

SLE is an autoimmune disease involving multiple organs, characterized by vast array of autoantibodies, (particularly ANAs) in which injury is caused by deposition of immune complexes and binding of antibodies to various cells and tissues.

- Systemic lupus erythematosus (SLE) is a progressive chronic autoimmune disease that results in inflammation and tissue damage
- Characterized by flares, spontaneous remission, and relapses
- Highly heterogeneous
- Can affect any part of the body
    - Often damages skin, joints, heart, kidneys, lungs, nervous system

Types

1) Systemic / Disseminated form	2) Discoid form
Acute and chronic inflammatory lesions	Chronic lesion
Lesions are widely scattered in body	Localised skin lesion involving bridge of nose and adjacent cheeks without any sytemic manifestation
There is presence of various nuclear and cytoplasmic autoantibodies in plasma.	Autoantibodies may or may not present

Spectrum of Autoantibodies in SLE

- Hallmark of SLE is production of **autoantibodies**.

Specificity of Autoantibody	% positive	Association with specific disease features
ANA	95-98	Found against DNA and RNA antigen, used as screening test
ds- DNA	40-60	Nephritis, specific for SLE
U1 RNP	30-40	more often in Sjogren's syndrome
SMITH (Sm) antigen	20-30	Sm antigen is a Core protein of small ribonucleoprotein particle, and is specific for SLE
Ro (SS-A)/(SS-B) nucleoprotein	30-50	congenital heart block, neonatal lupus
Phospholipid protein complex (anti- PL)	30-40	Antiphospholipid syndrome (a thrombotic complication) in 10% of SLE patient
Multiple nuclear antigen (generic ANA)	95-100	not specific
Histone		In drug induced lupus
Anti ribosomal P antibody		in CNS lupus

Immunofluorescence Patterns



Etio-Pathogenesis

1. The exact etiology of SLE is not known.tic Predisposition/ Factors
2. Immunological factors
3. Environmental factors

UV radiations
leads to apoptosis of cells
↓
Inadequate clearance of nuclear antigen
creates increased burden of nuclear Ag in all organs
↓
B and T cells with defective tolerance, stimulated by nuclear Ag,
start to form Ab against it, even though these are self Ag
↓
Ag-Ab complexes binds to Fc portion of dendritic cell and B cell
↓
Nucleic acid component engages TLR leading
Bcell to produce more autoantibodies
Dendritic cell to produce more Ty 1 IFN
↓
stimulating more apoptosis
↓
continuing vicious cycle



Mechanisms of Injury

Type III hypersensitivity reaction:
- Most of the systemic lupus is caused by immune complexes DNA- anti- DNA complexes can be detected in the glomeruli and small blood vessels.
- T cell infiltrates are also frequently seen in kidneys.

Type II hypersensitivity reaction:
- Autoantibodies specific for red cells, white cells and platelets opsonize these cells and promote their phagocytosis and lysis

Antiphospholipid antibody syndrome (APLA):
- Patients with antiphospholipid antibodies may develop venous and arterial thromboses.
- When it is
- Associated with SLE → secondary antiphosholipid antibody syndrome
- Not associated with SLE → Primary antiphospholipid antibody syndrome

LE cell Phenomenon

Principle →
- ANAs cannot penetrate the intact cells and thus cell nuclei should be exposed to bind them with the ANAs
- Binding of exposed nucleus with ANAs results in homogeneous mass of nuclear chromatin material → LE body or haematoxylin body.



- For demonstration of LE cell phenomenon in vitro→ blood sample is traumatized to expose the nuclei of blood leucocytes to ANAs. This results in binding of denatured and damaged nucleus with ANAs. The ANA-coated denatured nucleus is chemo tactic for phagocytic cells.
- If this mass is engulfed by a neutrophil, displacing the nucleus of neutrophil to the rim of the cell→ **LE cell**
- If the mass is phago-cytosed by a monocyte → **Tart cell.**


LE Cell

Tart cell

Clinical and pathologic manifestations

Clinical Manifestation	Prevalence in patients (%)
Hematologic	100
Fatigue	80-100
Arthritis/ Arthralgia/ Myalgia	80-90
Skin	85
Fever	55-85
Weight loss	60
Renal	50-70
Pleuritis	45
Neuropsychiatric	25-35
Pericarditis	25
Raynoud phenomenon	15-40
Gastrointestinal	20
Peripheral neuropathy	15
Ocular	10



Skin

- Most commonly in sun-exposed areas
- Oral/nasopharyngeal ulcers
- Alopecia
- Butterfly rash

CVS:

- Myocarditis
- Mitral / Aortic valve stenosis or regurgitation
- Libman – Sachs endocarditis
- Coronary artherosclerosis

- Musculoskeletal
    - Polyarthralgia with morning stiffness
    - Arthritis
        - Swan neck fingers
        - Ulnar deviation
        - Subluxation with hyperlaxity of joints
- Increased risk of bone loss and fracture

- Blood vessels: Acute nerotizing vasculitis of small arteries and arterioles and may progress into fibrosis and lumen narrowing
- Spleen: Spleenomegaly, capsular thickening and follicular hyperplasia
- Lungs: Pleuritis, plueral effusion, interstital fibrosis and so pulmonary hypertenti
- CNS: Neuropschycotic syndromes ascribes to vasculitis or non-immune occlusion of small vessels

Renal → lupus nephritis

- According to the WHO, **six patterns** of renal lesions are seen →
    - Class I: Minimal lesions
    - Class II: Mesangial lupus nephritis
    - Class III: Focal segmental lupus nephritis
    - Class IV: Diffuse proliferative lupus nephritis
    - Class V: Membranous lupus nephritis
    - Class VI: Sclerosing lupus nephritis

Class I: Minimal lesions



- Light microscopy → do not show any abnormality.
- Electron microscopy and immuno fluorescence microscopy → deposits within mesangium which consist of IgG and C3.

Class II: Mesangial lupus nephritis

- Light microscopy → increase in number of mesangial cells and amount of mesangial matrix.
- Electron microscopy and immunofluorescence → granular mesangial deposits of IgG and C3





Class III: Focal segmental lupus nephritis

- Light microscopy→ focal and segmental proliferation of endothelial and mesangial cells, together with infiltration by macrophages and sometimes neutrophils.
- Electron microscopy→ Subendothelial and subepithelial deposits of IgG, often with IgM or IgA and C3, are seen.

Class IV: Diffuse proliferative lupus nephritis

- Most severe and the most common form of lupus nephritis
- Light microscopy→ Diffuse proliferation of endothelial, mesangial, and sometimes epithelial cells, involving all glomeruli.
- Electron microscopy→ large electron-dense deposits in the mesangium and in the subendothelial region



Class V: Membranous lupus nephritis

- Light microscopy→ diffuse thickening of glomerular capillary wall
- Electron microscopy→ show subendothelial deposits of immune complexes



Class VI: Sclerosing lupus nephritis

- This is chronic kidney disease of SLE, similar to chronic GN.
- Most glomeruli are sclerosed and hyalinised



Criteria for diagnosis of SLE

CLINICAL CRITERIA	IMMUNOLOGICAL CRITERIA
1. Acute Cutaneous Lupus	12. ANA
2. Chronic Cutaneous Lupus	13. Anti-dsDNA
3. Oral/Nasopharyngeal ulcers	14. Anti-Sm
4. Non-scarring alopecia	15. Anti Phospholipid Antibody
5. Synovitis involving > 2 joints	16. Low Complement
6. Serositis	17. Direct Coombs Test
7. Renal manifestations	
8. Neurological manifestations	
9. Hemolytic anemia	
10. Leucopenia/ Lymphopenia	
11. Thrombocytopenia	

Classify a patient as having SLE if →

a) The patient satisfies four of the criteria, including at least one clinical criterion and one immunologic criterion OR
b) The patient has biopsy-proven nephritis compatible with SLE and with (+) ANA or (+) anti-dsDNA antibodies.

(6.05) HIV / AIDS

University Exam

Long Questions

1. Describe Structure, Mode of Transmission, Replication, Opportunistic Infections and Lab diagnosis of HIV

Short Questions

1. Pathogenesis of human immune deficiency virus (HIV) infection
2. HIV disease and opportunistic infections
3. Oppurtunistic infections in AIDS
4. Laboratory diagnosis of AIDS.
5. High risk groups of AIDS.
6. Replication of HIV virus
7. WHO clinical staging of HIV/AIDS
8. Structural genes and non-structural genes of HIV
9. NACO Strategy for HIV Diagnosis
10. HIV Genes, Antigens and Serotyping
11. Mode of Transmission of HIV
12. CD4 count in AIDS.

Very Short Questions

1. Four common neoplasms found inpatients with HIV infection.
2. Mention 4 opportunistic infections in AIDS.
3. Enlist 4 most common Infections found in AIDS patients
4. Name the fungal infections in AIDS.
5. Enlist 4 CNS lesions found in AIDS.

Overview

- Introduction
- History
- Structure
- HIV Genes and Antigens
- HIV Serotyping
- Mode of Transmission
- Receptor Attachment
- Replication
- Natural Course
- Opportunistic Infections
- WHO clinical staging of HIV/AIDS for adults
- High risk groups
- Kinetics of Immune Response
- Lab diagnosis
- NACO Strategy for HIV Diagnosis

Introduction

- Retroviruses are group of RNA viruses that possess a unique enzyme called reverse transcriptase which directs synthesis of DNA from the viral RNA after they infect a host cell.
- Globally 1st December is observed as World AIDS Day every year.

History

- HIV in humans was believed to be acquired from chimpanzee by the cross species infections of simian counterpart of HIV (simian immunodeficiency virus or SIVcpz) in rural Africa
- First case of AIDS was described from New York (USA) in 1981
- Discovery of HIV-1 is done at Pasteur Institute, Paris in 1983

Structure

- Spherical
- 80–110 nm in size



Envelope: Present → Lipoprotein

- Lipid part: host cell membrane derived
- Protein part: 2 components:

  a) GP 120 → projected as knob like spikes on the surface
  b) GP 41 → anchoring transmembrane pedicles.

Nucleocapsid:

- Capsid is icosahedral in symmetry
- Inner core encloses:

  - RNA: 2 identical copies of single-stranded positive sense linear RNA
  - Viral enzymes → reverse transcriptase, integrase and proteases

HIV Genes and Antigens

- 3 structural genes
- 6 non-structural or regulatory genes

Structural Genes

1. Gag
2. Pol
3. Env

1) gag gene

- Codes for the core and shell of the virus.
- Expressed as a precursor protein p55 which is cleaved into three proteins:
    - P18 → constitutes the matrix or shell antigen
    - p24 } constitute the core antigens
    - p15

2) pol gene

- Codes for viral enzymes
- Expressed as a precursor protein cleaved into→
    - p31 → integrase
    - p51 → reverse transcriptase
    - p66 → protease

3) env gene

- Codes for the envelope glycoprotein
- Precursor GP → (gp 160) which is cleaved into two components:
    - gp120: main receptor of HIV that binds to CD4 molecules on host cell to initiate the infection
    - gp41: fusion protein

Non-structural Genes

Essential regulatory genes:

- tat (transcriptional transactivator)
- nef (negative factor)
- rev

Accessory regulatory genes:

- vif (viral infectivity factor)
- vpr
- vpu

HIV Serotyping

Based on sequence differences in env gene → HIV comprises of two serotypes
1. HIV - 1
2. HIV - 2

1) HIV-1

Divided into three distinct groups:
a) M → dominant group worldwide, 11 subtypes or “clades” (A-K)
b) N
c) O

Geographical distribution

- Subtype A is common in West Africa
- Subtype B is predominant in Europe, America, Japan, and Australia

- Subtype C is the most common form worldwide (47%). It is also the dominant form in Southern and Eastern Africa, India, and China

2) HIV-2

- It comprises of eight subtypes (A–H)
- Mainly confined to Africa and few other places including India.
- Group A is the most common form.



Mode of Transmission

1. Blood transfusion
2. Sexual intercourse
    - Vaginal
    - Anal
    - Oral
3. Percutaneous → Injection drug abuse, Needle stick exposure
4. Parent to child

Route of transmission	Risk of transmission (Worldwide, %)	% of total transmission	
		World	India
Blood transfusion	**>90**	**5**	**1**
Sexual intercourse	**0.1-1**	**75**	**88.9**
Vaginal	0.05-0.1	60	87.4 (heterosexual)
Anal	0.065-0.5	15	1.5 (homosexual)
Oral	0.005-0.1	Rare	Not reported
Percutaneous			
Drug abuse	0.5-1.0	10	1.6%
Needle stick exposure	0.3	0.1	1
Parent to child	13–40	10	5.4

Sexual mode

- Most common mode of transmission (75%)
- But risk of transmission is minimal (0.1–1% per coitus)

Blood transfusion

- Least common mode of transmission (5%)
- But risk of transmission is maximum (90–95%)

Percutaneous transmission

- Needle stick injury, injection drug abuse and sharing razors or tattooing
- Less effective modes of transmission

Perinatal mode:

- Risk of transmission from mother to fetus is about 20–40%
- Transmission may occur at any time during pregnancy and breastfeeding but the risk is maximum during delivery

Receptor Attachment

Main receptor:

- HIV enters into the target cells by binding its gp120 to the CD4 receptor on host cell surface.
- CD4 molecules are mainly expressed on helper T cells and also on the surface of various other cells like monocytes, macrophages, Langerhans cells, astrocytes, keratinocytes and glial cells



Co-receptor

- Necessary for fusion of HIV to gain entry into the host cell.

- Usually, the chemokine receptors act as co-receptors for HIV and act by binding to gp120.

  Examples include:

  - CXCR4 molecules present on T lymphocytes
  - CCR5 molecules present on cells of macrophage lineage

Replication

Fusion:

Following attachment of receptor and coreceptor to gp120, fusion of HIV to host cell takes place; mediated by the fusion protein gp41

Penetration and uncoating:

After fusion, HIV nucleocapsid enters into the host cell cytoplasm, which is followed by uncoating and release of two copies of ssRNA and viral enzymes

Reverse transcription:

Viral reverse transcriptase mediates transcription of its ssRNA into ssDNA so that DNA-RNA hybrid is formed.

Integration:

The viral dsDNA gets integrated into the host cell chromosome; mediated by viral integrase. The integrated virus is called as provirus

Latency:

In the integrated state, HIV establishes a latent infection for variable period. However, HIV is different from other latent viruses as it is able to replicate even in latent state and is infectious to other neighboring cells.



Natural Course

About 80–90% of HIV, infected individuals are “typical progressors,” with a median survival time of approximately 10 years.

5 stages

1. Acute HIV Disease or Acute Retroviral Syndrome
2. Asymptomatic Stage (Clinical Latency)
3. Persistent Generalized Lymphadenopathy (PGL)
4. Symptomatic HIV Infection (AIDS-related Complex)
5. AIDS

1. Acute HIV Disease or Acute Retroviral Syndrome

- Initially, HIV destroys the infected T cells and spills over into bloodstream to cause primary viremia (or acute mononucleosis-like syndrome) → initial flu-like illness
- Significant drop in the number of circulating CD4 T cells

2. Asymptomatic Stage (Clinical Latency)

Adequate immune response
develops within 1 month in most of the patients
↓
Both effective cell-mediated immune response)
and humoral immune response come into play
↓
viremia drops down
↓
CD4 T cell count becomes normal
↓
Immune response cannot clear the infection completely
↓
HIV-infected cells persist in the lymph nodes
↓
High level of ongoing viral replication
↓
This period of clinical latency is variable, may last for
10 years but ranges from few months to 30 years
↓
Once the latency is broken
↓
disease progresses rapidly
↓
Death usually occurs within 2 years if left untreated

3. Persistent Generalized Lymphadenopathy (PGL)

- 25–30% of infected people who are otherwise asymptomatic develop lymphadenopathy.
- PGL is defined as enlarged lymph nodes of more than 1 cm size in two or more non-contiguous sites that persist for at least 3 months

4. Symptomatic HIV Infection (AIDS-related Complex)

- After variable period of clinical latency → CD4 T cell level starts falling.
- Patients develop constitutional symptoms such as:
    - Unexplained diarrhea, lasting for more than 1 month
    - Weight loss (>10%), fatigue, malaise and night sweat
    - Mild opportunistic infections such as oral thrush.

5. AIDS

Gradually patient moves towards the advanced end stage of HIV infection called AIDS

AIDS is characterized by:

- Rapid fall in CD4 T cell count (usually <200 cells/µL)
- High viral load
- Lymphoid tissue is totally destroyed and replaced by fibrous tissue
- Opportunistic infections set in secondary to profound immune suppression. Depending on the CD4 T cell count, various infections occur
- Development of neoplasia (e.g. CNS lymphoma)

Opportunistic Infections

- Tuberculosis is the most common opportunistic infection that occurs in HIV-infected people
- Fungal infections → candidiasis (oral thrush) and Pneumocystis jirovecii
- Viral infections → herpes simplex mucosal lesions and CMV retinitis
- Parasitic infections → Cryptosporidium parvum diarrhea, Toxoplasma encephalitis and Strongyloides stercoralis hyperinfection syndrome

Opportunistic infections associated with HIV infection and correlation with CD4 T cell counts



WHO clinical staging of HIV/AIDS for adults

- World Health Organization, revised 2007
- Based only on clinical conditions associated with the patient.
- It classifies HIV infection into 4 stages →

Clinical Stage 1
➢ Asymptomatic HIV infection ➢ Persistent generalized lymphadenopathy
Clinical Stage 2
➢ Unexplained moderate weight loss (<10%) ➢ Recurrent respiratory tract infections (sinusitis, tonsillitis, otitis media, pharyngitis) ➢ Herpes zoster ➢ Angular cheilitis ➢ Recurrent oral ulcers ➢ Papular pruritic eruptions ➢ Seborrheic dermatitis ➢ Fungal nail infection
Clinical Stage 3
➢ Unexplained severe weight loss (>10%) ➢ Unexplained chronic diarrhea: >1 month ➢ Unexplained persistent fever: 1 month ➢ Oral candidiasis ➢ Oral hairy leukoplakia ➢ Pulmonary tuberculosis ➢ Severe bacterial infections (pneumonia, empyema, etc.) ➢ Unexplained anemia, neutropenia or chronic thrombocytopenia

Clinical Stage 4
HIV wasting syndrome (Slim disease): Characterized by profound weight loss (>10%), chronic diarrhea (>1 month), prolonged unexplained fever (1 month)
Bacterial opportunistic infections: ➢ Recurrent severe bacterial infections ➢ Extrapulmonary tuberculosis ➢ Disseminated non-tubercular mycobacterial infection ➢ Recurrent septicemia (including non-typhoidal salmonellosis)
Viral opportunistic infections: ➢ Chronic HSV infection ➢ Progressive multifocal leukoencephalopathy ➢ CMV (retinitis, or infection of other organs)
Fungal opportunistic infections: ➢ Pneumocystis jirovecii pneumonia ➢ Esophageal candidiasis ➢ Extrapulmonary cryptococcosis (meningitis) ➢ Disseminated mycoses (histoplasmosis and coccidioidomycoses)
Parasitic opportunistic infections: ➢ Toxoplasma encephalitis ➢ Chronic intestinal cystoisosporiasis (>1 month) ➢ Atypical disseminated leishmaniasis
Neoplasia: ➢ Kaposi's sarcoma ➢ Invasive cervical cancer ➢ Lymphoma (cerebral, B-cell and non-Hodgkin)

High-risk Groups

Extremely high-risk group:

- Female sex workers
- Men who have sex with men
- Transgenders
- IV drug abuser group

Moderately high-risk group:

- Health care workers (via needle pricks or splashes injury)
- Hemophiliacs and other recipients of blood products
- People with other STIs (sexually-transmitted infections).

Kinetics Of Immune Response

Entry of the virus into the body
↓
Transient high-level viremia and p24 antigenemia
↓
Window period (3 to 12 weeks)
↓

Humoral response →formation of antibodies (IgM, IgA, IgG) against structural proteins
(gag: p15, p17, p24, p55; env: gp 41, gp 120, gp 160; and pol: p31, p51 and p66)
regulatory proteins (nef, rev, tat)
accessory proteins (vif, vpu and vpr)
↓
only IgG response is consistent and long lasting

Laboratory Diagnosis

- A number of moral, ethical, legal and psychosocial issues are associated with a positive HIV status.
- The disease is life long, outcome is fatal, no cure or vaccine is available and in majority transmission is through sexual contact.
- Hence, individuals known to be HIV infected are stigmatized and develop a fear of being discriminated and socially outcasted.

3Cs while performing the test for HIV

- Consent in written format should be taken before the test is done. The patient should be explained about the nature of the test being performed
- Confidentiality of a positive test result is a must. Patient name or the word "HIV positive" should not be written on the report form
- Counseling should be provided to motivate the individual to tell the spouse/family and induce behavioral change

Specific Tests for HIV Infection:

- Screening tests (antibody detection)
- Supplemental tests (antibody detection)
- Confirmatory tests

Non–specific Immunological Methods

Screening Assays

- ELISA
- Rapid/simple test

Antigens used →

a) HIV-1 specific (p24, gp 120, gp160, gp41)
b) HIV-2 specific gp36

Should be confirmed

Supplemental Tests

1. Western Blot
2. Lineimmune assay

- Highly specific antibody detection methods
- Used for validation of positive results of screening tests
- Expensive
- Need expertise to interpret

Confirmatory tests

1. Detection of p24 Core Antigen
2. Viral RNA Detection
3. DNA PCR
4. Isolation of the Virus from Blood or Tissues

Detection of p24 Core Antigen

p24 Ag is detected by 4th generation ELISA

p24 antigen becomes detectable
after 12–26 days of infection
↓
Lasts for 3–4 weeks
↓
Again, elevated during
late advanced stage of AIDS

Uses of p24 antigen detection test:

- For confirmation of diagnosis of HIV/AIDS
- Diagnosis of HIV during the window period
- To diagnose the late stage of HIV/AIDS (immune collapse) or CNS disease
- Monitoring the progress of HIV infection

Non-specific Tests

CD4 T cell count: By flow cytometry method.

Useful for:
- Assessing the risk of opportunistic infections
- Monitoring the response to antiretroviral therapy

NACO Strategy for HIV Diagnosis

- In poor countries → it is impracticable to confirm the result of HIV screening tests by PCR or western blot as these assays are expensive and available only at limited centers.
- NACO (National AIDS Control Organization, India) has formulated a strategic plan for HIV diagnosis.

Guidelines

- Depending on the situation, for which the test is done → Positive result of the first screening test should be either considered as such or confirmed by another one or two screening tests
- 3 screening tests should use different principles or different antigens. The same kit should not be used again
- First screening test should be highly sensitive, whereas second and third screening tests should have high specificity
- Supplemental or confirmatory tests should be used only when screening test results are equivocal/intermediate.

Strategy I

- Purpose: done for transfusion and transplantation safety; i.e. for the screening of the blood donors in blood banks
- Only one test should be done.
- If found reactive, then the unit of blood is destroyed.



Strategy IIa

- Purpose: It is done for sentinel surveillance of HIV infection to estimate the prevalence of infection
- UAT: Unlinked anonymous testing (UAT)→ screening of blood specimens taken for purposes other than HIV testing. Then the samples are permanently decoded of personal identifiers. This process occurs without informed consent
- Two tests format: Positive results of the first test should be confirmed by a second test. If the second test is negative, then it is reported as negative



Strategy IIb

- Purpose: It is followed for the diagnosis of HIV/AIDS in symptomatic patients
- Positive result of the first test should be confirmed by a second test.
- If the second test is negative, then a third test is done for confirmation.



Strategy III:

- Purpose: It is done for the diagnosis of asymptomatic HIV patients, antenatal screening and screening of patients awaiting surgeries

- Three tests format: All positive results in the first test should be confirmed by the second and third test. Positive report is sent only if all three test results are found reactive
- For indeterminate results of strategy IIB and III, (i.e. first test positive but second or third test negative), a repeat test is done after 14–28 days and the sample should be sent to the reference center



(6.06) AMYLOIDOSIS

University Exams

Long questions

1. Define and classify amyloid. Describe the physical and chemical nature of amyloid. Enumerate the special stains for amyloid.
2. Define and classify amyloidosis. Explain the gross and microscopic features of organs involved in secondary amyloidosis.
3. Describe pathogenesis, morphology and staining characteristics of amyloidosis.
4. Methods of demonstration of amyloid.

Short questions

1. Define amyloid. Give classification of amyloidosis.
2. Physical and chemical nature of amyloid
3. Special stains for amyloid and the result.
4. Pathogenesis of amyloidosis.
5. Classification amyloidosis and stains used for its demonstration.
6. Spleen in amyloidosis → morphology.
7. Gross and microscopic appearance of Kidney in amyloidosis.

Very Short questions

1. Precursor proteins of amyloid.
2. Types of localized amyloidosis.
3. Types of systemic amyloidosis
4. Ennumerate Special stains for amyloid and result.
5. Role of Congo red in Amyloidosis
6. Sago spleen
7. Lordoceous spleen

Overview

- Definition
- Physical nature
- Types of Amyloidosis
- Classification →

    1. Based on cause
    2. Based on extent of amyloid deposition

- AL } Introduction
- AA } Pathogenesis

- Diagnosis
- Staining Characteristics of Amyloid
- Organs affected



Definition

- A condition in which amyloid proteins are abnormally deposited extracellularly between cells in various organs / tissues.
- Amyloid → protein that has an alteration in its secondary structure → insoluble form

Physical nature

Electron microscopy →

Nonbranching fibrils with diameter 7.5-10 nm and indefinite length

X Ray crystallography / Infrared spectroscopy→

Cross β pleated sheet conformation



Types of Amyloidosis

AL	Light chao (lambda and kappa)	Plasma cell dyscrasias (multiple myeloma)
AA	SAA (serum amyloid associated protein)	chronic inflammatory conditions
Aβ_2m	β_2 microglobulin	Hemodialysis
ATTR	Transthyretin	Familial amyloidotic polyneuropathies, Senile cardiomyopathy
Aβ	Aβ protein precursor (AβPP)	Alzheimer's disease
APrP	Prion protein	Spongiform encephalopathy
A Cal	Calcitonin	Medullary carcinoma of thyroid
A IAPP	Islet amyloid polypeptide	Type II diabetes, Insulinomas
AANF	Atrial natriuretic factor	Isolated atrial amyloid

Classification

1) Based on cause

- Primary → with unknown cause and deposition is in disease itself (AL)
- Secondary → as a complication of some underlying known disease amyloidosis (AA)

2) Based on extent of amyloid deposition

- Systemic (Generalised) → involving multiple organs
- Localised → involving 1 organ

Systemic Amyloidosis

AMYLOID PROTEIN	PRECURSOR	SYNDROME OR INVOLVED TISSUE
AL	Immunoglobulin light chain	Primary, Myeloma associated
AA	Serum AA (SAA)	Reactive, chronic inflammatory condition, Familial mediterranean fever
Aβ_2m	β_2 microglobulin	Hemodialysis
ATTR	Transthyretin	Familial amyloidotic polyneuropathies

Localised Amyloidosis

AMYLOID PROTEIN	PRECURSOR	SYNDROME OR INVOLVED TISSUE
Aβ	Aβ protein precursor (AβPP)	Alzheimer's disease
APrP	Prion protein	Spongiform encephalopathy
AIAPP	Calcitonin	Medullary carcinoma of thyroid
A Cal	Islet amyloid polypeptide	Type II diabetes, Insulinomas
AANF	Atrial natriuretic factor	Isolated atrial amyloid

AL Protein

Introduction →

- Derived from immunoglobulin light chain (lambda and kappa (k)
- Seen in plasma cell dyscrasias (multiple myeloma)
- Primary systemic amyloidosis

Pathogenesis

Stimulus → multiple myeloma, B cell lymphoma, other plasma cell dyscrasias.
↓
Monoclonal proliferation of plasma cells (Monoclonal gammopathy)
↓
Excessive production of either lambda or kappa light chain
↓
Partial degradation occurs in macrophages
↓
Non-fibrillar components like AP and GAGs causes aggregation of fibril proteins
↓
AL



AA Protein

Introduction →

- Derived from larger precursor protein in serum called SAA (serum amyloid associated protein)
- SAA is an acute phase reactant protein synthesized in liver in response to chronic inflammatory conditions
- Secondary amyloidosis

Pathogenesis

Stimulus → chronic inflammation and some tumours
↓
Activated macrophages
↓
Cytokines (IL- 1 and 6)
↓
Levels of SAA elevated from liver
↓
Partial degradation in reticuloendothelial cells
↓
Non-fibrillar components like AP and GAGs causes aggregation of fibril proteins
↓
AA

Diagnosis



- Definitive investigation for amyloidosis → Biopsy.
- Best sites for biopsy → **rectum and abdominal fat pad**
- Other sites → salivary glands, gingiva, skin, tongue, bone marrow and stomach.

Staining Characteristics of Amyloid

Stain	Appearance
1. H&E	Pink, hyaline, homogeneous
2. Methyl violet / Crystal violet	Metachromasia: rose-pink
3. Congo red	Light microscopy: pink-red Polarising light: Apple green birefringence
4. Thioflavin-T/Thioflavin-S	Ultraviolet light: fluorescence
5. Immunohistochemistry (antibody against fibril protein)	Immunoreactivity: Positive

Congo Red

- Visible light → Pink red colour
- Polarised light → Apple-green birefringence (due to cross-B-pleated sheet configuration)



Visible light Polarised light

Congo red

Organs affected

1. Kidney
2. Heart
3. Spleen
4. Liver
5. Adrenal
6. Tongue
7. Brain

1. Kidneys

- Most common form of systemic amyloidosis.
- M**ost serious** form of organ involvement.
- Earliest pathological change → **thickening of GBM**





2. Heart→

- second most common presentation
- Major organ involved in senile systemic amyloidosis



3. Spleen

- Two patterns of deposition is seen –
- Sago spleen → Amyloid deposition in **splenic follicles (White pulp)**
- Lardaceous spleen → Amyloid deposition in **splenic sinuses (Red pulp)**

4. Liver

- Amyloid deposition in **space of Disse**



5. Adrenal →

- Deposits in zona glomarulosa.

6. Tongue →

- Macroglossia

7. Brain →

- In senile plaques of Alzheimer disease



Adrenal

Tongue



Brain

(7.01) NEOPLASIA (DEFINITION, NOMENCLATURE AND PREDISPOSING CONDITIONS)

University Exams

Long questions

1. Define-Neoplasia. Write the nomenclature of Neoplasia. Ennumerate Genetic and non-hereditary predisposing conditions of Neoplasia.

Short questions

1. Define-Neoplasia.
2. Differences between carcinoma and sarcoma.
3. Precancerous conditions.
4. Definition and two examples of sarcoma.

Overview

- Definition
- Nomenclature
- Precancerous lesions →
  1. Genetic predisposition →
     a) AD inherited cancer syndromes
     b) Defective DNA-repair syndromes
     c) Familial cancers
  2. Nonhereditary Predisposing conditions →
     a) Proliferation
     b) Chronic inflammations

Definition

- Term 'neoplasia' means **new growth**
- New growth produced → **neoplasm** or **tumour**
- Branch of science dealing with the study of neoplasms → **oncology**



Neoplasm

An abnormal mass of tissue

- Growth of which exceeds and is uncoordinated with that of normal tissues
- Persists in same excessive manner even after cessation of stimuli which evoked the change.

- Neoplastic cells **lose control and regulation of replication** and form an abnormal mass of tissue.

Nomenclature

a) Benign tumors
b) Malignant tumors

a) Benign tumors

- Suffix → **oma**

Papilloma - Benign tumor with finger-like projections

Adenoma - Benign epithelial tumor arising from glands or forming a glandular pattern



Exceptions
Malignant neoplasms with suffix - Oma –

1. Melanomas
2. Seminoma
3. Mesothelioma
4. Lymphoma

2) Malignant tumors



TISSUE OF ORIGIN	BENIGN	MALIGNANT
Epithelial Tumours 1. Squamous epithelium 2. Transitional epithelium 3. Glandular epithelium 4. Hepatocytes 5. Placenta (Chorionic epithelium)	Squamous cell papilloma Transitional cell papilloma Adenoma liver cell adenoma	Squamous cell carcinoma Transitional cell carcinoma Adenocarcinoma Hepatoma (Hepatocellular carcinoma) Choriocarcinoma

6. Neuroectoderm	Hydatidiform mole Naevus	Melanoma (Melanocarcinoma)
Mesenchymal Tumours 1. Adipose tissue 2. Adult fibrous tissue 3. Embryonic fibrous tissue 4. Cartilage 5. Bone 6. Synovium 7. Smooth muscle 8. Skeletal muscle 9. Blood vessels 10. Lymph vessels 11. Meninges 12. Lymphoid tissue 13. Nerve sheath	 Lipoma Fibroma Myxoma Chondroma Osteoma Benign synovioma Leiomyoma Rhabdomyoma Haemangioma Lymphangioma Meningioma Pseudo lymphoma Neurofibroma	 Liposarcoma Fibrosarcoma Myxosarcoma Chondrosarcoma Osteosarcoma Synovial sarcoma Leiomyosarcoma Rhabdomyosarcoma Angiosarcoma Lymphangiosarcoma Invasive meningioma Malignant lymphomas Neurogenic sarcoma

Precancerous lesions



1) Genetic Predisposition

a) Autosomal dominant inherited cancer syndromes
b) Defective DNA-repair syndromes
c) Familial cancers

a) Autosomal dominant inherited cancer syndromes

Mnemonic
Very Rich, Cute, Nice Men Hereditarily Like Familiar Females

Very : **V**on Hippel Lindau (VHL) syndrome causing renal cell cancer
Rich : **R**etinoblastoma
Cute : **C**owden syndrome (PTEN gene)
Nice : **N**eurofibromatosis 1, 2
Men : **M**elanoma, **M**EN 1 and2
Hereditarily : **H**NPCC
Like : **Li**-Fraumeni syndrome
Familiar : **Fam**ilial adenomatous polyposis
Females : Ovarian and breast tumor

b) Defective DNA-repair syndromes

- Predisposes to DNA instability when exposed to environmental carcinogens.
- **Autosomal recessive.**



Mnemonic

Big FAX Hai

Big - **B**loom syndrome
F - **F**anconi's syndrome
A - **A**taxia telengectesia
X - **X**eroderma pigmentosa
Hai - **H**NPPC

c) Familial cancers

- Cancers occurring with **high frequencies in families** but without distinct pattern of hereditary transmission.
- Onset → **early with multiple or bilateral tumors.**

2) Nonhereditary Predisposing Conditions

a) Proliferation
b) Chronic inflammations

a) Proliferation

Increase risk of developing malignancy since proliferating cells accumulate genetic mutations necessary for carcinogenesis.

Example of proliferations are:

1. Regeneration, e.g. liver cirrhosis to Hepatocellular carcinoma.
2. Metaplasia
3. Hyperplasia
4. Dysplasia

b) Chronic inflammations

Induces cancers by →
1. Increasing genomic instability by producing free radicals.
2. Produces cytokines to drive cell survival and proliferation

(7.02) TERATOMA, HAMARTOMA AND CHORISTOMA

University Exams

Short questions

1. Teratoma
2. Hamartoma
3. Choristoma

Overview

- Teratoma
- Not actual tumors (neoplasms) →
    - Hamartoma
    - Choristoma

Teratoma

All Tumors are composed of cells representative of a single germ layer.

Teratoma →

- Arise from totipotent cells
- Germ cell tumors composed of multiple cell types derived from 3 germ layers → ectoderm, mesoderm and endoderm
- Contain immature or fully formed tissue including teeth, hair, bone and muscle.
- Usually benign but may be malignant

Not actual tumors (neoplasms)



Hamartoma

- Focal developmental malformation
- Mass of disorganized but mature specialized cells indigenous / native to particular site



Hamartoma of lung



Hamartoma of Spleen

Choristoma

- Ectopic islands of normal tissue.
- Heterotopia but is not a true tumor

Osteocartilagerous Choristoma of tongue



Pancreatic choristoma of Stomach

(7.03) BENIGN VERSUS MALIGNANT TUMOURS

University Exams

Long questions

1. Define neoplasm. Explain differences between benign and malignant neoplasms.
2. Difference between benign & malignant tumors.
3. Characteristics of malignant tumours.
4. Characteristics of benign tumors.
5. Define Anaplasia. Classify neoplasia. Discuss histogenesis of benign and malignant tumors.

Short questions

1. Features of anaplasia
2. Differences between benign and malignant neoplasms.
3. Describe morphology of anaplastic tumor cells.
4. Morphology of malignant cells

Overview

- Rate of growth
- Clinical features
- Gross features
- Microscopic features → Features of anaplasia →

    1. Loss of polarity
    2. Pleomorphism
    3. N:C ratio
    4. Anisonucleosis
    5. Hyperchromatism
    6. Nucleolar changes
    7. Mitotic figures
    8. Tumor giant cells
    9. Cytoplasm increased mucin
    10. DNA aneuploidy

- Spread of tumours

Rate of growth

- Benign tumour cells → Less mitotic rate (more doubling time)
- Malignant tumour cells → Increased mitotic rate (less doubling time) and slower death rate ie. cell production exceeds cell loss.

Clinical Features

Benign tumours

- Slow growing
- Asymptomatic (e.g. subcutaneous lipoma)
- May produce serious symptoms (e.g. meningioma in the CNS)

Malignant tumours

- Grow rapidly
- Ulcerate on surface
- Invade locally into deeper tissues
- Spread to distant sites (metastasis)
- Systemic features such as weight loss, anorexia and anaemia.

Gross Features

Benign tumours

- Spherical or ovoid in shape.
- Encapsulated or well-circumscribed
- Freely movable
- More firm and uniform
- **Surrounding tissue compressed**

Malignant tumours

- Irregular in shape
- Poorly-circumscribed
- Extend into adjacent tissues.
- Sarcomas → fishflesh like consistency
- Carcinomas → firm
- **Surrounding tissue invaded**

Benign Tumor



Fibroadenoma

Malignant Tumor



Invasive ductal carcinoma of breast

Microscopic Features

Lack of differentiation
OR
Presence of Anaplasia

Differentiation

Extent of morphological and functional resemblance of tumour cells to corresponding normal cells.



Anaplasia

- Lack of differentiation
- Hallmark of malignant transformation
- Irreversible

Features of anaplasia

1. Loss of polarity
2. Pleomorphism
3. N:C ratio
4. Anisonucleosis
5. Hyperchromatism
6. Nucleolar changes
7. Mitotic figures
8. Tumor giant cells
9. Cytoplasm increased mucin
10. DNA aneuploidy



1) Loss of polarity

- Normally nuclei of epithelial cells are oriented along basement membrane → basal polarity.
- Tumour cells lose their basal polarity → **nuclei tend to lie away from basement membrane**

2) Pleomorphism →

- Variation in size and shape of tumour cells.

3) N:C ratio→

Nuclei are enlarged disproportionate to cell size so that N:C ratio is increased.

- In normal cells →1:4
- Anaplastic cells. →1:1

4) Anisonucleosis →

Nuclei show variation in size and shape in malignant tumour cells

5) Hyperchromatism →

Increased nuclear material or DNA is responsible for dark staining of the cells

6) Nucleolar changes →

Malignant cells have a **prominent nucleolus** in nucleus

7) Mitotic figures →

Tumour cells show **large number of normal or abnormal mitoses**

8) Tumor giant cells →

Important feature of malignant cells

9) Cytoplasm →

Increased mucin

10) DNA aneuploidy



Spread Of Tumours

1. Local invasion (direct spread)
2. Metastasis (distant spread)



Local invasion (direct spread)



Invasiveness and metastasis → 2 most important features to distinguish malignant from benign tumours.

- Benign tumour → Neither show local invasion nor Metastasis
- Malignant tumour → Show local invasion as well as Metastasis

- Metastasis → most reliable feature of a malignant tumor
- Invasiveness → 2nd most reliable sign of malignancy (after metastasis)

FEATURE	BENIGN	MALIGNANT
I. **Clinical And Gross Features**		
1. Boundaries	Encapsulated or well-circumscribed	Poorly-circumscribed and irregular
2. Surrounding tissue	Compressed	Invaded
3. Size	Usually small	Often larger
4. Secondary changes	Occur less often	Occur more often
II. Microscopic Features		
1. Basal polarity	Retained	Lost
2. Pleomorphism	Absent	Present
3. N / C ratio	Normal	Increased
4. Anisonucleosis	Absent	Present
5. Hyperchromatism	Absent	Present
6. Mitosis	May be present but are always typical mitoses	Mitotic figures Increased and are generally atypical and abnormal
7. Tumour giant cells	May be present but without nuclear atypia	Present with nuclear atypia
8. Chromosomal abnormalities	Infrequent	Invariably present
III. Growth Rate	Slow	Rapid
IV. Local Invasion	Compresses the surrounding tissues without Invading or Infiltrating them	Infiltrates and invades the adjacent tissues
V. Metastasis	Absent	Frequently present
VI. Prognosis	Local complications	Death by local and metastatic complications

(7.04) SPREAD OF TUMOURS (METASTASIS)

University Exams

Long questions

1. Define neoplasm. Discuss their modes of spread.
2. Define neoplasm. List important features of transcoelomic spread of neoplasm.
3. Define neoplasms. Describe various methods of spread of tumors. What is metastases?
4. Write the various pathways of spread of cancer. Describe the mechanisms of invasion and metastasis of tumor.
5. Enumerate modes of spread of malignant tumor. Describe in brief mechanism of Metastasis.

Short questions

1. Spread of tumors.
2. Metastasis (routes of spread of malignant tumors with examples).
3. Lymphatic spread of tumor.
4. Hematogenous spread of cancer.
5. Transcoelomic spread of malignant tumor.
6. Metastatic cascade.
7. Virchow's node.

Overview

- Introduction
- Routes of Metastasis →

    1. Lymphatic spread
    2. Hematogenous spread
    3. Trans coelomic spread →

        a) Direct seeding into body cavities or surface
        b) Spread via CSF
        c) Implantation

- Mechanism of Metastasis → 8 Steps

Introduction

- Spread of tumour by invasion in such a way that discontinuous secondary tumour mass/masses are formed at site of lodgment away from primary site
- Most reliable feature of a malignant tumor

    - Benign tumours do not metastasize
    - All Malignant tumours can metastasize **Except** →

        1. Gliomas of CNS
        2. Basal cell carcinoma of the skin

Routes of Metastasis

1. Lymphatic spread
2. Hematogenous spread
3. Trans coelomic spread



Remember

- Carcinomas metastasize by lymphatic route
- Sarcomas metastasize by hematogenous route

1) Lymphatic Spread

- Follows natural routes of lymphatic drainage producing regional nodal metastasis



- First node in a regional lymphatic that receives lymph flow from primary tumor → **sentinel lymph node.**
- Sentinel lymph node is useful in **breast cancer, vulvar cancer and melanoma.**
- Term 'sentinel lymph node' was first used by **Gould**



- In lymph node → tumor cells lodge in first in **subcapsular sinus**





Tumor cells in subcapsular sinus

Virchow's lymph node

- **Virchow's lymph node** is nodal metastasis to **left supraclavicular lymph node** from cancers of abdominal organs e.g. cancer stomach, colon, and gallbladder.
- It is **thoracic duct end node**
- Receives afferent lymphatic drainage from left head, neck, chest, abdomen, pelvis, and bilateral lower extremities, which eventually drains into jugulo-subclavian venous junction via thoracic duct



2) Hematogenous Spread



- Common route for **sarcomas**
- **Veins** are more commonly involved than arteries because veins have **thinner walls** that can be penetrated readily (In contrast arteries have thicker walls)

With **venous invasion** → tumor cells in blood follow venous flow draining site of neoplasm.

1. All portal area drainage flows to → liver → Most common site
2. All caval blood flows to → lung → 2nd most common site



3) Transcoelomic spread

a) Direct seeding into body cavities or surface

- Occurs when a malignant neoplasm penetrates into a **natural open field**

E.g. Carcinoma of stomach seeding to both ovaries **(Krukenberg tumour)**





b) Spread via CSF

Malignant tumour of ependyma and leptomeninges
↓
Release of tumour cells into CSF
↓
Metastases at other sites in CNS



c) Implantation

- Surgeon's scalpel, needles, sutures
- Direct prolonged contact of cancer of lower lip causing its implantation to apposing upper lip

Mechanism of Metastasis

8 steps



1) Clonal proliferation and angiogenesis→

From Population of monoclonal Tumor cells
↓
A subpopulation or clone of tumour become more aggressive
↓
Involved in development of metastasis
↓
Tumour heterogeneity

2) Tumour cell loosening

- Normal cells remain glued to each other due to presence of cell adhesion molecules (CAMs) i.e. E (epithelial)-cadherin.

In cancers
↓
loss or inactivation of E-cadherin and CAMs
↓
Loosening of cancer cells

3) Tumour cell - ECM interaction

Loosened cancer cells
↓
Have receptors for ECM proteins
↓
Attached to ECM proteins (laminin, fibronectin, vitronectin, and collagen)

4) Degradation of ECM

- Tumour cells Overexpress matrix-degrading enzymes → metalloproteinases (e.g. collagenases and gelatinase)

5) Entry of tumour cells into capillary lumen

- **Autocrine motility factor (AMF)** → a cytokine derived from tumour cells which stimulates receptor-mediated motility of tumour cells.
-

6) Thrombus formation

Tumour cells
↓
covered with constituents of circulating blood
↓
Forms thrombus
↓
provides nourishment to tumour cells and protects them
from immune attack by the circulating host cells

7) Extravasation of tumour cells

- Tumour cells attach to vascular endothelium
- Extravasate to extravascular space.

8) Survival and growth of metastatic deposit

- Extravasated malignant cells on lodgment in the right environment
- Grow further under influence of **growth factors** produced tumour

(7.05) PROTO-ONCOGENES – RAS, ABL- BCR

University Exams

Long questions

1. Describe molecular pathogenesis of cancer → Oncogenes, tumor suppressor genes
2. Proto - oncogenes & Cancer.
3. Describe role of RAS gene mutation in pathogenesis of cancer.
4. ABL – BCR Translocation

Short questions

1. Enumerate steps in molecular evolution of cancer from a normal cell.
2. Define Oncogenes. Write its classification
3. Philadelphia chromosome
4. t (9; 22) translocation

Very short questions

1. What is Proto- oncogene
2. Imatinib targeted therapy

Overview

- Genetic regulators of cancer
- Proto-oncogenes
    - Introduction
    - Classification
- RAS gene
- ABL-BCR hybrid gene

Genetic regulators of cancer

Normal cell growth → 4 regulatory genes:

1. Proto-oncogenes → growth-promoting genes
2. Anti-oncogenes or tumour suppressor genes → growth-inhibiting or growth suppressor genes.
3. Apoptosis regulatory genes →control programmed cell death.
4. DNA repair genes → regulate repair of DNA damage that has occurred during mitosis





Remember

In cancer →

1. Overstimulation of proto-oncogenes
2. Inhibition of tumour-suppressor genes

Proto-oncogenes

Introduction →

- Discovered by Harold Varmus and Michael Bishop.
- Normal genes required for cell proliferation
- Act under proper physiological stimuli
- Without physiological stimuli → silent → no replication.

Proto-oncogenes
(Normal genes required for cell proliferation and differentiation)
↓ **Mutation**
Oncogenes
(Genes promoting autonomous cell growth in cancer cells)
↓
Oncoproteins
(Proteins lacking regulatory control and responsible for promoting cell growth)

Classification



RAS gene

- RAS protein → signal transducing protein
- Point mutation of RAS family genes → most common abnormality of Proto-oncogenes in human tumor.

2 scenarios

- Normal function
- Mutation

Normal function

In Inactive state RAS proteins bind GDP
↓
Growth factors
↓
RAS proteins become activated by exchanging GDP for GTP
↓
Activated RAS recruits RAF-1 and also activates PI3K
↓
Mitosis
↓
GTPase Activating Proteins (GAP)
↓
GTP breaks into GDP
↓
RAS Back to inactive state
(Activated state is transient)



Mutation

Mutation in the RAS gene
↓
GTPase Activating Proteins (GAP) not working
↓
GTPase activity cannot be increased
↓
No breakdown of GTP
↓
Permanent activation of RAS
↓
Uncontrolled mitosis
↓
Cancer



Examples

K- RAS	Colon, lung and pancreatic tumors
N- RAS	Melanoma, blood tumors (AML)
H- RAS	Bladder and kidney tumors

ABL-BCR hybrid gene

- ABL gene → normal location on chromosome 9
- BCR gene→ normal location on chromosome 22

ABL gene → normal location on chromosome 9
↓
translocated to chromosome 22
↓
Fuses with BCR gene (breakpoint cluster region)
↓
Forms an ABL-BCR hybrid gene → Philadelphia chromosome on chromosome 22
↓
Abnormal signal transduction even without growth factors
↓
Uncontrolled mitosis
↓
Cancer



Example

- CML
- Treatment of CML → Specific tyosine kinase inhibitor drug → imatinib

(7.06) TUMOUR SUPPRESSOR GENES - RB, P53

University Exams

Long questions

1. Enumerate steps of malignant transformation of a cell. Discuss role of p53 gene in neoplasia.
2. Explain briefly the role of tumor suppression genes in oncogenesis. Discuss role of p53 gene in neoplasia.

Short questions

1. Mutation in p53 oncogene.
2. Li-Fraumeni syndrome
3. List five tumor suppressor genes and cancers caused by their mutations

Overview

- Classification
- P53 gene →
    - Introduction
    - Normal function
    - Mutation
    - Li-Fraumeni syndrome

Classification

GENE	ASSOCIATED HUMAN TUMOURS
1. **RB**	Retinoblastoma, osteosarcoma
2. **p53 (TP53)**	Most human cancers, common in Ca lung, head and neck, colon, breast
3. **TGF-b and its receptor**	Ca pancreas, colon, stomach
4. **APC and b-catenin proteins**	Ca colon
5. **Others**	
i. **BRCA 1 and 2**	Ca breast, ovary
ii. **VHL**	Renal cell carcinoma
iii. **WT 1 and 2**	Wilms' tumour
iv. **NF 1 and 2**	Neurofibromatosis type 1 and 2

P53 gene

Introduction →

- P53 is a tumour suppressor gene as well as DNA repair gene
- Located on **chromosome 17**

- **Guardian of genome**
- **Molecular policeman**

Normal function

UV radiation (DNA damage)
↓
P53 accumulate in cell
↓
P21 → Cell cycle arrest
GADD 45 → DNA repair
↓
successful → MDM-3 → Degradation of p53
Unsuccessful → BAX → Apoptosis



Mutation

Cells with mutation or loss of p53
↓
DNA damage
↓
p53 dependent genes not activated
↓
No cell cycle arrest
No DNA repair
↓
Mutant cells→ Expansion and additional mutations
↓
Cancer

Li-Fraumeni syndrome

- Individual inherit one mutant p53 allele and second acquired 'hit' inactivate normal p53 allele.
- Associated with development of **sarcoma, breast cancer, leukemia and brain tumors.**

(7.07) CARCINOGENESIS (PHYSICAL, CHEMICAL, BIOLOGICAL)

University Exams

Long questions

1. Describe Physicalcal carcinogenesis in detail.
2. Define carcinogenesis. Discuss mechanism of Radiation induced cancers
3. Describe Chemical carcinogenesis in detail.
4. Enumerate major chemical carcinogens. Describe mechanisms of action of chemical carcinogens.
5. Define and classify oncogenic viruses. Explain the mechanism involved in tumor production by viruses.
6. Define carcinogenesis. Discuss role of RNA viruses in carcinogenesis.
7. Define carcinogenesis. Discuss role of DNA viruses in carcinogenesis.
8. Give examples of oncogenic viruses and discuss oncogenesis by HPV
9. Give examples of oncogenic viruses and discuss oncogenesis by EBV

Short questions

1. Role of UV rays in Radiation induced cancers
2. Role of IR rays in Radiation induced cancers
3. Steps of Chemical carcinogenesis
4. Differences between Initiators and Promotors in Chemical carcinogenesis
5. Biological carcinogens.
6. Viral carcinogenesis or Oncogenic viruses.
7. Oncogenic deoxyribonucleic acid (DNA) viruses.
8. HPV-induced carcinogenesis.
9. Enumerate oncogenic viruses with example of cancer caused by each.

Very short questions

1. Mention four radiation-induced cancers.
2. Name any four chemical carcinogens.
3. Ennumerate DNA and RNA oncogenic viruses and name cancers caused by them
4. Ennumerate oncogenic parasites and name cancers caused by them
5. List diseases caused by Epstein-Barr virus.

Overview

- Introduction
- Types →

    a) Physical carcinogens →

        1. Ultraviolet Light
        2. Ionising Radiation

    b) Chemical carcinogens →

        1. Initiation
        2. Promotion
        3. Progression

        - Tests For Chemical Carcinogenicity

c) Biologic carcinogens →

- Carcinogenic Viruses → HPV, EBV
- Carcinogenic Bacterias
- Carcinogenic Parasites

Introduction

- Carcinogenesis → Mechanism of induction of tumours
- Carcinogens → Agents which can induce tumours



Types

a) Physical carcinogens
b) Chemical carcinogens
c) Biologic carcinogens

a) Physical Carcinogenesis

Also known as Radiation induced Carcinogenesis
1. Ultraviolet Light
2. Ionising Radiation

1) Ultraviolet Light

Source → **Sunlight**

3 subtypes

1. **UV-A** → 320-400 nm
2. **UV-B** → 280- 320 nm
3. **UV-C** → 200 - 280 nm

- **UV-C** gets filtered by ozone layer
- **UV-B** is the most carcinogenic UV ray to reach the earth

(UV-B is **Bad** for humans as it causes cancers)

Mechanism →

Exposure to UV rays
↓
Pyrimidine dimers in DNA
↓
Mutation in oncogenes and tumor suppressor genes
↓
Cancer

Cancers →

1. Basal cell carcinoma (most common)
2. Squamous Cell carcinoma
3. Melanoma

2) Ionizing Radiation

Source→ x-rays, y rays, α rays, β particles

Mechanism →

Exposure to ionizing radiation
↓
dislodge ions from water
↓
Formation of highly reactive free radicals
↓
DNA damage
↓
Cancer

Cancers →

1. All forms of leukaemias (except chronic lymphocytic leukaemia CLL)
2. Cancers of the thyroid (most commonly papillary carcinoma)
3. Others →skin, breast, ovary, uterus, lung, myeloma, and salivary glands

- Maximum sensitivity is at G2 stage

	Ultraviolet Light	Ionizing Radiation
Source	Sunlight (UV-B is Carcinogenic)	X-ray, a-ray, b-ray
Cancers	Skin Cancers → - BCC - SCC - Melanoma	All leukaemia's (except CLL)→ - AML - ALL - CML Thyroid tumours
Mechanism	Pyrimidine dimers in DNA	Reactive free Radical Formation

b) Chemical Carcinogenesis

3 stages →

1. Initiation
2. Promotion
3. Progression



1) Initiation

a) Metabolic activation
b) Reactive electrophiles
c) Target molecules mutations
d) Initiated cell

2) Promotion of Carcinogenesis

- Promoter carcinogens
    - Do not damage DNA per se (not mutagenic)
    - Enhance effect of initiators
    - Cause clonal proliferation and expansion of initiated (mutated) cells

- To produce tumour → Initiator should be followed by promoter



Sequence of initiator and Promotor

3) Progression of Carcinogenesis

- Stage when mutated proliferated cell shows **phenotypic features of malignancy.**

Tests For Chemical Carcinogenicity

Ames' Test
Evaluates ability of a chemical to induce mutation in mutant strain of Salmonella typhimurium that cannot synthesize histidine.



Chemical carcinogens	Associated cancer or neoplasm
• Alkylating agents • Androgens • Aromatic amines (dyes) (aniline dyes) • Arsenic • Asbestos • Estrogen • Ethyl alcohol • Tobacco (including smokeless) • Vinyl chloride • Chromium and nickel • Benzene • Nitrates • Aflatoxin	• AML, bladder cancer • Prostate cancer • Bladder cancer • Cancer of the lung, skin • Cancer of the lung, pleura, Peritoneum • Mesothelioma • Cancer of the endometrium liver breast • Cancer of the liver, esophagus, head and neck • Cancer of the upper aerodigestive tract, bladder • Liver cancer (angiosarcoma) • Lung cancer • Leukemia, Hodgkin's lymphoma • Gastric cancer • Liver cancer

c) Biologic Carcinogenesis

Carcinogenic Organisms

	Infectious agent	Tumors
Virus	HPV	Cervical, vulvar & penile cancers (squamous cell carcinoma)
	EBV	Nasopharyngeal carcinoma, Burkitt's lymphoma, Hodgkin's lymphoma
	HBV, HCV	Hepatocellular carcinoma
	HIV	Kaposi sarcoma, NHL, squamous cell carcinoma
	HTLV-1	Adult **T-** cell leukemia
	HHV-8	Kaposi sarcoma, primary effusion lymphoma
Bacteria	H. Pylori	Gastric Cancer
Parasite	Schistosomiasis	Bladder cancer (squamous cell)
	Clonorchis sinensis	Liver cancer (HCC), bile duct carcinoma (cholangiocarcinoma), Pancreatic cancer
	Opisthorchis viverrine	Bile duct carcinoma (cholangiocarcinoma)
	Fasciola hepatica	Bile duct carcinoma (cholangiocarcinoma)

Carcinogenic Viruses

Carcinogenic Viruses

RNA viruses	DNA viruses
• Human T cell leukemia virus-1 (HTLV-1) • Hepatitis C virus (HCV)	• Hepatitis B virus (HBV) • Human herpes virus 8 (HHV8) • Human papilloma virus (HPV) • Epstein Barr virus (EBV)

Human Papillomavirus (HPV)

- Low risk HPV (16,11) → cause benign squamous papilloma (warts) in humans.
- High-risk HPVs (16,18,31,33,45)→ cause squamous cell carcinomas of the cervix, Penis, anogenital region



Epstein-Barr Virus (EBV)

- Burkitt lymphoma
- Nasopharyngeal carcinoma
- T-cell lymphoma
- Subset of Hodgkin lymphoma (lymphocyte depleted)
- Natural killer (NK) cell lymphoma

EBV
↓
Enters epithelial cells of oropharynx and B-cells via CD21 receptor
↓
LMP-1 gene present in EBV integrate with human DNA
↓
Activation of NF-kB and JAK/STAT signaling pathways
↓
promoting B-cell survival and proliferation
↓
Acquisition of t (8;14) translocation
↓
Burkitt lymphoma





Carcinogenic Bacteria

- **H. pylori** → first bacterium classified as a carcinogen.
- Causes →

    1. Gastric adenocarcinomas
    2. Gastric lymphomas

Carcinogenic Parasites

1. Schistosomiasis → Bladder cancer (squamous cell)
2. Clonorchis sinensis →Liver cancer (HCC), bile duct carcinoma (cholangiocarcinoma), Pancreatic cancer
3. Opisthorchis viverrine → Bile duct carcinoma (cholangiocarcinoma)
4. Fasciola hepatica → Bile duct carcinoma (cholangiocarcinoma)

(7.08) CLINICAL FEATURES (TUMOUR LYSIS SYNDROME AND PARANEOPLASTIC SYNDROMES)

University Exams

Long questions

1. Local and Systemic manifestations of cancer
2. Paraneoplastic syndromes with four examples.
3. Malignant cachexia.
4. Tumour lysis syndrome

Short questions

1. Systemic effects of cancer.
2. Enumerate paraneoplastic syndromes with appropriate examples.

Very Short questions

1. Cachexia.
2. List five tumors which produce paraneoplastic syndrome.

Overview

- Clinical Features
- Local Effects →

    1. Compression
    2. Obstruction
    3. Tissue destruction
    4. Infarction, ulceration, haemorrhage

- Systemic Effects →

    1. Cacer cachexia
    2. Fever
    3. Tumour lysis syndrome
    4. Paraneoplastic syndrome

Clinical Features



Local Effects

1) Compression→

Some benign tumours due to their critical location have more serious consequences e.g. pituitary adenoma

2) Mechanical obstruction →

Benign and malignant tumours in gut may produce intestinal obstruction.

3) Tissue destruction →

Malignant tumours infiltrate and destroy vital structures

4) Infarction, ulceration, hemorrhage



Systemic Effects

1. Cacer cachexia
2. Fever
3. Tumour lysis syndrome
4. Paraneoplastic syndrome

1) Cancer Cachexia

- Due to anorexia and increased nutritional demands of the tumour.
- Certain cytokines such as **tumour necrosis factor a (TNF-a), IL-1 and INF-g** play a contributory role in cachexia



2) Fever

- Tumour cells themselves elaborate **pyrogens.**



3) Tumour Lysis Syndrome

- Caused by extensive destruction of a large number of rapidly proliferating tumour cells

Tumor lysis syndrome is associated with →

1. Burkitt lymphoma
2. Acute lymphoblastic leukemia (ALL)
3. Chronic tumors (CLL)
4. Uncommonly solid tumors

Characterized by →

1. Hyperuricemia → due to increased turnover of nucleic acids → uric acid precipitation in the kidney → acute renal failure.
2. Hyperkalemia → due to release of intracellular potassium → Arrythmias
3. Hyperphosphatemia → due to release of intracellular phosphate
4. Hypocalcemia → due to complexing of calcium with elevated phosphate



TUMOR CELL LYSIS



4) Paraneoplastic Syndrome

- A group of conditions developing in patients with advanced cancer which are neither explained by direct and distant spread of the tumour
- 10 to 15% of the patients with advanced cancer develop paraneoplastic syndromes
- Sometimes PNS may be the earliest manifestation of a latent cancer.

CLINICAL SYNDROME	UNDERLYING CANCER	MECHANISM
1. **Endocrine Syndrome**		
i. Hypercalcaemia	-Lung (sq. cell Ca), kidney, breast, Adult T-cell leukaemia-lymphoma	-Parathormone-like protein -Vitamin D
ii. Cushing's syndrome	-Lung (small cell carcinoma), pancreas, neural tumours	-ACTH or ACTH-like substance
iii. Inappropriate anti-diuresis	-Lung (small cell Ca), prostate, intracranial tumour	-ADH or atrial natriuretic factor
iv. Hypoglycaemia	-Pancreas (islet cell tumour), mesothelioma, fibrosarcoma	-Insulin or insulin-like substance
v. Carcinoid syndrome	-Bronchial carcinoid tumour, carcinoma pancreas, stomach	-Serotonin, bradykinin
vi. Polycythaemia	-Kidney, liver, cerebellar haemangioma	-Erythropoietin
2. **Neuromuscular Syndromes**		
i. Myasthenia gravis	-Thymoma	-Immunologic
ii. Neuromuscular disorders	-Lung (small cell Ca), breast	-Immunologic
3. **Osseous, Joint, Soft Tissue**		
i. Hypertrophic osteoarthropathy	Lung	Not known
ii. Clubbing of fingers	Lung	Not known
4. **Haematologic Syndromes**		
i. Thrombophlebitis (Trousseau's phenomenon)	Pancreas, lung, GIT	Hypercoagulability
ii. Non-bacterial thrombotic endocarditis	Advanced cancers	Hypercoagulability
iii. Disseminated intravascular coagulation (DIC)	AML, adenocarcinoma	Chronic thrombotic phenomena
iv. Anaemia	Thymoma	Unknown
5. **Gastrointestinal Syndromes**		
i. Malabsorption	Lymphoma of small bowel	Hypoalbuminemia
6. **Renal Syndromes**		
i. Nephrotic syndrome	Advanced cancers	Renal vein thrombosis, systemic amyloidosis
7. **Cutaneous Syndromes**		
i. Acanthosis nigricans	Stomach, large bowel	Immunologic
ii. Seborrheic dermatitis	Bowel	Immunologic
iii. Exfoliative dermatitis	Lymphoma	Immunologic
8. **Amyloidosis**		
i. Primary	Multiple myeloma	Immunologic (AL protein)
ii. Secondary	Kidney, lymphoma, solid tumours	AA protein

- Most common tumor associated with paraneoplastic syndrome → small cell carcinoma of lung
- Most common paraneoplastic syndrome → Hypercalcemia (caused by parathyroid hormone related peptide by squamous cell carcinoma lung)
- Most common endocrinopathy → Cushing's syndrome (due to production of ectopic ACTH by small cell carcinoma lung)

(7.09) GRADING AND STAGING OF CANCERS

University Exams

Short questions

1. Grading and staging of cancer.
2. Grading of tumors.
3. TNM Staging
4. Differentiate Grading and staging of cancer.

Overview

- Introduction
- Grading
- Staging
- TNM Staging

Introduction

Grading' and 'staging' → 2 systems **to predict tumour behavior and guide therapy** after a malignant tumour is detected.

- Grading → gross appearance and microscopic degree of differentiation of tumour
- Staging → extent of spread of tumour within patient

Remember

- Grading → done on patholologic basis
- Staging → done on clinical grounds

Grading

Gross appearance and microscopic degree of differentiation of the tumour

Broders' grading →

- **Grade I:** Well-differentiated (less than 25% anaplastic cells)
- **Grade II:** Moderately-differentiated (25-50% anaplastic cells)
- **Grade III:** Moderately-differentiated (50-75% anaplastic cells)
- **Grade IV:** Poorly-differentiated or anaplastic (more than 75% anaplastic cells)



Staging

- Extent of spread of tumour within patient.
- Assessed by 3 ways—

    a) By clinical examination
    b) By investigations
    c) By pathologic examination of the tissue removed

2 important staging systems currently followed are:

1. TNM staging
2. AJC staging

TNM staging

3 components → **T, N and M**

- **T0 to T4**: In situ lesion to largest and most extensive primary tumour.
- **N0 to N3**: No nodal involvement to widespread lymph node involvement.
- **M0 to M2:** No metastasis to disseminated haematogenous metastases.

Tumor Size T	Tumor size < 2 cm T1	Tumor size 2-5 cm T2	Tumor size > 5 cm T3	Tumor extends to skin or chest wall T4
Lymph Nodes N	**NO** No lymph node metastasis	**N1** Metastasis to ipsilateral, movable, axillary LNs	**N2** Metastasis to ipsilateral fixed axillary, or IM LNs	**N3** Metastasis to infraclavicular/ supraclavicular LN, or to axillary and IM LNs
Metastasis M	**M0** No distant Metastasis	**M1** Distant metastasis	LNs= Lymph Nodes; IM= Internal Mammary	



Stage I Stage II Stage III Stage IV

UICC stage	TNM	Category	5-year survival (%)
I	T_1, N_O, M_0	Early cancer (operable, curable)	84
II	T_1, N_1, M_0 T_2, $N0_1$, M_0	Early cancer (operable, curable)	71
III	Any T, $N_{2\text{-}3}$, M_0 Any N, M_0	Locally advanced breast cancer (Neoadjuvant treatment to make operable)	48
IV	Any T, N, M_1	Metastatic (Not curable, Inoperable, Palliative care)	18

(7.10) LABORATORY DIAGNOSIS OF CANCER (TUMOUR MARKERS)

University Exams

Long questions

1. Laboratory diagnosis of cancer.
2. What do you understand by tumor markers? Discuss them in detail.

Short Answer

1. Discuss the role of tumor markers and their role in diagnosis.
2. Histochemistry.
3. FNAC (Fine needle aspiration cytology)

Very short questions

1. Immunohistochemistry and its role in diagnosis of tumors-list four tumors.
2. Enumerate six tumor markers giving one example of tumor associated with each.
3. Exfoliative cytology
4. Alpha feto protein.

Overview

Diagnosis Of Cancer →

1. Histological method →
    a) Paraffin embedded technique
    b) Frozen section
2. Cytological method →
    a) Exfoliative cytology
    b) FNAC
3. Histochemistry and Cytochemistry
4. Immunohistochemistry (IHC)
5. Electron microscopy
6. Tumour markers

1) Histological Methods

a) Paraffin embedded technique
b) Frozen section

a) Paraffin embedded technique

Steps →

i. Fixation
ii. Processing
iii. Embedding
iv. Cutting And Trimming
v. Staining
vi. Examination

Fixation

Processing

Embedding

Cutting And Trimming


Staining

Examination

b) Frozen section

- Procedure is generally carried out when patient is undergoing surgery
- Unfixed tissue is used
- Tissue is embedded in ice instead of paraffin wax → frozen section
- Cryostat machine is used



2) Cytological Methods

2 types →

a) Exfoliative cytology → Study of cells shed off into body cavities
b) FNAC (Fine needle aspiration cytology) → Study of cells by putting a fine needle into lesion

a) Exfoliative cytology

- **Microscopic examination of shed, desquamated cells from body surfaces** or cells harvested by rubbing or brushing a lesional tissue surface.

- Buccal mucosa
- Junction of the hard & soft palate
- Dorsum of the tongue

- Floor of the mouth and
- The lower labial region.

Exfoliative cytology is used for

1. Cervical Carcinoma
2. Endotnetrial Carcinoma
3. Bronchogenic Carcinoma
4. Bladder Carcinoma
5. Prostatic Carcinoma
6. Stomach Carcinoma

b) FNAC

- Cell morphology and nuclear features is well seen
- Cell architecture is not seen





FNA consist of four steps →

1. Palpation

2. Aspiration
3. Smear preparation
4. microscopy

	Surgical Biopsy	FNAC
Diagnosis	Histopathological	Cytopathological
Diagnostic difficulty	Narrow	Broad
Anaesthetic	Yes	No
Length of procedure	More than 5 min	Less than 5 min
Report available	2-4 days	2-4 hrs
False positive	None	Rare
Cost	Relatively high	Economical
Trauma	Yes	Little if any

3) Histochemistry and Cytochemistry

1. Basement membrane/collagen	- Periodic acid-Schiff (PAS) - Reticulin - Van Gieson - Masson's trichrome
2. Glycogen	- PAS with diastase loss
3. Glycoproteins, glycolipids, glycomucins	- PAS with diastase persistence
4. Acid mucin	- Alcian blue
5. Argyrophilic/argentaffin granules	- Silver stains
6. Nucleolar organiser regions (NORs)	- Colloidal silver stain

4) Immunohistochemistry (IHC)

TUMOUR	IMMUNOSTAIN
1. Epithelial tumours (Carcinomas)	• Pankeratin (HMW-K, LMW-K) • Epithelial membrane antigen (EMA) • Carcinoembryonic antigen (CEA) • Neuron-specific enolase (NSE)
2. Mesenchymal tumours (Sarcomas)	• Vimentin (general mesenchymal) • Desmin myogenic) • Myoglobin (for skeletal myogenic) • a-1-anti-chymotrypsin (for malignant fibrous histiocytoma) • CD34 (endothelial marker) • MB-45 (most specific)
3. Melanoma	• Vimentin • S-100
4. Lymphoma	• Leucocyte common antigen (LCA/CD45) • Pan-B (Immunoglobulins, CD20) • Pan-T (CD3) • RS cell marker for Hodgkin's (CD15, CD30)
5. Neural and neuroendocrine tumours	• Neurofilaments (NF) • NSE • GFAP (for glial tumours) • Chromogranin (for neuroendocrine)



Anticytokeratin immunoperoxidase stain of a tumor of epithelial origin (carcinoma)

5) Electron Microscopy

- Ultrastructural examination of tumour cells



6) Tumour Markers

- Some tumor produce markers that can be measured in the serum or urine or other body fluids.
- Tumor markers → biochemical indicators of presence of a tumor
- Many of these products are produced by normal body cells too
- Thus, biochemical estimation of the product in blood reflects total substance and not by the tumour cells alone.
- So, lack sensitivity as well as specificity
- Use as an adjunct to pathologic diagnosis and for prognostic and therapeutic purposes

Clinical applications

- Screening in general population
- Diagnosis in symptomatic patients
- Estimating tumor volume
- Prognostic indicator of disease progression
- Monitoring responses to therapy
- Detecting recurrence of cancer

Tumor Markers	Tumor Types
Hormones	
Human chorionic gonadotropin	Trophoblastic tumors, nonseminomatous testicular tumors
Calcitonin	Medullary carcinoma of thyroid
Catecholamine and metabolites	Pheochromocytoma and related tumors
Oncofoetal Antigens	
a-Fetoprotein	Liver cell cancer, nonseminomatous germ cell tumors of testis
Carcinoembryonic antigen	Carcinomas of the colon, pancreas, lung, stomach, and heart
Isoenzymes	
Prostatic acid phosphatase	Prostate cancer
Neuron-specific enolase	Small-cell cancer of lung, neuroblastoma
Specific Proteins	
Immunoglobulins	Multiple myeloma and other gammopathies
Prostate-specific antigen and prostate-specific membrane antigen	Prostate cancer
Mucins and Other Glycoproteins	
CA-125	Ovarian cancer
CA-19-9	Colon cancer, pancreatic cancer
CA-15-3	Breast cancer

(8.01) OBESITY

University Exams

Short questions

1. Describe etiology and pathogenesis of Obesity
2. Write Parameters for assessment of obesity
3. Body mass index (BMI)

Overview

- Introduction
- Parameters for assessment of obesity
- Etiology
- Pathogenesis
- Effects Of Obesity

Introduction

- Excess of adipose tissue that imparts health risk
- A body weight of 20% excess over ideal weight for age, sex and height is considered a health risk.

Parameters for assessment of obesity

1. BMI (Quetlets index)
2. Broca's index
3. Lorentz's formula
4. Corpulence index (HT. independent)
5. Skin fold thickness

1. BMI (Quetlets index)
   - Internationally accepted method of assessment of obesity
     Most widely used method to gauge obesity
   - BMI (Quetlets index) = $\frac{\text{Wt (Kg)}}{\text{Ht}^2\text{(m)}}$
   - A cut-off BMI value of 30 is used for obesity in both men and women.

Classification	BMI (Global)
Underweight	**<18.5**
Grade I	17.00-18.49
Grade II	16.00-16.99
Grade III	< 16
Normal range	**18.50-24.99**
Overweight	**>25.00**
Preobese	25.00-29.99
Obesity- I	30.00-34.99
Obesity- II	35.00-39.99
Obesity- III (Morbid)	>40.00

2. Broca's index = Ht - 100 = Ideal wt

3. Lorentz's formula = Ht(cm) -100 - $\frac{\text{Ht (cm)} - 150}{\text{2 (women) or 4 (men)}}$

4. Corpulence index (HT. independent) = $\frac{\text{Actual Wt}}{\text{Desirable Wt}}$ < 1.2

5. Skin fold thickness
   - **Noninvasive method** for body assessment fat
   - 4 sites – **biceps / midtriceps / Subscapular & suprailiac region**
   - Measured by – **Herpenden skin calliper**
   - Single best measurement – **Midtriceps**

     Obesity Cut – off →

     - > 18mm (boys)
     - > 32mm (girls)

Etiology

Obesity results when caloric intake exceeds utilisation.

Imbalance occur in following situations →
1. Inadequate pushing of oneself away from dining table causing overeating.
2. Insufficient pushing of oneself out of chair leading to inactivity and sedentary life style.
3. Genetic predisposition to develop obesity.
4. Diets derived from carbohydrates and fats
5. Secondary obesity → underlying diseases → hypothyroidism, Cushing's disease, insulinoma and hypothalamic disorders.

Pathogenesis



Effects Of Obesity

1. Hyperinsulinaemia
2. Type 2 diabetes mellitus
3. Hypertension
4. Hyperlipoproteinaemia
5. Atherosclerosis
6. Nonalcoholic fatty liver disease (NAFLD)
7. Cholelithiasis
8. Hypoventilation syndrome (Pickwickian syndrome)
9. Osteoarthritis
10. Cancer



Hypoventilation syndrome (Pickwickian syndrome)

Obese individuals
↓
Carbon dioxide retention and hypoxia
↓
Polycythaemia and Hypersomnolence

(Mr. Pickwick was a character, the fat boy, in Charles Dickens' Pickwick Papers → Term Pickwickian syndrome was first used by Sir William Osler for sleep apnea syndrome).

(9.01) CANCERS OF INFANCY AND CHILDHOOD

University Exams

Long questions

1. Describe etiology, Molecular pathogenesis, Clinical features, Gross and microscopy of Retinoblastoma.
2. List three common pediatric malignant tumors. Describe Molecular pathogenesis, Clinical features, Gross, microscopy of any 3 pediatric malignant tumors.

Short questions

1. List four common pediatric malignant tumors.
2. Retinoblastoma gene.
3. Knudson's two hit hypothesis.
4. Types of Retinoblastoma.
5. Molecular pathogenesis of Retinoblastoma.
6. Trilateral Retinoblastoma.
7. Gross and microscopy of Retinoblastoma.
8. Gross and microscopy of Wilms tumour.
9. Gross and microscopy of Neuroblastoma.
10. Differentiate Familial and sporadic retinoblastoma.
11. Rosettes in retinoblastoma.
12. Differentiate Neuroblastoma and Wilms tumour.

Overview

- Common Cancers of infancy and childhood
  1. Retinoblastoma
  2. Wilms tumour
  3. Neuroblastoma

  } Introduction, Arises from, Age, Gender, Molecular pathogenesis, Clinical features, Gross, Microscopy, Spread, Prognosis

Common Cancers of infancy and childhood

0-4 Years	5-9 Years	10-14 Years
Leukemia	Leukemia	
Retinoblastoma		
Neuroblastoma	Neuroblastoma	
Wilms tumor		Renal cell carcinoma
Hepatoblastoma	Hepatocellular carcinoma	Hepatocellular carcinoma
Soft tissue sarcoma (especially rhabdomyosarcoma)	Soft tissue sarcoma	Soft tissue sarcoma
	Ewing sarcoma	Ewing sarcoma
		Osteogenic sarcoma
Teratoma		
Central nervous system tumors	Central nervous system tumors	
	Lymphoma	Lymphoma, including Hodgkin lymphoma

(1) Retinoblastoma

Introduction →

Most common primary intraocular malignancy of childhood

Arises from →

Primitive retinal cells

Age →

< 3 years of age

Gender →

M = F

Molecular pathogenesis

- Mutation of RB gene
- Located on long arm (q) of **chromosome 13 (chromosome 13q14)**

Normally

Growth inhibitors	Growth factors
↓	↓
Inactivates cyclins and CDKS	Activates cyclins and CDKS
↓	↓
Hypophosphorylation of RB	Hyperphosphorylation of RB
↓	↓
RB becomes active	RB becomes inactive
↓	↓
Form an inactive complex with transcription factor EF-2	Dislocation of RB from transcription factor EF-2
↓	↓
Blocks cell division (no mitosis)	Cell division (mitosis)





Mutation in RB

Mutation in RB
↓
Permanent inactivation of RB
↓
Permanent Dislocation of RB from transcription factor EF-2
↓
Abnormal Cell division (mitosis)
↓
Cancer

Active RB gene →

- **Active** in hypophosphorylated state
- Prevents replication by forming an inactive complex with transcription factor EF-2.
- Blocks cell division
- Inhibiting the cell cycle at G1 → S phase

Inactive RB gene →

- Phosphorylation of RB gene causes **inactivation** of RB gene
- Dissociation of RB from EF-2.
- Activation of EF-2
- Cell replication

Mutation in RB →

- Permanent inactivation of RB

Knudson's two hit hypothesis

- Retinoblastoma develops when both the normal alleles of the RB genes are inactive

Types of Retinoblastoma

1. Familial / Hereditary retinoblastoma
2. Sporadic/non- hereditary retinoblastoma



1. Familial retinoblastoma
   - comprises 40% of cases
   - bilateral
   - **somatic cells inherit one mutant RB gene from a carrier parent (i.e. germline mutation). The other mutation occurs after birth.**
   - Besides retinoblastoma, children inheriting mutant RB gene have **200 times greater risk of development of other cancers eg osteosarcoma, breast, colon and lungs.**

2. Sporadic retinoblastoma
   - constitutes 60% of cases
   - unilateral
   - **acquired both the somatic mutations in the two alleles after birth.**

Heritable vs Nonheritable Retinoblastoma

	Inherited	Sporadic
Family history	Positive, but affected child may represent a new mutation	No affected relatives
Average age at presentation	8 months	24 months
Tumor distribution	Usually bilateral and multifocal	Unilateral
Increased risk for other primary tumors	Osteosarcoma, other sarcomas, melanoma, bladder cancer	None
Mutational origin	One germline and one somatic	Both somatic
Comprises	40% of cases	60% of cases

Trilateral RB

Bilateral RB with ectopic intracranial RB
(pineal gland or parasellar region



Clinical Features

1. White reflex in pupillary area (cat's eye)
2. Squint
3. Cataract/ Bulging eye/ large eye

Unilateral leukocoria

Gross

Patterns of Growth



Microscopy

- Tumour shows wide areas of necrosis and calcification

1. **Flexner-Wintersteiner rosettes** → small tumour cells arranged around a lumen with their nuclei away from the lumen
2. **Homer-Wright rosettes** → radial arrangement of tumour cells around central neurofibrillar structure



Spread



Prognosis

- **95%** survival rate
- Poor prognostic factors→

  a) Size of the tumor
  b) Optic nerve involvement
  c) Extra-ocular spread
  d) Older age at diagnosis

(2) Wilms Tumour



Introduction →

- Also known as Nephroblastoma
- Most common abdominal malignant tumour of young children

Arises from →

Primitive renal epithelial and mesenchymal components

Age →

3-5 years of age

Gender →

M = F

Molecular Pathogenesis

Mutation of **Wilms' tumour associated gene, WT1 gene** (chromosome 11p13)

Clinical Features

a) Palpable abdominal mass in a child detected by mother
b) Abdominal swelling
c) Abdominal pain

Gross

- Tumour is large, spheroidal, replacing most of the kidney.
- Solitary and unilateral (5-10% bilateral tumour)
- On cut section → variegated appearance



Microscopy

Triphasic tumour → 3 components →

1. **Blastemal** → small, round, blue, anaplastic tumour cells
2. **Epithelial** → Abortive tubules and poorly-formed glomerular structures
3. **Stromal** → Mesenchymal elements such as smooth and skeletal muscle, cartilage and bone, fat cells and fibrous tissue



Spread

- Spreads via blood → especially to lungs

Prognosis

- The prognosis of tumour with combination therapy of nephrectomy, post-operative irradiation and chemotherapy is good
- 5-year survival now is 80-90%

(3) Neuroblastoma

Introduction →

- Most common extracranial solid tumor of childhood
- Most common malignant tumor of infancy

Arise from →

Cells of neural crest that form the adrenal medulla and sympathetic ganglia

Sites →

- Adrenal gland
- Sympathetic chain
- Neck
- Thorax
- Retroperitoneum
- Pelvis

Age →

1 to 2 years of age

Gender →

M = F

Molecular pathogenesis

- Autosomal dominant pattern of inheritance

    - Amplification of the N-MYC oncogene
    - Hereditary neuroblastoma predisposition gene chromosome 16p
    - Deletion of the short arm of chromosome 1

Clinical features

a) **Constitutional symptoms**: Anorexia, weight loss, malaise, fever
b) **Pain**

    a) Most common presenting symptom
    b) Due to local spread &/or metastatic disease

c) **Abdominal lump**
d) **Respiratory compromise:** in young infants with massive hepatomegaly.
e) **Mimic pheochromocytoma:**

    a) Paroxysmal hypertension, palpitations, flushing, and headache.



Gross

- Tumour is large, soft and lobulated mass
- Extensive areas of necrosis and haemorrhages
- Cut surface → grey white



Microscopy

Tumour cells →

- Small, round and oval, slightly larger than lymphocytes
- Have scanty
- Poorly-defined cytoplasm
- Hyperchromatic nuclei

- Homer-Wright's rosettes (pseudorosettes)→ have a central fibrillar eosinophilic material surrounded by radially arranged tumour cells.

Spread

Spreads via blood → especially to bones, bone marrow, liver

Prognosis

Favorable prognostic features are →
a) Age of child below 2 years.
b) Extra-abdominal location of tumour than abdominal masses.
c) Patients in clinical stage I or stage II
d) Favourable genetic features → hyperdiploidy, lack of amplification of N-myc oncogene

Features	Retinoblastoma	Wilm's tumous	Neuroblastoma
Introduction	MC primary intraocular malignancy of childhood	Most common abdominal malignant tumour of young children	• MC extracranial solid tumor of childhood • MC malignant tumor of infancy
Arise from	Primitive retinal cells	Primitive retinal epithelial cells	Neutral crest from adrenal medulla or sympathetic ganglion
Age	<3 Yr.	3-5 Yr.	1-2 Yr.
Gender	M = F	M = F	M = F
Molecular Pathogenesis	Mutation in RB gene (Ch. 13)	Mutation in WT-1 gene (Ch. 11)	Amplification of N-Myc gene

Features	Retinoblastoma	Wilm's tumous		Neuroblastoma
C/F	White reflex, Squint, Catract, Glancoma	Abdominal mass, Pain		Abdominal mass, Pain
Gross	• Endophytic • Exophytic • Mixed	• Solitary, U/L, C/s varigated		Large lobulated, calcification
Microscopy	• FW rosettes • HW rosettes	• Triphasic	• Blastemal • Epithelial • Stromal	• HW Rosettes
Spread	• Direct • Lymphatic • Blood	• Blood		• Blood
Prognosis → 5 Yr. Survival	• 95%	• 80-90%		• 82%

(10.01) BONE MARROW ASPIRATION AND BIOPSY

University Exams

Long questions

1. Describe methods of Bone marrow examination. What is the importance of study of Bone marrow in hematological disorders.

Short questions

1. Indications and contraindications of bone marrow aspiration.
2. Differences between Bone marrow aspiration and Bone marrow Biopsy

Very Short questions

1. Bone marrow aspiration
2. Bone marrow Biopsy

Overview

- Introduction
- Haematopoiesis
- Haematopoietic organs
- Bone marrow examination
    - Preferred sites
    - Indications
    - Absolute Contraindications
- Methods→

1. Aspiration
2. Trephine biopsy

} Needles, Method, Information

- Bone marrow aspiration vs Biopsy

Introduction

Circulating blood contains 3 main types of mature blood cells—

a) RBCs (erythrocytes)
b) WBCs (leucocytes)
c) Platelets (thrombocytes)



Haematopoiesis

- Production of formed elements of blood
- RBCs, WBCs, platelets → derived from **Pluripotent Hematopoietic stem cells (HSC)**
- **CD34 positive**





Haematopoietic Organs

Age	Predominant site for erythropoiesis
2nd- 10th week of gestation (mosoblastic stage)	Yolk sac
10th-24 week of gestation (hepatic stage)	Liver
24 weeks -till birth (myeloid stage)	Bone marrow throughout skeleton
Birth to puberty	Bone marrow throughout skeleton
After puberty (after 16-18 years)	Bone marrow of vertebrae, ribs, sternum, skull, pelvis, and proximal ends of humerus & femur





Bone Marrow Examination

- Provides an invaluable diagnostic help in hematological disorders
- A peripheral blood smear examination must always precede bone marrow examination

Preferred site

a) Posterior iliac crest → In adults and children
b) Anteromedial area of tibial head → In infants
c) Anterior superior iliac crest → obese patients
d) Sternum (aspiration only)





Indications

(Mnemonic → **MM FF AA LL**)

Definitive diagnosis of→

- **M**ultiple myeloma
- **M**yelodysplastic syndrome
- **F**ever of unknown origin.
- Atypical **f**ungal and parasitic disorders such as histoplasmosis, leishmaniasis, cryptococcus, and Q fever.
- **A**nemia Unexplained
- **A**neuploidy in neonates
- **L**eukemia
- **L**ymphoma

Absolute contraindications

(Mnemonic → **BD**)

- Severe **b**leeding diatheses such as severe hemophilia
- Severe **d**isseminated intravascular coagulopathy

Methods

Bone Marrow Aspiration

Aspirate of contents of bone marrow drawn under pressure by puncturing marrow cavity.

Trephine Biopsy

Removal of a small core (1-2cm long) of BM under local anesthetic.

Aspiration

Needle →

Salah bone marrow aspiration needle



Method

- Salah bone marrow aspiration needle is used
- Suction of marrow via needle
- Smears are prepared immediately
- Smears are air-dryed
- Smears fixed in 95% methanol
- Romanowsky technique for staining
- Pearl's stail to assess reticuloendothelial stores of iron

Information

- Cellularity
- Details of developing blood cells (i.e. normoblastic or megaloblastic, myeloid, lymphoid, macrophages and megakaryocytic)
- Ratio between erythroid and myeloid cells
- Storage diseases
- Fungi (e.g. histoplasmosis) and parasites (e.g. malaria, leishmaniasis, trypanosomiasis).



Trephine Biopsy

Needle →

Jamshidi trephine needle



Method

a) Performed by a Jamshidi trephine needle
b) A core of tissue from periosteum to bone marrow cavity is obtained
c) Fixed in formalin
d) Decalcified
e) Paraffin block is prepared
f) Cutting with microtome
g) Stained with haematoxylin and eosin



Information

a) Overall marrow architecture
b) Cellularity
c) Presence or absence of infiltrates

- But is less valuable than aspiration for individual cell morphology

Bone marrow aspiration vs Biopsy

Feature	Aspiration	Trephine Biopsy
1. Site	Sternum, posterior iliac crest	Posterior iliac crest
2. Instrument	Salah BM aspiration needle	Jamshidi trephine needle
3. Stains	Romanowsky, Perl's' reaction for iron	Haematoxylin and eosin
4. Time	Within 1-2 hours	Within 1-7 days
5. Morphology	Better cellular morphology of aspiration smears but marrow architecture is indistinct	Better marrow architectural pattern but cell morphology is indistinct

(10.02) RETICULOCYTES

University Exams

Short questions

1. What is reticulocyte? Mention its morphology and staining method.
2. Significance of reticulocytosis and reticulocytopenia

Very Short questions

1. Reticulocyte-normal and abnormal values.
2. Reticulocyte count- utility.
3. Four conditions where reticulocyte increases.
4. Stains for Reticulocytes.

Overview

- Erythropoiesis
- Introduction
- Stains of Reticulocytes
- Reticulocyte count
- Reticulocytosis versus Reticulocytopenia

Erythropoiesis





Introduction

- Juvenile red cells devoid of nuclei
- Slightly larger than RBCs
- Polychromatic (having a blue color)
- Contain ribosomal RNA



Stains of Reticulocytes

Supravital stains →

1. New methylene blue
2. Brilliant cresyl blue

} Deep blue reticulofilamentous material





Reticulocyte count

- Percentage of reticulocytes among red cells present in peripheral blood
- **Indicator of erythropoeitic activity of bone marrow**

Normal Values →

- Infants → 2-6%
- Adults and children → 0.5-2.5%
- Absolute reticulocyte count → 25,000-75,000/mL

Reticulocytosis versus Reticulocytopenia

Reticulocytosis	Reticulocytopenia
Increased RBC production	Decreased RBC production
Reticulocyte Count > 1.5%	Reticulocyte Count < 0.5%
Conditions • Acute blood loss or hemorrhage • Postsplenectomy • Microangiopathic Anemia • Autoimmune Hemolytic Anemia • Hemoglobinopathy (Sickle cell anemia and Thalassemia) • Post anemia Treatment like Folate Supplementation, Iron Supplementation and vitamin B_{12} Supplementation.	**Conditions** • Aplastic anemia • Bone marrow infiltration • Bone marrow suppression (Sepsis/Chemotherapy radiotherapy) • Blood transfusion • Liver disease • Disordered RBC maturation (Iron, B_{12}, Folate Deficiency, Hypothyroidism, Sideroblastic Anemia or Anemia of Chronic Disease)

(10.03) RBC INDICES AND ESR

University Exams

Short questions

1. Mean corpuscular volume (MCV)
2. Mean corpuscular haemoglobin (MCH)
3. Mean corpuscular haemoglobin concentration (MCHC)
4. Red cell distribution width (RDW)
5. Hematocrit or Packed cell volume (PCV)
6. Factors determining value of ESR.
7. Principle and methods of determining ESR.

Very Short questions

1. Normal values of RBC indices.
2. Ennumerate diseases causing Raised ESR
3. Ennumerate diseases causing Low ESR

Overview

- Red Cell Indices→

1. Mean corpuscular volume (MCV)
    1. Mean corpuscular haemoglobin (MCH)
    2. Mean corpuscular haemoglobin concentration (MCHC)
    3. Red cell distribution width (RDW)
    4. Hematocrit or Packed cell volume (PCV)

- Erythrocyte Sedimentation Rate (ESR) →
    - Definition
    - Principle
    - Factors
    - Phases
    - Methods
    - Diseases with abnormal ESR

Red Cell Indices

1. Mean corpuscular volume (MCV)
2. Mean corpuscular haemoglobin (MCH)
3. Mean corpuscular haemoglobin concentration (MCHC)
4. Red cell distribution width (RDW)
5. Hematocrit or Packed cell volume (PCV)

1) Mean cell volume (MCV)

- Average volume of red blood cell.
- Normal value → **82-96 fL (femtolitres)**

$$\text{MCV} = \frac{\text{PCV in L/L}}{\text{RBC Count/L}}$$

- Increased in → megaloblastic anemia
- Decreased in →iron deficiency anemia

2) Mean cell hemoglobin (MCH)

- Hemoglobin content per RBC.
- Normal value → **27-33 Pg**

$$\text{MCH} = \frac{\text{Hb/L}}{\text{RBC Count/L}}$$

-
- Decreased in → **microcytic hypochromic anemia.**
- Increased in → **megaloblastic anemia.**

3) Mean cell hemoglobin concentration (MCHC)

- Average hemoglobin concentration in a given volume of packed RBC.
- Normal value is **33-37 gm/Dl**

$$\text{MCHC} = \frac{\text{Hb/L}}{\text{PCV in L/L}}$$

-
- Decreased in →**microcytic hypochromic anemia.**
- Increased in → **hereditary spherocytosis.**
- Remains normal in → **megaloblastic anemia.**

4) Red blood cell distribution width (RDW)

- Degree of anisocytosis (size variation of RBCs).
- Normal value is **11.5 to 14.5.**

- Increased in → **megaloblastic anemia, iron deficiency anemia and immune hemolytic anemia.**
- Remains normal in → **thalassemia.**



5) Hematocrit or packed cell volume (PCV)

- Blood volume proportion occupied by RBC
- Normal values →
  - 45% for men
  - 42%for women
- Increased in → Increase in the number or size of RBC
- Decreased in → Decrease in the number or size of RBC



HEMATOCRIT

Red Cell Indices

a) MCV = $\frac{\text{PCV in L/L}}{\text{RBC count/L}}$ Normal value = 82-96 fL

b) MCH = $\frac{\text{Hb/L}}{\text{RBC count/L}}$ Normal range = 27-33 Pg.

c) MCHC = $\frac{\text{Hb/dl}}{\text{PCV in L/L}}$ Normal value = 33-37 gm/dL

d) RDW = Degree of Anisocytosis Normal value = 11.5 to 14.5.
e) PCV = Blood volume proportion occupied by RBC
Normal value = 45% for men and 42%for women

Erythrocyte Sedimentation Rate (ESR)

Definition→

ESR is used as an index for presence of an active disease which could be due to many causes.

Principle →

When well-mixed anticoagulated blood is placed in a vertical tube
↓
Erythrocytes tend to fall towards the bottom of the tube/pipette
↓
Form a packed column in the lower part of the tube in a given time
↓
Rate at which they fall is ESR



Factors



1. Rouleaux formation → When a number of erythrocytes aggregate in form of rouleaux and settle down → their area is much less than that of sum of area of constituent corpuscles.
2. Concentration of fibrinogen → causes increased viscosity of plasma. Concentration of fibrinogen parallels ESR.
3. Concentration of globulins → decreases negative charge of RBCs that tends to keep them apart → promoting rouleaux formation → retards ESR. Conditions in which albumin is low, ESR is higher.
4. Length of the tube → If length of tube is more → RBCs will have to travel a longer distance → ESR is lower than when length of tube is short and vice versa.
5. Bore of the tube → If bore of tube is more → negative charge which keeps the RBCs apart will be less → ESR will be more and vice versa.
6. Position of the tube → If tube or pipette is not vertical → RBCs will have to travel less distance → ESR will be more.

Phases

- ESR by all methods is expressed as mm first hour
- 3 phases occur in sequence within 1 hour →

1. Phase of rouleaux formation → In initial 10 minutes→ process of rouleaux formation occurs
    1. Phase of settling → In next 40 minutes→ settling of RBCs occurs at a constant rate.
    2. Phase of packing → In last 10 minutes→ sedimentation slows and packing of RBCs to bottom occurs and it continues slowly later too.



Methods

1. Westergren's method
2. Wintrobe's method
3. Micro ESR method
4. Automated methods

1) Westergren's Method

- Open at both ends
- Straight pipette 30 cm long
- Internal bore diameter of 2.5 mm
- Calibrated from 0-200 mm from top to bottom

Normal values →
- Males → 3-5 mm 1st hour
- Females → 4-7 mm 1st hour



Westereren's pipette

Advantages →
a) More sensitive method.
b) Easy to fill and clean Westergren's pipette

Disadvantages →
a) Requires more amount of blood.
b) Dilution of blood in anticoagulant affects ESR.
c) Filling of blood by mouth pipetting should be strictly discouraged.

2) Wintrobe's Method

- Glass tube closed at one end.
- 110mm long
- Internal bore diameter of 2.5 mm.
- Graduated on both sides: from 0 to 10 cm on one side and 10 to 0 cm on the other side

Normal values →

- Males → 0-7 mm 1st hour
- Females → 0-15 mm 1st hour

Advantages →
1. Requires small amount of blood.
2. No dilution with anticoagulant,
3. PCV can also be done by same tube.

Disadvantages →
1. Because of short column and choice of anticoagulant → it is not as sensitive as Westergren's method.
2. Addition of more anticoagulant can lower ESR.
3. ESR of more than 100 mm cannot be measured.



Wintrobe's tube & pasteur pipette

Diseases with abnormal ESR

Diseases Causing Raised ESR	Diseases Causing Low ESR
1. Tuberculosis 2. SABE 3. Acute myocardial infarction 4. Rheumatoid arthritis 5. Shock 6. Anaemias 7. Chronic liver diseases 8. Multiple myeloma 9. Pregnancy 10. Ankylosing spondylitis	1. Polycythaemia 2. Spherocytosis 3. Sickle cell anaemia 4. Congestive heart failure 5. New-born and infant • Hypofibrinogenemia

(10.04) ANEMIA → DEFINITION, CLASSIFICATION, C/F

University Exams

Long questions

1. Define anemia. Write etiological and morphological classification of anemias
2. Define anemia and write its Clinical features. Enumerate various causes of macrocytic anemia.
3. Define anemia. Classify anemia according to mechanisms of its production.

Short questions

1. Etiological classification of anemias
2. Morphological classification of anemias
3. Clinical features of anemia

Very Short questions

1. Anisocytosis
2. Poikilocytosis
3. Enumerate various causes of microcytic anemia.
4. Enumerate various causes of macrocytic anemia.
5. Enumerate various causes of normocytic anemia.

Overview

- Definition
- Morphology of RBC
- Variations in Size
- Variations in Colour
- Abnormal shapes of RBC
- Classification of Anaemias
- Clinical Features

Definition

- Anaemia is defined as reduced haemoglobin concentration in blood below the lower limit of the normal range for the age and sex of the individual.
- The lower extreme of the normal haemoglobin is taken as →
    - 13.6 g/dl for males
    - 12.0 g/dl for females
    - 15 g/dl for newborns



Morphology of RBC

- Shape→ biconcave (due to spectrin)
- Size → diameter of 7 to 8 um.
  Biconcave shape is

- Colour → Hb. is located peripherally, leaving an area of central pallor equal to 30-35% of diameter of the cells (Central 1/3rd pallor)

Variations in Size

- RBC of normal size → **Normocytic**
- When red cell diameter is greater than 9 µm they are referred as **macrocytes**
- When red cell diameter is less than 6 µm they are referred as **microcytes**
- **On smear size of RBC is compared with small lymphocyte**


Normocytes


Microcytes


Macrocytes

Variations in Colour

- RBC with normal hemoglobin content (color) (Central 1/3rd pallor) → **Normochromic**
- When area of central pollar is greater than 50% of diameter, it is referred as **hypochromic**
- Variation is size of RBCs is know as **anisocytosis**
- Variation in shape of RBCs is know as **poikilocytosis**


Normochromic


Hypochromic


Anisocytosis


Poikilocytosis

Abnormal shapes of RBC

Target cells		• Thalassemia, • Post splenectomy
Bite cells		• G6PD deficiency
Sickle cell		• Sickle cell anemia
Teardrop cell		• Myelofibrosis
Spurr cell (Acanthocyte)		• Abetalipoproteinemia
Burr cell (Echinocyte)		• Uremia
Rouleaux formation		• Multiple myeloma
RBC agglutination		• Cold autoimmune haemolytic anemia • Paroxysmal cold hemoglobinuria

Classification of Anaemias

1. Morhological Classification
2. Pathophysiological Classification

1) Morphologic Classification

1. Microcytic, hypochromic
2. Normocytic, normochromic
3. Macrocytic, normochromic



Microcytic/hypochromic



Normocytic/Normochromic



Macrocytic/Normochromic

2) Pathophysiological/ Etiological Classification







Clinical Features



Symptoms →

1. Tiredness
2. Easy fatiguability
3. Generalised muscular weakness
4. Lethargy and headache.
5. In older patients → symptoms of cardiac failure, angina pectoris, intermittent claudication, confusion and visual disturbances

Signs →

1. Pallor

    - Most common and characteristic sign
    - Seen in mucous membranes, conjunctivae and skin.

2. Cardiovascular system

    - A hyperdynamic circulation may be present with tachycardia, collapsing pulse, cardiomegaly, midsystolic flow murmur, dyspnoea on exertion,congestive heart failure.

3. Central nervous system

    - Attacks of faintness, giddiness, headache, tinnitus, drowsiness, numbness and tingling sensations of the hands and feet.

NO PALLOR

PALLOR IN ANEMIA

NO PALLOR

PALLOR IN ANEMIA

PROMINENT CREASES ON THE PALM

PALE PALMAR CREASES IN ANEMIA

4. Ocular manifestations
    - Retinal haemorrhages may occur if there is associated vascular disease or bleeding diathesis.
5. Reproductive system
    - Menstrual disturbances such as amenorrhoea and menorrhagia and loss of libido
6. Gastrointestinal system
    - Anorexia, flatulence, nausea, constipation and weight loss may occur.

(10.05) HEMOLYTIC ANEMIAS → DEFINITION, CLASSIFICATION, LAB DIAGNOSIS

University Exams

Long questions

1. Classify heamolytic anemias. Discuss laboratory investigations for diagnosis of heamolytic anemia.
2. Write Characteristic features of haemolytic anaemia. Discuss approach to laboratory diagnosis in a case of heamolytic anemias.
3. Discuss broad mechanisms of hemolytic anemia.

Short questions

1. Laboratory tests in hemolytic anemias.
2. Peripheral smear blood picture of hemolytic anemia.
3. How do you classify hemolytic anemias?
4. Intravascular vs Extravascular hemolysis
5. Characteristic features of haemolytic anaemia

Very Short questions

1. Ennumerate examples of Intravascular and Extravascular hemolysis

Overview

- Introduction
- Characteristic features
- Types
  - Intravascular haemolysis
  - Extravascular haemolysis
- Extravascular versus Intravascular Haemolysis
- Classification
- Lab Diagnosis

Introduction

Haemolytic anaemias are defined as anaemias resulting from an increase in the rate of red cell destruction



Characteristic features

1. A shortened red cell life span → premature destruction of red cells.
2. Inceased erythropoiesis in the marrow → to compensate the loss of red cells → increased reticulocyte count.
3. Accumulation of products of hemoglobin catabolism → Increased bilirubin □ Jaundice

Types

2 Types→

1. Intravascular haemolysis
2. Extravascular haemolysis





1) Intravascular haemolysis

- Less common
- RBC's undergo lysis in the circulation
- Release their contents into plasma
- Plasma haemoglobin rises → haemoglobinemia
- Part of it excreted in the urine → haemoglobinuria
- Some of the hemoglobin is reabsorbed in kidney and is converted to hemosiderin which is excreted → hemosiderinuria
- Free hemoglobin is oxidized to methemoglobin → Methemoglobinemia
- Free hemoglobin in plasma bound to haptoglobin and this complex is rapidly cleared by mononuclear phagocytic system -> Decreased heptoglobin
- During hemolysis LDH released in blood →Increased serum LDH
- Excess heme catabolized to bilirubin within mononuclear phagocytic system → Mild Increase indirect bilirubin and jaundice and gall stones



2) Extravascular haemolysis

- More common
- RBCs are taken up by cells of RE system of spleen (major site), liver or bone marrow where they are destroyed and digested
- Plasma haemoglobin level is not raised.

- Hemoglobinemia
- Hemoglobinuria
- Hemosiderinuria
- Methemoglobinemia

} All absent

- Because little hemoglobin escapes into plasma → heptoglobin does not decrease
- LDH → mildly raised
- Jaundice (due conversion of heme into indirect bilirubin)
- Increased urinary urobilinogen
- Increased fecal stercobilinogen
- Splenomegaly and Hepatomegaly



Extravascular versus Intravascular Haemolysis

Features	Extravascular Haemolysis	Intravascular Haemolysis
1. Frequency	More common	Less common
2. Location	RE system organs (spleen, bone marrow, liver)	In peripheral blood
3. Iron stores	Increased	Decreased
4. Serum ferritin	Normal to high	Low
5. Plasma haemoglobin	Absent	Present
6. Methaemoglobin	Absent	Present
7. Haemoglobinuria	Absent	Present
8. Haemosiderinuria	Absent	Present
9. Unconjugated hyperbilirubinaemia	Markedly elevated	Mildly elevated
10. Serum LDH	Mildly elevated	Markedly elevated

Classification



Lab Diagnosis

1) Tests of Increased Red Cell Breakdown

1. Serum bilirubin → raised
2. Urine urobilinogen → raised
3. Faecal stercobilinogen → raised

4. Serum haptoglobin → reduced
5. LDH → raised
6. Evidences of intravascular haemolysis in the form of → haemoglobinaemia, methaemoglobinaemia, haemoglobinuria, and haemosiderinuria.

2) Tests of Increased Red Cell Production

1. Reticulocytosis → most useful initial test of marrow erythroid hyperplasia.
2. Bone marrow shows erythroid hyperplasia
3. X-ray of bones shows evidence of expansion of marrow space especially in tubular bones and skull.



Reticulocytosis


Erythroid Hyperplasia


X-Ray

3) Tests for Shortened Red Cell Lifespan

1. Tested by **51Cr** labelling method.

4) Morphology of RBC

Red cell morphology		Hemolytic
	Polychromasia	
	Reticulocyte (supra-vital stain)	
	Spherocyte	Hereditary spherocytosis, Autoimmune hemolytic Anemia
	Elliptocyte	Hereditary elliptocytosis
	Stomatocyte	Liver disease
	Sickle cell	Sickle cell anemia
	Fragments	Microangiopathy, HUS, TTP, Cardiac valve, DIC
	Blister cell	G6PD deficiency
	Spur cell	Severe liver disease

(10.06) HEREDITARY SPHEROCYTOSIS

University Exams

Long questions

1. Mention two conditions where you get spherocytes in the peripheral smear. Discuss the etiology and diagnosis of hereditary spherocytosis.
2. Describe pathogenesis and laboratory diagnosis of hereditary spherocytosis.

Short questions

1. Osmotic fragility test
2. Discuss molecular pathology and morphology of hereditary spherocytosis.

Very Short questions

1. Draw peripheral smear diagram of hereditary spherocytosis.
2. Why MCHC is increased in hereditary spherocytosis.

Overview

- Introduction
- Pathogenesis
- Lab diagnosis

Introduction

- Autosomal dominant inheritance
- RBC membrane is abnormal

Pathogenesis

1. **Spectrin** →The chief protein component responsible for **biconcave shape**
2. **Ankyrin and band 4.2** →Binds spectrin to band 3
3. **Band 3** →A transmembrane ion transport protein.
4. **Band 4.1** →Binds spectrin to glycophorin A, a transmembrane protein.



Spectrin → 2 subunits a and β

- This spectrin is attached to cell membrane at two sites→
    1. Spectrin binds to ion transporter, **band 3** of membrane with **ankyrin and band 4.2**
    2. Spectrin binds to **glycophorin A** of the membrane by **protein 4.1 and actin**
- Mutation most commonly involves the gene coding for → **Ankyrin > Band-3 > Spectrin >Band 4.2 (also called paladin)**



Remember

- Most common defect in hereditary Spherocytosis is in Ankyrin.
- Most common defect in hereditary elliptocytosis is in spectrin.

Mutation
↓
Loss of membrane cytoskeleton proteins (ankyrin, spectrin, Band 3, 4.2)
↓
Reduced membrane stability
↓
Loss of membrane relative to cytoplasm
↓
Cells become microspherocytes (smallest possible diameter for a given volume)
↓
Trapped in to spleen (Due to reduced deformability)
↓
RBC are phagocytosed by RE cells
↓
Extravascular hemolysis



Lab diagnosis

1. Extravascular hemolysis
2. Morphology of RBC

    - Microspherocytes



3. Increase in MCHC → due to dehydration caused by loss of K+ and water.
    - Hereditary spherocytosis is the only important anemia in which MCHC is increased.
4. Special test
    1. Osmotic fragility test / Pink test
        - RBCs lyses more readily in solutions of low salt concentration i.e. osmotic fragility is increased
    2. Direct Coombs test is negative so as to distinguish it from acquired spherocytosis of AIHA in which it is positive.



(10.07) G6PD DEFICIENCY ANEMIA

University Exams

Long questions

1. Describe pathogenesis, types and laboratory diagnosis of G6PD deficiency Anemia

Short questions

1. G6PD deficiency state.
2. Heinz bodies
3. Bite cells

Overview

- Introduction
- Pathogenesis
- Types
- Lab diagnosis

Introduction

- X-linked recessive disorder
- Characterized by hemolytic anemia on exposure to oxidative stress

Oxidant stress →

a) Viral and bacterial infections
b) Certain drugs (antimalarials Primaquine, sulfonamides, nitrofurantoin, aspirin, vitamin K)
c) Metabolic acidosis
d) Ingestion of fava beans (favism)

Pathogenesis

- Normally RBCs are protected from oxidant injury by **reduced glutathione**
- Reduced glutathione is generated from oxidized glutathione and reducing equivalent for this reaction is provided by **NADPH.**
- This NADPH is generated in HMP shunt by enzyme **Glucose-6-phosphate dehydrogenase**



Normal person
↓
Oxidant stress
↓
G6PD present
↓
NADPH is generated in HMP shunt
↓
Oxidized glutathione is converted to Reduced glutathione
↓
Reduced glutathione protects RBCs from oxidant injury
↓
No hemolysis

Person with G6PD deficiency
↓
Oxidant stress
↓
NADPH is not generated in HMP shunt
Oxidized glutathione is not converted to Reduced glutathione
RBCs are not protected from oxidant injury
↓
Intravascular and extravascular hemolysis

Types

1. **Type-1** (mildest form)
2. **Type-2** (moderately sever)
3. **Type-3** (most severe form)

Lab diagnosis

a) Intravascular and extravascular hemolysis
b) Morphology of RBC → Heinz bodies, Bite cells

Morphology

- Heinz bodies
    - Precipitate of denatured Hb. in RBC due to oxidative stress)
    - Revealed by supravital staining.
- Bite cells (eaten RBCs by splenic macrophages)






Heinz bodies on
supravital stain
Bite cell

(10.08) PAROXYSMAL NOCTURAL HEMATURIA (PNH)

University Exams

Long questions

1. Describe pathogenesis, clinical features and laboratory diagnosis of Paroxysmal Noctural Haemoglobinuria

Short questions

1. Screening test for Paroxysmal Noctural Haemoglobinuria
2. Confirmatory test for Paroxysmal Noctural Haemoglobinuria

Very Short questions

1. Ham test
2. Triad of Paroxysmal Noctural Haemoglobinuria

Overview

- Introduction
- Pathogenesis
- Clinical features → Triad
- Lab diagnosis →
    - Screening test
    - Confirmatory test

Introduction

- Caused by an acquired intrinsic defect in the cell membrane.
- Results from acquired mutation that inhibits the synthesis of Glycosylphospatidylinositol (GPI)

Pathogenesis

- 2 anchoring proteins called complement regulating proteins →
    a) Decay accelerating factor (DAF, CD55)
    b) A membrane inhibitor of reactive lysis (MIRL, CD59)





These proteins are anchored to cell membrane of blood cells by **GPI**
CD55 and CD 59
↓
Anchored to cell membrane of blood cells by GPI
↓
Prevent activation of complement (During sleep)
↓
Prevent complement mediated lysis of blood cells
↓
No hemolysis



Mutation of PIG-A (phosphatidyl inositol glycan) gene
↓
Synthesis of glycosyl phosphatidyl inositol (GPI) does not occur
↓
Absence of complement regulating proteins (CD55, CD59) on blood cells
↓
Blood cells sensitive to complement mediated lysis (during sleep)
↓
Intravascular hamolysis

Clinical features → Triad

a) Hemolysis →

- Paroxysmal (intermittent attacks)
- Nocturnal (occurs in the night) → Because during sleep the pH of blood gets slightly reduced and acidic medium leads to activation of the complement.

b) Pancytopenia

- Because causative somatic mutations occur in pluripotent stem cells → all cells are sensitive to complement mediated lysis → pancytopenia → Anemia, Leucopenia, Thrombocytopenia

c) Thrombosis → Due to absence of CD-59 on platelets → externilization of phosphotidylserine → thrombosis

Lab diagnosis

a) Screening test
b) Confirmatory test

1) Screening test

1. Ham test → Acidic pH will activate complement pathway for RBC lysis.



2. Sucrose lysis test → Sucrose will reduce pH and this will activate complement pathway for hemolysis.



2) Confirmatory test

Flow cytometry →

- Bimodal distribution of the red cells



(10.09) AUTOIMMUNE HEMOLYTIC ANEMIA (AIHA)

University Exams

Long questions

1. Describe Types, pathogenesis and Causes of Autoimmune Haemolytic Anaemia (AIHA)

Short questions

1. Coomb's Test.
2. Pathogenesis of Warm Autoimmune Haemolytic Anaemia
3. Pathogenesis of Cold Autoimmune Haemolytic Anaemia

Very Short questions

1. Donath-Landsteiner antibody

Overview

- Introduction
- Types →
  - Warm AIHA
  - Cold AIHA

  } Introduction, Pathogenesis, causes
- Lab diagnosis

Introduction

- Caused by autoantibodies directed against a red cell antigen

Types

- Warm Antibody AIHA
- Cold Antibody AIHA

1) Warm Antibody AIHA

Introduction →

- **Most common** form of AIHA
- Antibody react with RBC at **37°C.**
- Most causative antibodies are of the **IgG** class
- **Extravascular hemolysis** occur.

Pathogenesis





Causes

1. Autoimmune disorders → SLE, RA, Ulcerative colitis, Scleroderma, Antiphospholipid antibodies.
2. Chronic lymphocytic leukemia (CLL)
3. Hodgkin disease
4. Non-Hodgkin lymphoma
5. Multiple myeloma
6. Thymoma
7. Drugs → Methyldopa, penicillin, Cephalosporine, quinidine.
8. PAN

2) Cold Antibody AIHA

2 conditions:

1. Cold agglutinin disease
2. Paroxysmal Cold haemoglobinuria (PCH)

a) Cold agglutinin disease

Introduction →

- Antibodies react with RBC at 0° to 4°C.
- Caused by cold agglutinin IgM antibodies.
- Extravascular hemolysis occur.

Pathogenesis

IgM antibody directed against the I antigen of RBC
↓
bind to I antigen of RBC at 4 degree C
↓
Fix complement on RBC (transient reaction)
↓
C3b acts as an opsonin
↓
It enhances the phagocytosis of RBCs in phagocytic system of liver and spleen
↓
Extravascular hemolysis



b) Paroxysmal cold haemoglobinuria (PCH)

Introduction →

- Antibodies react with RBC at 0° to 4°C
- Autoantibodies are of IgG class→ also known as Donath-Landsteiner antibody
- Intravascular hemolysis occur.

Pathogenesis

IgG (Donath-Landsteiner) antibody directed against the P antigen of RBC

↓

bind to P antigen of RBC at 4 degree C

↓

Antigen antibody complex on the red blood cell activates complement

↓

Large amount of membrane attack complex (MAC)forms

↓

Destroys red cells directly in blood vessel

↓

Intravascular hemolysis



Causes

- Syphilis
- Mycoplasma
- Mumps
- Measles
- Flu syndrome

	Warm AIHA	Cold AIHA CAD	Cold AIHA PCH
Temperature	37^0C	4^0C	4^0C
Ab.	IgG	IgM	IgG (Donath-Landsteiner)
Hemolysis	Extravasular	Extravasular	Intravascular

Lab diagnosis

1. Intravascular haemolysis as well as Extravascular haemolysis
2. Morphology

    - **Prominent spherocytosis** in the peripheral blood film.



3. Special test

    - Positive direct and indirect Coombs test



(10.10) SICKLE CELL ANEMIA

University Exams

Long questions

1. Give the classification of anemias. Discuss pathogenesis and laboratory diagnosis of sickle cell disease.
2. Classify hemolytic anemias. Describe sickle cell anemia.

Short questions

1. Clinical features of Sickle cell anemia
2. Pathogenesis of sickle cell disease.
3. Factors determining rate of sickling
4. Lab. investigation in sickle cell disease.
5. Why Sickle cell anemia gives Protection against falciparum malaria
6. Qualitative and Quantitative Haemoglobinopathies

Very Short questions

1. Diagnosis of sickle cell anemia.
2. HbS
3. Enumerate tests for sickling in sickle cell anemia.
4. Principle of Sickling/sickle cell test
5. Screening of sickle cell anemia
6. Hb. Electrophoresis in sickle cell anemia

Overview

- Haemoglobin types
- Haemoglobinopathies
- Sickle Cell Anaemia→
    - Introduction
    - Pathogenesis
    - Consequences of Sickling
    - Factors determining rate of sickling
    - Clinical features
    - Protection against falciparum malaria
    - Screening
    - Diagnosis

Haemoglobin types



Haemoglobinopathies

2 types:

- ✓ Qualitative disorders → there is structural abnormality in synthesis of haemoglobin e.g. sickle cell syndrome
- ✓ Quantitative disorders → there is quantitatively decreased globin chain synthesis of haemoglobin e.g. thalassaemias

Sickle Cell Anaemia

Introduction →

- Autosomal recessive disorder
- Homozygous state of HbS in RBCs (SS)

Sickle Cell Anaemia
↓
Presence of sickle haemoglobin (HbS) in the RBCs (Qualitative disorders)
↓
RBCs with HbS develop sickling when they are exposed to low oxygen tension

Pathogenesis

Haemoglobin
↓
β-globin chain gene mutation (chromosome11)
↓
Single point mutation at sixth position of β-globin chain (mis-sense mutation)
↓
Substution of a valine residue for a glutamic acid residue
↓
Sickle hemoglobin (HbS) instead of HbA



Consequences of Sickling

1. Vaso occlusion of microcirculation
2. Haemolytic anaemia
3. Increased MCHC

Deoxygenation of HbS-containing RBC
↓
Polymerisation of deoxygenated HbS
↓
Elongated rod-like polymers align
↓
Distort RBC into sickle shape (Sickling)
↓
Express higher levels of adhesion molecules (abnormally sticky)
↓
Vaso occlusion of microcirculation

With repeated episodes of deoxygenation and sickling
↓
Membrane damages permanently
↓
RBC become irreversibly sickled
↓
Difficulty in passing through splenic sinusoids → Spleenic Sequestration
↓
Rapid phagocytosis
↓
Extravascular Haemolysis → Haemolytic anaemia

With membrane damage
↓
Water comes out of the cell
↓
Intracellular dehydration
↓
Increased MCHC





Factors determining rate of sickling

1. Amount of HbS and interaction with other Hb

    - **In homozygous individuals**→ all Hb. is HbS → sickling.
    - **In heterozygous**→ only 40% of Hb. is HbS → Sickling does not occur because HbA (remaining 60%) has an inhibitory effect on polymerization of HbS
    - **Fetal hemoglobin HbF** also has inhibitory effect on sickling of HbS → Newborns do not manifest until 6 months of age

2. Hemoglobin concentration of RBC→ Greater MCHC→ greater is sickling
3. Intracellular dehydration →increases the MCHC → faciliatates sickling
4. Decrease in Ph → Acidosis→Reduces the oxygen affinity of Hb. → Increasing the fraction of deoxygenated HbS→ faciliatates sickling
5. Length of time RBCs exposed to low oxygen tension→ Organs having slow or sluggish circulation (bone and spleen) have an increased chance of sickling

Clinical Features

1. Anaemia →

    - Irreversible sickle cells have difficulty in passing the splenic sinusoids→ sequestration→ rapid phagocytosis.
    - Associated with Jaundice and reticulocytosis.

2. Vaso-occlusive phenomena →

    - Reversible sickle cells → express higher levels of adhesion molecules → abnormally sticky → occlusion of microcirculation
    - Most common manifestations of sickle cell anemia.

A) Bone

a) Dactylitis or inflammation of bones of hands and feet → Hand foot syndrome
b) Fish mouth appearance of vertebra → due to vaso occlusive crisis of vertebral arteries.
c) Prominent cheek bones
d) Crew cut appearance of skull → due to marrow expansion causing bone resorption



B) Lungs →

- Acute chest syndrome characterized by cough, fever and chest pain

C) Brain →

- Seizures or stroke

D) Skin →

- Leg ulcers

E) Penis →

- Stagnation in corpora cavernosa leads to priapism

F) Aplastic crises →

- Occurs from the infection of red cell progenitors by parvovirus B19 → sudden worsening of the anemia

G) Spleen

- Initial stages → **splenomegaly** due to congestion and trapping of RBCs in sinusoids
- Prolonged hypoxia and infarction → **Autosplenectomy** → increases susceptibility to infection with capsulated organisms like Hemophilus influenzae, Pneumococcus, etc.

Protection against falciparum malaria

Patients with HbS are relatively protected against falciparum malaria

Malarial parasites consume O2
↓
Decrease intracellular pH
↓
Promote sickling
↓
Distorted RBCs cleared rapidly by phagocytes in spleen
↓
Keep parasite loads down

Screening

Sickling Test



Diagnosis

1) Peripheal smear

a) Sickle cells
b) Target cells
c) Features of splenic atrophy such as presence of Howell-Jolly bodies



2) ESR

- Increased in all anemia except in sickle cell anemia where **ESR is decreased** because there is no rouleaux formation.

3) Haemoglobin electrophoresis

- HbS moves Slowly towards Anode.
- HbA moves faster towards Anode

Interpretation→
- Formation of 1 band → sickle cell anemia
- Formation of 2 bands → sickle cell carrier or trait.



(10.11) THALASSEMIA

University Exams

Long questions

1. Classify heamolytic anemia. Give an account of thalassemia.
2. Classify thalassemia syndromes. Write a note on molecular pathogenesis of β Thalassaemia and describe the blood picture in β Thalassaemia major.
3. β -thalassemia – Pathogenesis, Types, Clinical features, Lab diagnosis
4. α Thalassaemia – Pathogenesis, Types, Clinical features, Lab diagnosis

Short questions

1. Discuss hematological findings in thalassemia major
2. β-thalassemia - Lab diagnosis
3. α Thalassaemia - Types and Clinical features

Very Short questions

1. Classification of thalassemia.
2. Peripheral blood picture in thalassemia.

Overview

- Introduction
- Classification

α Thalassaemia→

- Definition
- Molecular pathogenesis
- Types
- Clinical features

β Thalassaemia →

- Definition
- Molecular pathogenesis
- Types
- Clinical features
- Lab diagnosis

Introduction

- Quantitative abnormalities of globin chain synthesis
- Reduced synthesis of one or more of the globin polypeptide chains.

Classification

1. **α** Thalassaemia→
   Absent or reduced synthesis of **α globin** chain **(chromosome 16)** with normal β -chain synthesis.
2. β Thalassaemia→
   Absent or reduced synthesis of **β globin** chain **(chromosome 11)** with normal α -chain synthesis



1) α thalassemia

Definition →

- Absent or reduced production of α globin chain
- Reduced formation of HbA in RBCs

Molecular pathogenesis

- Deletion of one or more of α -chain genes located on short arm of chromosome 16.

Types

TYPE	Hb	Electrophoresis	Genotype	Clinical Syndrome
1. Hydrops foetalis	3-10 gm/dl	Hb Barts	Deletion of four α-genes	Fatal in utero or in early infancy
2. HbH disease	2-12 gm/dl	HbF, HbH	Deletion of three α-genes	Haemolytic anaemia
3. α-Thalassaemia trait	10-14 gm/dl	Almost normal	Deletion of two α-genes	Microcytic hypochromic blood picture but no anaemia
4. α-Thalassaemia Carrier	Normal	Normal	Deletion of One α-genes	Asymptomatic





Clinical features

- α –thalassaemia carrier → Clinically asymptomatic
- α-thalassaemia trait → Minimal or no anemia
- HbH→ HbH has extremely high affinity for oxygen → tissue hypoxia → moderately severe anemia→ occasional blood transfusion.
- Hydrops fetalis (Most dangerous) → In fetus→ excess y-globin chains form tetramers → Hb barts → high affinity for oxygen → Severe tissue anoxia → Intrauterine fetal death.

2) β thalassemia

Definition →

- Absent or reduced production of β globin chain
- Reduced formation of HbA in RBCs.

Molecular pathogenesis

Point Mutations of β-globin gene resulting from single base changes

3 types of mutations →

a) Splicing mutations (most common)
b) Chain terminator mutations
c) Promoter region mutations

Types

TYPE	HB	Electrophoresis	Genotype	Clinical Syndrome
1. β-Thal major	<5 gm/dl	HbA (0-50%), HbF(50-98%)	β^{thal}/ β^{thal}	Severe anaemia, requires transfusions
2. β-Thal intermedia	5-10 gm/dl	Variable	Multiple mechanisms	Severe anaemia, but regular transfusions not required
3. β-Thala minor	10-12 gm/dl	HbA2(4-9%), HbF (1-5%)	β^{A}/ β^{thal}	Usually, asymptomatic



Clinical features

1. Anaemia → starts within first 4-6 months of life
2. Hepatosplenomegaly → due to extramedullary haematopoiesis
3. Expansion of bones → due to marked erythroid hyperplasia leading to thalassaemic facies and malocclusion of the jaw.
4. Iron overload → due to repeated blood transfusions → damage to the endocrine organs (slow rate of growth, delayed puberty, diabetes mellitus) and damage to the liver and heart.

Lab Diagnosis

1) Peripheral smear



Microcytic Anisocytosis Poikilocytosis Target cells Nucleated RBls

hypochromic RBC



Basophilic Stippling
Target Cells
Poikilocytosis

2) Osmotic fragility test

Decreased osmotic fragility → Increased resistance to saline haemolysis



3) NESTROF Test

Naked Eye Single Tube Red cell Osmotic Fragility (NESTROF) test

- blood samples (1 of control and 1 of patient) are added to 2 tubes with 0.35% saline.
- After 30 min a white paper with a black line is placed behind both tubes.
- RBCs in control sample undergo hemolysis → black line is visible
- RBCs in thalassemia resistant→ black line is not clearly visible.



4) Haemoglobin electrophoresis

- Almost complete absence of HbA
- Since β chains are not produced but y chains are synthesized normally → increased amounts of HbF (90%).
- Increased amount of HbA2

5) Skull X-ray

Widening of the diploe gives rise to →

- Crew cut appearance on skull X-ray
- Hair on end appearance



(10.12) IRON DEFICIENCY ANEMIA (IDA)

University Exams

Long questions

1. Define anemia. Give etiologic classification of anemias. Describe peripheral blood smear and bone marrow picture and list laboratory investigations in iron deficiency anemia.
2. Describe the etiology, clinical features and the laboratory diagnosis of iron deficiency anemia.

Short questions

1. Laboratory diagnosis of iron deficiency anemia (peripheral blood smear and bone marrow findings).
2. Blood picture in iron deficiency anemia.
3. Role of Hepcidine in Iron regulation
4. Clinical features of iron deficiency anemia.

Very Short questions

1. Peripheral blood smear findings in iron deficiency anemia
2. Ennumerate Factors of Iron absorption
3. Earliest Response to therapy in iron deficiency anemia

Overview

- Introduction
- Normal iron metabolism
- Regulation of iron absorption
- Factors of Iron absorption
- Etiology
- Clinical features
- Lab diagnosis
- Differential diagnosis
- Treatment
- Response to therapy

Introduction

Most common cause of anemia worldwide and in India



Normal iron metabolism

Iron in diet is in Fe3+ (ferric) state
↓
Reduced to Fe2+ (ferrous) iron in gastric acid
↓
Transported across apical membrane by divalent metal transporter 1 (DMT1)
↓
Transported from cytoplasm across basolateral membrane by Ferroportin
↓
coupled to oxidation of Fe2+ iron to Fe3+ iron
↓
Fe3+ iron binds to plasma protein transferrin
↓
Transferrin delivers iron to red cell progenitors in marrow
↓
Extra iron is stored in liver in form of ferritin



Remember

- Iron is transported is blood in combination with a glycoprotein transferrin.
- Iron is stored as ferritin or haemosiderin

Regulation of iron absorption

Increases in intrahepatic iron levels
↓
Hepcidin is synthesized and released from liver
↓
Hepcidin binds to ferroportin
↓
Degrade it
↓
Inhibits iron transfer from enterocyte to plasma
↓
Iron becomes trapped within duodenal cells
↓
Iron is lost as these cells are sloughed

When body is replete with iron
↓
High hepcidin levels
↓
Inhibit iron absorption into blood

When body iron stores is low
↓
Hepcidin synthesis falls
↓
Facilitates iron absorption into the blood

Factors of Iron absorption

Factors increasing absorption	Factors decreasing absorption
• Ferrous form (Fe2+) • Acid (HCI) in the stomach • Ascorbic acid • Amino acid and sugars in the food • Physiological conditions (pregnancy and hypoxia)	• Ferric form (Fe3+) • Achlorhydria (absence of HCI secretion) • Alkaline food (pancreatic secretions) • Phytates, tannates and phosphates in diet • Tetracyclines and EDTA • Inflammatory disorders

Etiology

Dietary lack	Impaired absorption	Increased requirement	Chronic blood loss
• Infants • Children • Low socioeconomic status • Elderly	• Steatorrhea • Sprue • Chronic diarrhea • Gastrectomy	• Growing infants and children • Pregnant females • Premenopausal women	• GIT (peptic ulcer, gastric cancer, hemorrhoids, hookworm disease) • Urinary tract (renal, pelvic or bladder cancers) • Genital tract (uterine cancer, menorrhagia)

Clinical Features

1) Anaemia

- Weakness
- Fatigue
- Dyspnoea on exertion
- Palpitations
- Pallor of the skin, mucous membranes and sclerae.
- Older patients may develop angina and congestive cardiac failure.

2) Epithelial tissue changes

a) Nails (koilonychias or spoon-shaped nails)
b) Tongue (atrophic glossitis)
c) Mouth (angular stomatitis)

Koilonychia



Normal tongue | Atrophic glossitis | Angular stomatitis

3) Plummer-Vinson syndrome



Lab Diagnosis



1) Blood picture

a) Hb. concentration → Fall
b) RBCs

- Hypochromic
- Microcytic
- Anisocytosis
- Poikilocytosis
- Target cells
- Ring / pessary cells




Microcytic hypochromic RBCs


Anisocytosis and Poikilocytosis


Target Cells

c) Reticulocyte count → slightly low
d) Indices

- Diminished MCV
- Diminished MCH
- Diminished MCHC

e) Leucocytes → TLC and DLC normal
f) Platelets → normal

2) Bone marrow findings

a) Marrow cellularity

- Increased due to erythroid hyperplasia
- Myeloid-erythroid ratio decreased

b) Erythropoiesis

- Small normoblasts (micronormoblast)
- Cytoplasmic maturation lags behind nuclear maturation

c) Marrow iron (Prussian blue reaction) → Deficient



Prussian blue stain (Iron Stores)



Normal — Iron deficiency Anemia

3) Biochemical findings

a) Serum iron level → Low (50 µg/dl)
b) Serum ferritin level → Low (Indicating poor tissue iron stores)
c) Total iron binding capacity (TIBC) → High
d) Transferrin saturation → Low (below 15%)



SERUM IRON BOUND TO TRANSFERRIN — UNSATURATED SERUM IRON BINDING CAPACITY

Differential Diagnosis

Test	Iron Deficiency	Chronic Disorders	Thalassaemia Minor	Sideroblastic Anaemia
1. MCV, MCH, MCHC	Reduced	Low normal-to-reduced	Very low	Very low (except MCV raised in acquired type)
2. Serum iron	Reduced	Reduced	Normal	Raised
3. Serum ferritin	Reduced	Raised	Normal	Raised (complete saturation)
4. TIBC	Raised	Low-to-normal	Normal	Normal
5. Marrow-iron stores	Absent	Present	High	High
6. Iron in normoblasts	Absent	Absent	Present	Ring sideroblasts
7. Hb electrophoresis	Normal	Normal	Abnormal	Normal

Treatment

a) Correction of disorder causing the anemia
b) Correction of iron deficiency →

1. Oral therapy →

    a) Ferrous sulfate
    b) Ferrous fumarate
    c) Ferrous gluconatev

2. Parenteral therapy → Indications →

    a) Intolerance to oral iron therapy
    b) Malabsorption
    c) Post-operative cases
    d) Cases requiring a rapid replenishment of iron stores e.g. in women with severe anaemia a few weeks before expected date of delivery.

- Common preparation is iron dextran → single intramuscular injection or intravenous infusion

Response to therapy

Earliest hematological evidence of improvement → Increase in reticulocytes

(10.13) SIDEROBLASTIC ANEMIA

University Exams

Short questions

1. Define and classify anemia. Write Pathogenesis and Laboratory diagnosis of Sideroblastic Anaemias

Very Short questions

1. Ring Sideroblast
2. Types of Sideroblast

Overview

- Introduction
- Pathogenesis
- Types of Sideroblast
- Types of Sideroblastic Anaemias
- Lab diagnosis
- Differential diagnosis
- Treatment

Introduction

Body has adequate amount of iron but is unable to corporate it into hemoglobin.

Pathogenesis

Mitochondrial Enzymes defect activity in erythroblasts
↓
Iron enters into mitochondria
↓
Cannot be utilized to synthesize heme
↓
Iron accumulates in mitochrondria of erythroblasts
↓
Ringed sideroblast

Types of Sideroblast

- Type 1 sideroblasts: < 5 siderotic granules in cytoplasm.
- Type 2 sideroblasts: ≥ 5 siderotic granules, but not in perinuclear distribution.
- Type 3 or ringed sideroblasts: ≥ 5 siderotic granules in perinuclear distribution.





Types of Sideroblastic Anaemias

a) Hereditary sideroblastic anaemia → AX-linked disorder → defective enzyme activity of aminolevulinic acid (ALA) synthetase required for haem synthesis.
b) Acquired sideroblastic anaemia →

- Previous chemotherapy
- Irradiation
- Myelodysplasia
- Myelproliferative disorders
- Drugs→ Alcohol, isoanizid, choramphenicol, pyridoxine deficiency, lead poisoning

Lab Diagnosis



1) Blood picture

a) Haemoglobin → Fall
b) RBC → Hypochromic anaemia which may be microcytic or normocytic (dimorphic).
c) Reticulocyte count → normal but may be slightly low
d) Indices → (MCV, MCH and MCHC) → Reduced in hereditary type but MCV is often raised in acquired type.
e) Leucocytes → TLC and DLC normal
f) Platelets → normal

2) Bone marrow findings

a) Marrow cellularity

- Increased due to erythroid hyperplasia
- Myeloid-erythroid ratio decreased

b) Erythropoisis →

- Erythroid hyperplasia

c) Marrow iron

- Raised
- Ring sideroblasts



3) Biochemical Findings

a) Serum iron level → Raised
b) Serum ferritin level → Raised
c) Total iron binding capacity (TIBC) → normal 33%
d) Transferrin saturation → Increased

Differential Diagnosis

- Iron deficiency Anemia
- Thalassaemia
- Anemia of Chronic disorders

Treatment

1. Removal of the offending agent.
2. No definite treatment is available for hereditary types
3. Pyridoxine → administered routinely to all cases of sideroblastic anaemia
4. Blood transfusions

(10.14) ANEMIA OF CHRONIC DISEASE (AOCD)

University Exams

Long questions

1. Define and classify anemia. Write Pathogenesis and Laboratory diagnosis of Anaemia of chronic disorders (AOCD)

Short questions

1. Pathogenesis of Anaemia of chronic disorders
2. Describe investigations in Anaemia of chronic disorders
3. Causes of Anaemia of chronic disorders

Overview

- Introduction
- Etiology
- Pathogenesis
- Lab diagnosis
- Differential diagnosis
- Treatment

Introduction

- In patients of chronic systemic diseases in which anaemia develops secondary to a disease process but there is no actual invasion of the bone marrow.
- Severity of anaemia is usually **directly related** to the primary disease process.
- Anaemia is **corrected only if the primary disease is alleviated**

Etiology

1. Anaemia in chronic infections/inflammation

   a) Infections e.g. tuberculosis, lung abscess, pneumonia, osteomyelitis, subacute bacterial endocarditis, pyelonephritis.
   b) Non-infectious inflammations e.g. rheumatoid arthritis, SLE, vasculitis, dermatomyositis, scleroderma, sarcoidosis, Crohn's disease.
   c) Disseminated malignancies e.g. Hodgkin's disease, disseminated carcinomas and sarcomas.

2. Anaemia of renal disease e.g. uraemia, renal failure
3. Anaemia of hypometabolic state e.g. endocrinopathies, protein malnutrition, scurvy, pregnancy, liver disease.

Pathogenesis





Lab Diagnosis



1) Blood picture

a) Haemoglobin → Fall
b) RBC → Normocytic normochromic
c) Reticulocyte count → normal but may be slightly low
d) Indices → MCHC is slightly low.
e) Leucocytes → TLC and DLC normal
f) Platelets → normal

2) Bone marrow findings

a) Marrow cellularity

- Increased due to erythroid hyperplasia
- Myeloid-erythroid ratio decreased

b) Erythropoisis → Normal erythroid maturation
c) Marrow iron → Raised

3) Biochemical Findings

a) Serum iron level → Reduced
b) Serum ferritin level → Increased
c) Total iron binding capacity (TIBC) → Normal or decreases
d) Transferrin saturation → Decrease

4) Others

a) Acute phase reactants (gamma-globulin, C3, haptoglobin, a1-antitrypsin and fibrinogen) → Raised
b) ESR → Raised

Differential Diagnosis

a) Iron deficiency Anemia
b) Thalassaemia
c) Sideroblastic Anemia

Treatment

- Treat underlying cause

(10.15) MEGALOBLASTIC ANEMIA

University Exams

Long questions

1. Define and classify anemia. Write Pathogenesis and Laboratory diagnosis of megaloblastic anemia.
2. Enumerate causes of megaloblastic anemia. Describe its peripheral blood and bone marrow findings.

Short questions

1. Laboratory diagnosis of megaloblastic anemia (peripheral blood smear and bone marrow findings).
2. Megaloblast.
3. Describe investigations in macrocytic anemia.
4. Describe PS and BM findings in Megaloblastic anemia.
5. Draw a well labelled diagram of peripheral smear in megaloblastic anemia.
6. What is schilling test?

Very Short questions

1. Causes of megaloblastic anemia.
2. Effects of vitamin B12 deficiency on the nervous system.

Overview

- Introduction
- Etiology
- Pathogenesis
- Clinical features
- Lab diagnosis
- Treatment

Introduction

- Impaired DNA synthesis
- Vit B12 and folic acid are required for DNA synthesis.
- So megaloblastic anaemia is due to deficiency of vit. B12 and folic acid

Etiology

Vitamin B12 deficiency

1. Decrease intake: Inadequate diet, alcoholism, infancy.
2. Impaired absorption: Malabsorption state, intestinal diseases.
3. Increased requirement: Pregnancy, infancy, disseminated cancer.

Folate deficiency

1. Decrease intake: Inadequate diet, alcoholism, infancy.
2. Impaired absorption: Malabsorption state, intestinal diseases.
3. Increased requirement: Pregnancy, infancy, disseminated cancer.

Pathogenesis

Inadequate DNA synthesis
↓
Defective nuclear maturation
↓
Maturation of nucleus lag behind to that of cytoplasm
↓
Nuclear/Cytoplasmic asynchrony
↓
Formation of megaloblasts and macrocytes





- Vit. B12 is also involved in conversion of **methylmalonyl CoA to succinyl CoA** which is required for the formation of **normal neuronal lipids.**
- So, deficiency of vitamin B12 (but not of folic acid) results in **neurological features**

Clinical Features

1. Anaemia → In both vit. B12 and folate def.
2. Neurologic manifestations → only in vit. B12 def. Not in folate def.

    - Subacute combined degeneration of the spinal cord
    - Peripheral neuropathy
    - Numbness, paraesthesia, weakness, ataxia, poor finger coordination and diminished reflexes.

Lab Diagnosis



1) Blood picture

a) Haemoglobin concentration → Fall
b) RBCs

- Macrocytosis
- Macroovalocytes
- Anisocytosis
- Poikilocytosis
- Tear drop cells
- Basophilic stippling
- Cabott Ring
- Howell-jolly bodies







c) Reticulocyte count → Low to normal

- Absolute values
- Elevated MCV
- Elevated MCH

d) Normal or reduced MCHC
e) Leucocytes

- Hypersegmented neutrophils (having more than 5 nuclear lobes) → First manifestation of megaloblastic anemia

f) Platelets → Bizarre forms



Hyper segmented neutrophils
(6 lobes)



2) Bone marrow findings

a) Marrow cellularity

- Hypercellular
- Erythroid hyperplasia
- Decreased myeloid-erythroid ratio

b) Erythropoiesis

- Megaloblastic erythropoiesis
- Nuclear maturation lags behind that of cytoplasm

c) Marrow iron →Increase



Special Tests for for vit. B12 deficiency

Schilling test

- Radioisotope used for labeling B12 is either 58Co or 57Co.



1. 1 mg of unlabelled vitamin B12 **('cold' B12)** is given by **intramuscular route** (to saturate binding sites)
2. 1 mg of radioactively labelled vitamin B12 **('hot' B12)** is given by **oral dose**
3. The patient is kept fasting for a further period of 2 hours
4. **Urinary excretion of B12** is estimated

Interpretation→

- In normal individuals → 24-hour urinary excretion is >10% of the oral dose of 'hot' B12
- Patients with B12 deficiency → 24-hour urinary excretion is < 10% of 'hot' B12

Stage I: With IF

If the 24-hour urinary excretion of 'hot' B12 is low→

- Test is repeated using the same procedure as in stage I but in addition high oral dose of IF

Interpretation→
If the 24-hour urinary excretion of 'hot' B12 is **now normal (>10% of the oral dose of 'hot' B12) → IF deficiency (Pernicious anaemia)**

Stage II: with antibiotics

If the 24-hour urinary excretion of 'hot' B12 is still low→

- Test is repeated after a course of treatment with antibiotics or anti-inflammatory drugs.

Interpretation→
If the 24-hour urinary excretion of 'hot' B12 is now normal (>10% of the oral dose of 'hot' B12) → Malabsorption or Bacterial overgrowth

Stage III: with Pancreatic enzymes.

If the 24-hour urinary excretion of 'hot' B12 is still low→

- Test is repeated after a course of treatment with Pancreatic enzymes.

Interpretation→
If the 24-hour urinary excretion of 'hot' B12 is now normal (>10% of the oral dose of 'hot' B12) →Chronic pancreatitis

Schilling Test

	^{58}Co-Cbl	W/intrinsic Factor	After 5 Days of Antibiotics	W/Pancreatic Enzymes
Pernicious Anemia	Reduced	Normal	Reduced	Reduced
Bacterial overgrowth	Reduced	Reduced	Normal	Reduced
Chronic pancreatitis	Reduced	Reduced	Reduced	Normal

Special Tests for for folate deficiency

Urinary excretion of FIGLU

- Folic acid is required for conversion of formiminoglutamic acid (FIGLU) to glutamic acid in the catabolism of histidine.
- On oral administration of histidine→ urinary excretion of FIGLU is increased if folate deficiency is present



Treatment

- Vitamin B12 deficiency → Hydroxycobalamin as intramuscular injection 1000 µg for 3 weeks
- Folic acid deficiency → Oral folic acid 5 mg tablets daily for 4 months.

(10.16) PERNECIOUS ANEMIA

University Exams

Short questions

1. What is Pernecious anemia. Write its Pathogenesis, morphological features and Lab diagnosis

Very Short questions

1. Pathogenesis of Pernecious anemia
2. Lab diagnosis of Pernecious anemia

Overview

- Introduction
- Pathogenesis
- Morphological features
- Lab diagnosis
- Treatment

Introduction

- Pernicious anemia is megaloblastic anemia due to vit. B12 deficiency (due to IF deficiency)
- Intrinsic factor is secreted from **parietal cells of stomach.**

Pathogenesis

Autoantibodies against gastric parietal cells
↓
Atrophy of gastric mucosa
↓
IF absent
↓
No absorption of Vitamin B 12
↓
Vitamin B 12 deficiency
↓
Pernecious anemia



Morphologic features

- Gastric atrophy
- Intestinalization of stomach (replacement of gastric mucosa by muscus-secreting globlet cells)
- All features of megaloblastic anemia→ Neurologic abnormalities such as peripheral neuropathy and spinal cord damage
- Associated with increased risk of gastric cancer

Lab diagnosis

- Schilling test

Treatment

- Parenteral vitamin B12 replacement therapy
- Corticosteroids

(10.17) APLASTIC ANEMIA AND PURE RED CELL APLASIA (PRCA)

University Exams

Short questions

1. What is Aplastic anemia. Write its Etiology, Pathogenesis, Clinical features and Lab diagnosis

Very Short questions

1. Four causes of aplastic anemia.
2. Peripheral blood picture in aplastic anemia.
3. Dry tap
4. Pure Red Cell Aplasia (PRCA)

Overview

- Aplastic anaemia →
  - Introduction
  - Etiology
  - Pathogenesis
  - Clinical features
  - Lab diagnosis
  - Treatment
- Pure red cell aplasia (PRCA)
  - Introduction
  - Causes

Aplastic anaemia

Introduction →

- Aplastic anaemia result from **aplasia of the bone marrow**
- In aplastic anaemia, there is **pancytopenia** i.e. simultaneous presence of
  1. Anaemia
  2. Leucopenia
  3. Thrombocytopenia



Etiology

1. Primary aplastic anaemia
   a) Fanconi's anaemia (congenital)
   b) Immunologically-mediated (acquired)
2. Secondary aplastic anaemia

a) Drugs e.g. with antimetabolites (methotrexate), mitotic inhibitors (daunorubicin), alkylating agents (busulfan), nitroso urea, anthracyclines.
b) Toxic chemicals e.g. benzene derivatives, insecticides, arsenicals.
c) Infections e.g. infectious hepatitis, EB virus infection, AIDS, other viral illnesses.

Pathogenesis

Aplasia of bone marrow
↓
Sufficient reduction in no. of haematopoietic pluripotent stem cells
↓
Lower counts of all 3 blood cells types
a) Red blood cells (anemia, reticulocytopenia)
b) WBCs (leukopenia/ neutropenia)
c) Platelets (thrombocytopenia)
↓
Pancytopenia with hypocellular bone marrow

Clinical features

1. Anaemia → mild progressive weakness and fatigue.
2. Thrombocytopenia → Haemorrhage from various sites such as from skin, nose, gums, vagina, bowel, CNS and retina.
3. Leucopenia →Infections
4. The lymph nodes, liver and spleen are generally not enlarged.

Laboratory findings

1. Blood picture
2. Bone marrow picture

1) Blood picture

a) Haemoglobin → Fall
b) Red cells → Normocytic normochromic anaemia
c) Reticulocyte → reduced or zero
d) WBC → Leucopenia The absolute granulocyte count is particularly low (below 1500/µl) with relative lymphocytosis.
e) Platelet → Thrombocytopenia

2) Bone marrow picture

Cellularity →

- A bone marrow aspirate → 'dry tap'
- A trephine biopsy → Hypocellular or aplastic marrow due to replacement by fat.
- Haematopoietic stem cells bearing CD34 marker are markedly reduced or absent.



Healthy bone marrow



Aplastic anemia
(Dry tap)

Treatment

1. General management

a) Identification and elimination of cause
b) Supportive care → blood transfusions, platelet concentrates

2. Specific treatment

   a) Marrow stimulating agents such as androgen
   b) Immunosuppressive therapy

3. Bone marrow transplantation

   - Considered in severe cases under the age of 40 years where the HLA matched donor is available.

Pure red cell aplasia (PRCA)

Introduction →

- Selective failure in production of erythroid elements in bone marrow but with normal granulopoiesis and megakaryocytopoiesis.
- Patients have normocytic normochromic anaemia and reticulocytopenia with normal WBC and platelet count.

Causes

1. Congenital PRCA (Blackfan-Diamond syndrome)

   - Mutation in a ribosomal RNA processing gene → RPS19

2. Acquired PRCA

   - Thymoma (most common)

3. Chronic B19 parvovirus infections

(10.18) PANCYTOPENIA

University Exams

Short questions

1. Pancytopenia → define, causes
2. Pancytopenia with hypocellular bone marrow:
3. Pancytopenia with hypercellular bone marrow:

Overview

- Introduction
- Causes

Introduction

- **Cytopenia** → Reduction in either of cellular component of blood
- **Pancytopenia** → Reduction in all 3 cell lines of blood
    - **Hemoglobin** < 13.5 (M)/ 11.5 (F) g/dl
    - **Total leukocyte count** < 4000 /cumm
    - **Platelet** < 150,000/cumm



Causes



Pancytopenia with hypocellular bone marrow:

- Aplastic anemia
- Fanconi anemia
- Dyskeratosis congenital

Pancytopenia with hypercellular marrow:

- Myelodysplasia
- PNH
- Myelofibrosis
- hairy cell leukemia
- megaloblastic anemia
- bone marrow lymphoma
- SLE
- Hypersplenism
- Sarcoidosis
- alcohol

(11.01) INTERPRETATION OF DLC

University Exams

Very Short questions

1. Leukocytosis (definition and causes).
2. Neutrophilia and Neutropenia
3. Causes for lymphocytosis and Lymphopenia
4. Name six causes for monocytosis.
5. Normal and abnormal absolute eosinophil count values.
6. Normal eosinophil count, two causes for eosinophilia.
7. Conditions associated with eosinophilia.
8. Conditions associated with Basophilia.

Overview

- Introduction
- Types of WBCs
- Hematopoiesis
- Normal counts

    1. Neutrophils → Neutrophilia, Neutropenia
    2. Lymphocytes → Lymphocytosis, Lymphopenia
    3. Monocytes → Monocytosis
    4. Eosinophil → Eosinophilia
    5. Basophils → Basophilia

Introduction

Hematology → study of various cells and components of blood→

1. RBCs (erythrocytes)
2. WBCs (leucocytes)
3. Platelets (thrombocytes)

Type of WBC



Hematopoiesis





Normal Counts

	ABSOLUTE COUNT
TLC Adults	4,000-11,000/ml
DLC In Adults Polymorphs (neutrophils) 40-75% Lymphocytes 20-50% Monocytes 2- 1 0% Eosinophils 1-6 % Basophils <1%	 2,000-7,500/ml 1 ,500-4,000/ml 200-800/ml 40-400/ml 10-100/ml

1) Neutrophils (Polymorphs)



- Normal count → 1500 to 7500 cells/mm3
- Characteristic dense nucleus → **2-5 lobes**
- Pale Cytoplasm → numerous fine **violet pink granules**



Granules → 2 types

Primary or azurophilic granules →
- Large and coarse
- Appear early at promyelocyte stage.
- Contain → hydrolases, elastase, myeloperoxidase, cathepsin-G, cationic proteins, permeability increasing protein, and microbicidal protein called defensins.

Secondary or specific granules→
- Smaller and more numerous
- Appear later at myelocyte stage
- Contain → lactoferrin, NADPH oxidase, histaminase, vitamin B12 binding protein, and receptors for chemoattractant and for laminin.

Neutrophilia

Above 7,500 /mm3

Causes →
1. Acute infections and inflammations
2. Intoxication → e.g. uremia, diabetic ketosis, eclampsia, poisonings by chemicals and drugs.
3. Acute hemorrhage
4. Acute hemolysis
5. Disseminated malignancies
6. Myeloproliferative disorders

Neutropenia

Below 1,500 /mm3

Causes →
1. Infections → e.g. typhoid, paratyphoid, brucellosis, influenza, measles, viral hepatitis, malaria, kala-azar etc.
2. Drugs induce aplasia of bone marrow → e.g. antimetabolites, nitrogen mustards, benzene, ionising radiation.
3. Hematological diseases → e.g. pernicious anemia, aplastic anemia, cirrhosis of the liver with splenomegaly, SLE, Gaucher's disease.

2) Lymphocytes

- Normal count → 1500 to 3500 cells/mm3.
- Round or slightly indented nucleus with coarsely-clumped chromatin
- Scanty basophilic cytoplasm.

- Plasma cells are derived from B lymphocytes
- Nucleus of plasma cell is eccentric and has cart-wheel pattern of clumped nuclear chromatin.



Lymphocytosis

Above 3,500 /mm3

Causes
1. Acute infections → e.g. pertussis, infectious mononucleosis, viral hepatitis, infectious lymphocytosis.
2. Chronic infections → e.g. brucellosis, tuberculosis, secondary syphilis.
3. Haematopoietic disorders → e.g. lymphocytic leukemias, lymphoma, heavy chain disease.

Lymphopenia

Below 1,500/µm

Causes
1. Severe bone marrow failure
2. Corticosteroid and immunosuppressive therapy
3. Widespread irradiation

3) Monocytes

- Normal count → 200 to 1000 cells/mm3.
- Largest mature leucocyte
- Large, central, oval, notched or indented or horseshoe-shaped nucleus

Monocytosis

Above 1000/mm3

Causes →
1. Bacterial infections → e.g. TB, SABE, syphilis.
2. 2. Viral infections.
3. Protozoal and rickettsial infections → e.g. malaria, typhus, trypanosomiasis, kala-azar.
4. Haematopoietic disorders → e.g. monocytic leukemia, lymphomas, myeloproliferative disorders
5. Granulomatous diseases e.g. sarcoidosis, IBD
6. Collagen-vascular diseases

4) Eosinophils



- Normal count → 50 to 500 cells/mm3.
- 2 nuclear lobes.
- Deep red staining granules in cytoplasm

Eosinophilia

Above 500/mm3

Causes →
1. Allergic disorders → e.g. Asthma, urticaria, angioneurotic oedema, hay fever, drug hypersensitivity.
2. Parasitic infestations → e.g. intestinal parasitism.
3. Skin diseases → e.g. pemphigus, dermatitis herpetiformis, erythema multiforme.
4. Loeffler's syndrome.
5. Pulmonary infiltration with eosinophilia (PIE) syndrome.
6. Tropical eosinophilia.

5) Basophils



- Normal count → 10 to 100 cells/mm3.
- Coarse intensely basophilic granules

Basophilia

Above 100/µl

Causes →
1. Chronic myeloid leukemia
2. Polycythemia vera
3. Myxedema
4. Ulcerative colitis
5. Hodgkin's disease
6. Urticaria pigmentosa

(11.02) LEUKEMOID REACTION

University Exams

Short questions

1. Leukemoid Reaction
1. Differentiate Leukemoid Reaction from CML

Overview

- Definition
- Types →

1. Myeloid
2. Lymphoid

} Causes, Lab Findings, Differential diagnosis

Definition

- Reactive excessive leukocytosis in peripheral blood resembling that of leukemia in a subject who does not have leukemia.
- Blood picture confusing with leukemia
- Clinical features of leukemia such as splenomegaly, lymphadenopathy and hemorrhages are absent

Types

1. Myeloid
2. Lymphoid

1) Myeloid Leukaemoid Reaction

Causes →

1. Infections e.g. staphylococcal pneumonia, disseminated tuberculosis, meningitis, diphtheria, sepsis, endocarditis, plague, infected abortions etc.
2. Intoxication e.g. eclampsia, mercury poisoning, severe burns.
3. Malignant diseases e.g. multiple myeloma, myelofibrosis, Hodgkin's disease, bone metastases.
4. Severe hemorrhages and severe hemolysis.

Lab Findings

1. Leukocytosis → not exceeding 100,000/µl.
2. Proportion of immature cells → metamyelocytes, myelocytes, and blasts fewer than 5%
3. Infective cases show → toxic granulation and Dohle bodies in cytoplasm of neutrophils.
4. Neutrophil / Leucocyte alkaline phosphatase (NAP or LAP) score in cytoplasm of mature neutrophils in leukemoid reaction is high and is very useful to distinguish it from CML
5. Cytogenetic studies → negative Philadelphia chromosome i.e. t (9; 22) or BCR-ABL fusion



Leukemoid reaction

Toxic granules
Dohle bodies- consist of ribosomes and endoplasmic reticulum

Differential diagnosis

CML

	Feature	Leukemoid Reaction	CML
1	TLC	25,000-100,000/ml	> 1 00,000/ml
2	DLC	• Dominant cells PMN's • Basophils normal	• All maturation stages • Basophilia present
3	NAPscore	Elevated	Reduced
4	Philadelphia chromosome	Absent	Present
5	ABL-BCR fusion gene	Absent	Present
6	Major etiology	Infections, intoxication, disseminated malignancy, severe hemorrhage	Genetic factors, radiations, certain drugs and chemicals
7	Organ infiltration	Absent	May be present
8	Massive splenomegaly	Absent	Present

2) Lymphoid Leukemoid Reaction

Causes →

- Infections e.g. infectious mononucleosis, cytomegalovirus infection, pertussis, chickenpox, measles, infectious lymphocytosis, tuberculosis.

Lab Findings

1. Leukocytosis → not exceeding 100,000/µl.
2. DLC → mature lymphocytes simulating blood picture of CLL.

(11.03) LEUKEMIAS AND LYMPHOMAS (DEFINITION AND CLASSIFICATION)

University Questions

Short questions

1. Define Leukemia and Write its classification
2. Define Lymphoma and Write its classification

Very Short questions

1. Classification of Leukemia
2. Classification of Lymphoma

Overview

- Introduction
- Types of Leukemia
- Myeloblast Versus Lymphoblast
- Lymphoma
- WHO Classification of Lymphoma

Introduction



Types of Leukemia



Acute leukemia

Malignant disease of bone marrow stem cells in which →

- Bone marrow: Diffuse replacement with proliferating neoplastic blast cells that fail to mature.
- Blast cells more than 20% (WHO criteria) of nucleated cells in marrow. (Normally blast cells are less than 5% of nucleated cells in the marrow)
- Peripheral blood: Abnormal numbers and forms of immature WBCs.

- Have a rapidly downhill course.

Chronic leukemias

- Chronic leukemias have easily recognizable late precursor series of leucocytes circulating in large number as predominant leukemic cell type
- Have more indolent behaviour.

Myeloblast Versus Lymphoblast



	Myeloblast	Lymphoblast
Size	Larger	Smaller
Cytoplasm	Moderate	Scanty
Auer rod	May be present	Absent
Nuclear chromatin	Fine	Coarse
Nucleoli	Prominent, 1-4	Indistinct

Lymphoma

- Proliferations that arise as discrete tissue masses (e.g. within lymph nodes, spleen, or extra nodal tissues).



WHO Classification of Lymphoma

- WHO Classification of Myeloid neoplasm
- WHO Classification of Lymphoid neoplasm

WHO CLASSIFICATION OF MYELOID NEOPLASMS

I. MYELOPROLIFERATIVE DISEASES

1. Chronic myeloid leukaemia (CML), {Ph chromosome t(9;22) (q34;11), BCR/ABL-positive}
2. Chronic neutrophilic leukaemia
3. Chronic eosinophilic leukaemia/ hypereosinophilic syndrome
4. Chronic idiopathic myelofibrosis
5. Polycythaemia vera (PV)
6. Essential thrombocythaemia (ET)
7. Chronic myeloproliferative disease, unclassifiable

II. MYELODYSPLASTIC/MYELOPROLIFERATIVE DISEASES

1. Chronic myelomonocytic leukaemia (CMML)

III. MYELODYSPLASTIC SYNDROME (MDS)

1. Refractory anaemia (RA)
2. Refractory anaemia with ring sideroblasts (RARS)
3. Refractory cytopenia with multilineage dysplasia (RCMD)
4. RCMD with ringed sideroblasts (RCMD-RS)
5. Refractory anaemia with excess blasts (RAEB-1)
6. RAEB-2
7. Myelodysplastic syndrome unclassified (MDS-U)
8. MDS with isolated del 5q

IV. ACUTE MYELOID LEUKAEMIA (AML)

1. Acute myeloid leukaemias with recurrent genetic abnormalities
   - i. AML with t(8;21)(q22;q22); RUNX1-RUNX1T1
   - ii. AML with inv (1 6) (p 13.1; q22) or t (1 6;16) (p13.1; q22); CBF & P-MYH 11
   - iii. Acute promyelocytic leukaemia (M3) with t (15;17) (q22; q!2); PML7RAR-CC and variants
   - iv. AML with t (9; l 1) (p22; q23); MLLT3-MLL
   - v. AML with t (6, -9) (p23; q34); DEK-NUP214
   - vi. AML with Inv (3) (q21q26.2) or t (3;3) (q21; q26.2); RPN1-EVI1
   - vii. AML (megakaryoblastic) with t (1;22) (pl 3; ql3); RBM15-MKL1
   - viii. AML with mutated NPM1
   - ix. AML with mutated CEBPA
2. Acute myeloid leukaemia with myelodysplasia related changes
3. Therapy related acute myeloid leukaemia
   - i. Alkylating agent related
   - ii. Topoisomerase II inhibitor related (some may be lymphoid)
4. Acute myeloid leukaemia, not otherwise categorised
   - i. AML minimally differentiated (MO)
   - ii. AML without maturation (M 1)
   - iii. AML with maturation (M2) *
   - iv. Acute myelomonocytic leukaemia (M4)
   - v. Acute monoblastic and monocytic leukaemia (M5a, M5b)
   - vi. Acute erythroid leukaemia (M6)
   - vii. Acute megakaryoblastic leukaemia (M7)
   - viii. Acute basophilic leukaemia

ix. Acute panmyelosis with myelofibrosis

5. Myeloid sarcoma
6. Myeloid proliferations related to Down's syndrome

   i. Transient abnormal myelopoiesis
   ii. Myeloid leukaemia associated with Down's syndrome

7. Blastic plasmacytoid dentritic cell neoplasm

V. ACUTE BIPHENOTYPIC LEUKAEMIA

WHO CLASSIFICATION OF LYMPHOID NEOPLASM				
Precursor B-cell neoplasm	Precursor T-cell neoplasm	Peripheral B-cell neoplasm	Peripheral T-cell and NK-cell neoplasm	Hodgkin's lymphoma
• B-cell acute lymphoblastic leukemia/ lymphoma (B-ALL)	• T-cell acute lymphoblastic leukemia/ lymphoma	• Chronic lymphocytic leukemia/small lymphocytic lymphoma • Mantle cell lymphoma • Follicular lymphoma • Marginal zone B-cell lymphoma • Burkitt's lymphoma • Hairy cell leukemia • Diffuse large B-cell lymphoma • Splenic and nodal marginal zone B-cell lymphoma • Plasmacytoma/plasma cell myeloma • Lymphoplasmacytic lymphoma	• Mycosis fungoides/Sezary syndrome • Large granular lymphocytic lymphoma • Anaplastic large cell lymphoma • Enteropathy associated T-cell lymphoma • Hepatosplenic dt- cell lymphoma • NK-cell leukemia • Angioimmunoblastic T-cell lymphoma • Extranodal NK/T-cell lymphoma	• Classical • Nodular sclerosis • Mixed cellularity • Lymphocyte-rich • Lymphocyte depleted • Lymphocyte predominant

(11.04) CHRONIC MYELOID LEUKEMIA (CML)

University Exams

Long questions

1. Define and classify leukemias. Discus peripheral blood smear, bone marrow, biochemical findings and the characteristic chromosomal abnormality associated with chronic myeloid leukemia.
2. Define and classify leukemias. Describe the clinical features and laboratory diagnosis of Chronic myeloid leukemia (CML).

Short questions

1. Differentiate between leukemoid reaction and CML.
2. Philadelphia chromosome.
3. Laboratory diagnosis of chronic myeloid leukemia
4. Blood picture in chronic myeloid leukemia
5. Chromosomal abnormality in chronic myeloid leukemia
6. t (9;22) / ABL-BCR Translocation
7. Phases of CML
8. Prognostic index in CML

Very Short questions

1. Special stains used to differentiate leukemoid reaction and CML.
2. Peripheral smear findings in chronic myeloid leukemia

Overview

- Introduction
- Age
- Pathogenesis
- Phases of CML
- Clinical features
- Lab diagnosis
- Treatment
- Prognostic factors

Introduction →

Characterized by→

1. Leukocytosis with granulocytic immaturities
2. Basophilia
3. Splenomegaly
4. Distinct chromosomal abnormality - Philadelphia (Ph') chromosome.

Age →

Old age (> 50 years)

Pathogenesis

- Translocations t (9;22) / ABL-BCR Translocation → Philadelphia chromosome
- ABL gene → normal location on chromosome 9
- BCR gene → normal location on chromosome 22



In Translocation

ABL gene (normal location on chromosome 9)
↓
Translocated to chromosome 22
↓
Fuses with BCR gene
↓
ABL-BCR hybrid gene → Philadelphia chromosome
↓
210 KD fusion protein
↓
Abnormal (tyrosine kinase) → signal transduction even without growth factors
↓
Uncontrolled mitosis
↓
CML

Phases of CML

CML (Triphasic leukemia)		
I. Chronic phase	**II. Accelerated phase**	**III. Blast crisis**
Blast cells < 10%	Blast cells 10-19%	Blast cells > 20%

Clinical Features

- Due To Bone Marrow Failure→

  a) Anemia → pallor, lethargy, dyspnea.
  b) Bleeding manifestations → spontaneous bruises, petechiae, bleeding from gums
  c) Infections

- Hypermetabolism → weight loss, lassitude, anorexia, night sweats.
- Splenomegaly → massive

Lab diagnosis





1) Blood Picture

a) Anemia→ Normocytic normochromic
b) Thrombocytopenia
c) White blood cells →

- Marked leukocytosis (approx. 200,000/µl or more)
- Immature Neutrophils ((band forms, metamyelocytes, myelocytes, promyelocytes)
- Basophilia
- Eosinophilia

2) Bone Marrow Examination

a) Cellularity→ Hypercellularity (proliferating myeloid cells)
b) Myeloid cells → predominate in bone marrow with increased myeloid-erythroid ratio
c) Erythropoiesis → Reduced
d) Megakaryocytes → Reduced

3) Cytogenetics

- Show characteristic chromosomal abnormality → Philadelphia (Ph) chromosome → t (9;22) → BCR/ ABL fusion gene

4) Cytochemistry

Neutrophil alkaline phosphatase (NAP) → distinguish CML from Leukemoid reaction
- CML → NAP scores Reduced
- Leukemoid reaction → NAP scores elevated



Treatment

1. Treatment of Anemia and Hemorrhage

    a) Blood transfusions
    b) Platelet transfusions

2. Imatinib oral therapy

    a) Inhibits signal transduction BCR/ABL fusion protein
    b) Induces apoptosis in BCR/ ABL-positive cells and eliminates them

3. Allogenic bone marrow transplantation

Prognostic factors

Index to assess prognosis of CML

Sokal index
a) Age
b) Platelet count
c) Spleen size
d) Circulating blasts
e) **Cytogenetic changes**

Hasford system
a) Age
b) Platelet count
c) Spleen size
d) Circulating blasts
e) **Basophils and eosinophils**

(11.05) ACUTE MYELOID LEUKEMIA (AML)

University Exams

Long questions

1. Define and classify leukemias. Discus peripheral blood smear, bone marrow, biochemical findings and chromosomal abnormality associated with Acute myeloid leukemia.
2. Define and classify leukemias. Write FAB classification of AML. Describe the clinical features and laboratory diagnosis of Acute myeloid leukemia (AML).

Short questions

1. Describe peripheral blood smear and bone marrow aspiration findings in Acute Myeloid leukemia.
2. Laboratory diagnosis of Acute Myeloid leukemia.
3. Blood picture in Acute Myeloid leukemia.
4. FAB Classification of Acute Myeloid leukemia
5. Prognostic factors of AML

Very Short questions

1. Cytochemistry of acute Myeloid leukemias.
2. Peripheral smear findings in acute myeloid leukemia.
3. Enlist cytochemical stains for typing of leukemias.
4. Chloroma or granulocytic sarcoma
5. Aeur rods
6. Faggot cells

Overview

- Introduction
- Age
- Pathogenesis
- Classification
- Clinical features
- Lab diagnosis
- Treatment
- Prognostic factors

Introduction →

Infiltration of malignant myeloid cells into blood, bone marrow and other tissues.

Age →

15 to 40 years

Pathogenesis

a) t (8;21)
b) t (15;17)
c) inv (16)

Classification

FAB Classification

FAB CLASS	OLD NAME
M0:	Minimally differentiated AML
M1:	AML without maturation
M2:	AML with maturation
M3:	Acute promyelocyte leukemia → **DIC**
M4:	Acute myelomonocytic leukemia (Naegeli type)
M5:	Acute monocytic leukemia (Schilling type)
M6:	Acute erythroleukemia (DiGuglielmo's syndrome)
M7:	Acute megakaryocyte leukemia

Clinical Features

1. Due To Bone Marrow Failure→

   a) Anemia → pallor, lethargy, dyspnea
   b) Bleeding manifestations → spontaneous bruises, petechiae, bleeding from gums
   c) Infections

2. Due to organ infiltration by leukemic cells →

   a) Bones→ Pain and tenderness e.g. sternal tenderness
   b) Lymphadenopathy
   c) Splenomegaly
   d) Hepatomegaly
   e) Kidney → Leukemic infiltration
   f) Gum hypertrophy → Common in **AML M4 and AML M5**
   g) Chloroma or granulocytic sarcoma → localized tumor forming mass occurring in skin of orbit (Proptosis)→ common in **AML M2**

Gum hypertrophy
(M4 and M5)

Chloroma
(M2)

Lab diagnosis



1) Blood Picture

a) Anaemia→ Normocytic normochromic
b) Thrombocytopenia (Acute promyelocytic leukemia (M3) associated with DIC).

c) White blood cells →

- Exceed 100,000/µl
- Myeloblasts with Auer rods present (Faggot cells→ Maximum in M3)



2) Bone Marrow Examination

a) Cellularity → Hypercellular
b) Leukemic cells

- Myeloblasts more than 20% of cells in marrow
- Auer rods present in myeloblasts

c) Erythropoiesis → Reduced
d) Megakaryocytes → Reduced

3) Cytogenetics

a) M2 → t (8;21)
b) M3 → t (15;17)
c) M4 → inv (16)

4) Cytochemistry

	AML	ALL
Myeloperoxidase (MPO)	Positive (Negative in M0)	Negative
Sudan Black	Positive	Negative
Non-specific esterase (NSE)	Positive in monocytic series (M3 ,M4 and M5)	Negative
Periodic acid-Schiff (PAS)	Negative (Positive in M6)	Positive
Acid phosphatase	Negative (Diffuse reaction in M4 and M5)	Focal positivity

Treatment

1. Treatment of Anemia and Hemorrhage

   a) Blood transfusions
   b) Platelet transfusions

2. Cytotoxic Drug Therapy

   a) Cytosine arabinoside
   b) Anthracyclines (daunorubicin, adriamycin)
   c) 6-thioguanine

   - Promyelocytic leukemia (M3) → **Tretinoin (retinoic acid)**

3. Allogenic bone marrow transplantation

Prognostic factors

Good prognosis	Bad prognosis
• Age <40 years • M2, M3, M4 forms of AML • Blast cell with Auer rods • TLC < 25 X 10^9/L • t (15;17), t (8;21), inv 16 • Leukemia without preceding MDS	• Age <2 years or >55 years • M0, M6, M7 forms of AML • Complex karyotypes • TLC > 100X10^9/L • Deletions 5q, 7q • AML with preceding MDS or anticancer drug exposure

(11.06) ACUTE LYMPHOBLASTIC LEUKEMIA (ALL)

University Exams

Long questions

1. Define and classify leukemias. Discus peripheral blood smear, bone marrow, biochemical findings and chromosomal abnormality associated with Acute lymphoblastic leukemia.
2. Define and classify leukemias. Describe the clinical features and laboratory diagnosis of Acute lymphoblastic leukemia (ALL).

Short questions

1. Describe the bone marrow findings in ALL including special stains.
2. Describe peripheral blood smear and bone marrow aspiration findings in Acute lymphoblastic leukemia (ALL).
3. Laboratory diagnosis of Acute lymphoblastic leukemia.
4. Blood picture in Acute lymphoblastic leukemia (ALL).
5. Prognostic factors of ALL

Very Short questions

1. Cytochemistry of Acute lymphoblastic leukemia.
2. French-American-British (FAB) classification of ALL
3. Peripheral smear findings in Acute lymphoblastic leukemia.

Overview

- Introduction
- Age
- Pathogenesis
- Classification
- Clinical features
- Lab diagnosis
- Treatment
- Prognostic factors

Introduction →

ALL composed of immature, precursor B (Pre-B) or T (Pre-T) lymphocytes → lymphoblast.

Age →

1. Pre B-cell ALL → 3-5 years
2. Pre T-cell ALL → Adolescent male

Pathogenesis

Pre B cell ALL

- Hyperdiploidy → most common
- Hypodiploidy
- Loss-of function mutations → PAX5, E2A, or EBF genes.
- t (9;22) translocations (BCR and ABL)

Pre T cell ALL

- Gain-of function mutations → NOTCH 1 gene

Remember

t(9;22); (BCR-ABL protein):

B-ALL

- BCR-ABL protein is 190 kDa Stronger tyrosine kinase activity.

CML

- BCR-ABL protein of 210 kDa Weaker tyrosine kinase activity.

Classification

FAB Classification

L1 ALL	L2 ALL	L3 ALL
Commonest type Best prognosis 	Next common type Worse prognosis. 	Rarest type Worst prognosis. 

Clinical Features

1. Due To Bone Marrow Failure→

    a) Anemia → pallor, lethargy, dyspnea
    b) Bleeding manifestations → spontaneous bruises, petechiae, bleeding from gums
    c) Infections

2. Due to organ infiltration by leukemic cells →

    a) Bone pain and tenderness
    b) Lymphadenopathy
    c) Splenomegaly
    d) Hepatomegaly
    e) Compression of mediastinal vessels or airway → In Pre-T ALL
    f) Meningeal involvement
    g) Testicular involvement

Lab Diagnosis



1) Blood Picture



a) Anemia
b) Thrombocytopenia
c) White blood cells →

- Leukocytosis
- Large number of circulating lymphoblasts

ACUTE LYMPHOBLASTIC LEUKEMIA



LYMPHOBLAST
Large cell with round to oval nucleus, with coarse, clumped chromatin, and inconspicuous nucleoli and scanty cytoplasm

2) Bone Marrow Examination

a) Cellularity → Hypercellular
b) Leukemic cells → 30 % blasts in bone marrow (FAB diagnostic criteria)
c) Erythropoiesis → Reduced
d) Megakaryocytes → Reduced

3) Cytogenetics

- Hyperdiploidy
- Hypodiploidy
- Trisomy 4, 7,10
- t (9;12)
- t (9;22)
- t (4;11)

4) Cytochemistry

	AML	ALL
Myeloperoxidase (MPO)	Positive (Negative in M0)	Negative
Sudan Black	Positive	Negative
Non-specific esterase (NSE)	Positive in monocytic series (M3 ,M4 and M5)	Negative
Periodic acid-Schiff (PAS)	Negative (Positive in M6)	Positive
Acid phosphatase	Negative (Diffuse reaction in M4 and M5)	Focal positivity

Treatment

1. Treatment of Anemia and Hemorrhage

    a) Blood transfusions
    b) Platelet transfusions

2. Chemotherapy

    a) Vincristine
    b) Prednisolone
    c) Anthracyclines (daunorubicin, adriamycin)
    d) L-asparaginase

3. Allogenic bone marrow transplantation

Prognostic factors

Criteria	Good	Bad
1. Age	2-10 years	<2 years; > 10 years
2. Gender	Female	Male
3. Race	White	Black
4. CNS, mediastinum, testicular involvement	Absent	Present
5. Peripheral blood blast count	<100000	>100000
6. Cytogenetics	a. Hyperdiploidy (>50 chromosomes) b. Trisomy 4, 7, 10 c. t (9:12) and t (12:21)	a. Hypodiploidy (<50 chromosomes) b. t (8:14) c. t (1:19) d. t (9:22) e. t (4:11)

(11.07) CHRONIC LYMPHOCYTIC LEUKEMIA (CLL)

University Exams

Long questions

1. Define and classify leukemias. Discus peripheral blood smear, bone marrow biopsy and Immunophenotyping associated with Chronic Lymphocytic leukemia (CLL)
2. Define and classify leukemias. Describe the clinical features and laboratory diagnosis of Chronic Lymphocytic leukemia (CLL)

Short questions

1. Peripheral blood smear of CLL
2. Bone marrow biopsy of CLL
3. Immunophenotyping of CLL
4. Laboratory diagnosis of Chronic Lymphocytic leukemia
5. Poor Prognostic factors of CLL
6. Small Lymphocytic Lymphoma (SLL)

Very Short questions

1. Smudge cells

Overview

- Introduction
- Age
- Pathogenesis
- Clinical features
- Lab diagnosis
- Treatment

Introduction →

CLL and SLL are identical neoplasms arise due to an **abnormal neoplastic proliferation of B cells.**

Age →

Median age of **60 years**

Pathogenesis

a) Deletion of 13q (Most common)
b) Deletion of 11q
c) Deletion of 17p
d) Trisomy 12q

Clinical Features

1. Due To Bone Marrow Failure→

    a) Anemia → pallor, lethargy, dyspnea
    b) Bleeding manifestations → spontaneous bruises, petechiae, bleeding from gums
    c) Infections

2. Enlargement of superficial lymph nodes
3. Splenomegaly and hepatomegaly

Lab Diagnosis



1) Blood Picture



a) Anemia → Normocytic normochromic
b) Thrombocytopenia
c) White blood cells →

- Leukocytosis (50,000-200,000/ µl).
- 90% of leucocytes → mature small lymphocytes.
- Smudge or basket cells (degenerated forms) are present due to damaged nuclei of fragile malignant lymphocytes.

CHRONIC LYMPHOCYTIC LEUKEMIA



2) Immunophenotyping

- CLL is a tumor of B-cells
- Therefore, it expresses the B-cell markers such as **CD19, CD20 and surface IgM and IgD**
- In addition, **CD 23 and CD5** are also present

3) Lymph Node Biopsy

- Effacement of normal nodal architecture by a **diffuse proliferation of small, round lymphoid cells** with coarse chromatin, indistinct nucleoli, and scant cytoplasm
- **Scattered nodules (pseudofollicles)** composed of medium sized and large lymphoid cells with dispersed chromatin and distinct nucleoli are observed.
- The diffuse proliferation of small lymphoid cells with pseudofollicles is pathognomonic for SLL.



Lymph node biopsy in CLL

Treatment

- Palliative and symptomatic



(11.08) POLYCYTHEMIA VERA

University Exams

Short questions

1. Diagnostic criteria, Clinical features and Lab Diagnosis of Polycythaemia Vera
1. Pathophysiology of Polycythaemia Vera

Overview

- Definition
- Pathophysiology
- Diagnostic criteria (WHO) → Major and Minor
- Clinical features
- Lab Diagnosis

Definition

Clonal disorder characterised by increased production of all myeloid elements resulting in Pancytosis →

- Increased red cells
- Increased granulocytes
- Increased Platelets

Pathophysiology

Tyrosine kinase JAK2 mutation
↓
Removes autoinhibitory control
↓
Activates kinases
↓
Increased production of all myeloid elements
↓
Pancytosis



Diagnostic criteria (WHO)

a) 3 major criteria
b) 1 minor criterion

a) Major criteria

1. Hb >16,5 g/dL in men, >16.0 g/dL in women or
   Hct >49% in men, >48% in women or
   Red cell mass >25% above mean normal predicted value.
2. Bone marrow biopsy showing hypercellularity for age with trilineage growth (panmyelosis) including prominent erythroid, granulocytic, and megakaryocytic proliferation with pleomorphic, mature megakaryocytes
3. Presence of JAK2 exon 12 mutation

b) Minor criteria

1. Subnormal serum erythropoietin level.

Diagnostic criteria (WHO)

All 3 major criteria
OR
First 2 major criteria and minor criterion

Clinical features

1. Erythrocytosis →
   - Plethoric and cyanotic due to vascular stagnation and deoxygenation.
   - Headache, dizziness and hypertension
2. Basophilia with histamine release → gastrointestinal symptoms like peptic ulceration and intense pruritus.
3. High cell turnover → hyperuricemia, symptomatic gout and urate stones
4. Platelet dysfunction and abnormal blood flow → Bleeding and thrombotic events.

Lab Diagnosis

1. Raised Hb. concentration (above 17.5 g/dl in males and 15.5 g/dl in females).
2. Erythrocytosis (above 6 million/µl in males and 5.5 million/µl in females).
3. Haematocrit (PCV) above 55% in males and above 47% in females.
4. Mild to moderate Leukocytosis (15,000 - 25,000/µl) with basophilia and raised NAP scores.
5. Thrombocytosis with defective platelet function.
6. Bone marrow examination → erythroid hyperplasia or panhyperplasia.



(11.09) MYELODYSPLASTIC SYNDROME

University Exams

Short questions

1. Etiology, Clinical features and Lab Diagnosis of Myelodysplastic Syndromes
1. Types and Morphology of Myelodysplastic Syndromes

Overview

- Definition
- Etiology
- Types
- Clinical Features
- Lab diagnosis
- Morphology

Definition

- Hematopoietic clonal stem cell disorders having →
    - Maturation defects associated with ineffective haematopoiesis
    - Progressive cytopenia
    - Dysplasia in one or more cell lines
    - Risk of development of AML
- Also known as preleukaemic syndromes or dysmyelopoietic syndromes.

Etiology

a) Monosomy 5 and 7
b) Deletion of chromosome 5q, 7q and 20q.
c) Trisomy 8

Remember →

- Most common cytogenetic change in pediatric MDS is monosomy 7
- Most common cytogenetic change in adult MDS is deletion of 5q
- Most common cytogenetic change in MDS (overall) is deletion of 5q

Types

1. Primary or Idiopathic MDS → seen in patients older than 50 years.
2. Secondary or therapy related MDS (t-MDS) → appears 2 to 8 years after myelo suppressive chemotherapy or radiotherapy→ poor prognosis

Remember

- All types of MDS can transform to AML
- AML transformed from MDS will carry bad prognosis

Clinical Features

1. Anemia appreciated by pallor, fatigue and weakness.

2. Fever.
3. Weight loss.
4. Splenomegaly seen in 20% cases of MDS.

Lab diagnosis

Cytopenia in peripheral blood

1. RBCs: Mild to moderate degree of macrocytic or dimorphic anemia.
2. WBCs: low total leukocyte count.
3. Platelets: thrombocytopenia, large hypogranular or giant platelets.

Morphology

- Hypercellular bone marrow.
- Most characteristic finding is dysplastic differentiation of all 3 lineages→

1) Erythroid lineage

a) Ringed erythroblast: Perinuclear Iron accumulation in a ring like pattern due to iron inside mitochondria.
b) Oval macrocytes in peripheral blood.
c) Erythrocyte precursor with nuclear bridging.


Ringed erythroblast



Oval macrocytes


Erythrocyte precursor

2) Granulocytic lineage

a) Dohle bodies→ Coarse blue granule of cytoplasm made up of dilated rough endoplasmic reticulum.
b) Neutrophil with hyper segmented nucleus
c) Pseudo Pelger Huet anomaly → Neutrophil with bilobed nucleus.


Dohle bodies


hyper segmented nucleus


Pseudo Pelger Huet anomaly

3) Megakaryocyte lineage

- Pawn-Ball Megakaryocyte→ Separated lobes of nuclei in megakaryocytes.



Pawn-Ball Megakaryocyte

(11.10) HODGKIN'S LYMPHOMA (HL)

University Exams

Long questions

1. Classify lymphomas and write on Hodgkin's lymphoma.
2. Classify Hodgkin's disease. Describe the pathological features of nodular sclerosis Hodgkin's disease.
3. Classify Hodgkin's lymphoma. List two important features of mixed cellularity type.
4. Classify Hodgkin's disease. Describe the pathological features of lymphocytic rich and lymphocytic depletion Hodgkin's disease
5. A 45-year-old lady presented with painless neck swelling since 3 months associated with loss of weight and fever. Fine needle aspiration cytology (FNAC) of the same showed a polymorphous population of lymphocytes, eosinophils, plasma cells and large binucleate cells with nucleus having owl-eye appearance. What is your diagnosis? Classify the disease in question and write its staging

Short questions

1. Define & classify Hodgkin's Lymphoma.
2. Describe types of Reed-Sternberg cell.
3. Ann Arbor Staging of Hodgkin's Lymphoma.
4. Morphology of Hodgkin's lymphoma mixed cellularity type.
5. Morphology of Hodgkin's lymphoma Nodular sclerosis type.
6. Morphology of lymphocytic rich Hodgkin's disease
7. Morphology of lymphocytic depletion Hodgkin's lymphoma
8. Morphology of Non classical Hodgkin's lymphoma

Very Short questions

1. Classification of Hodgkin's disease.
2. Appearance of diagnostic cell of Hodgkin lymphoma (morphology of Reed-Sternberg cell).
3. Reed-Sternberg cell and its variants.
4. Name the different types of Reed Sternberg cells. Mention the types of Hodgkin's lymphoma where they are seen.

Overview

- Introduction
- Types of RS Cells
- WHO Classification
- Ann Arbor Staging
- Differences between Hodgkin's lymphoma and Non-hodgkin's lymphoma

Introduction

- Arises in a single lymph node and follow orderly spread to anatomically contiguous lymphoid tissue
- Presence of Reed-Sternberg (RS) cells

Types of RS Cells

Classical R S Cell



Lacunar cell

Lymphohistiocytic variant

Pleomorphic RS cell

WHO Classification

Based on Immunophenotyping of RS cells

1. Classical HD → RS cells positive for CD 15 and CD 30
2. Non-classical /Nodular lymphocytic predominance type → RS cells positive for CD 20 and BCL-6 but are negative for CD 15 and CD 30.



Classic subtypes	Non classical
• Nodular sclerosis • Mixed cellularity • Lymphocyte-rich • Lymphocyte depletion	• Lymphocyte predominant

Nodular sclerosis	Mixed cellularity	Lymphocyte rich	Lymphocyte depleted	Lymphocyte predominant (non-classical HL)
– MC type	– MC type in India	–	– Associated with HIV	–
– M = F	– M > F	– M>F	– M>F	– M > F
– Lacunar cell RS cell	– Classical RS cells	– Classical RS cells	– Pleomorphic RS cell	– LH (popcorn cells) RS cell
– CD15 + CD30+	– CD15 + CD30+	– CD15 + CD30 + and CD20-	– CD15 + CD30+	– CD20 + BCL6 +, EMA + – CD15—, CD30-
– No association with EBV	– Associated with EBV	– Associated with EBV	– Associated with EBV	– Not associated with EBV
– Excellent prognosis	– Prognosis very good	– Good to excellent prognosis	– Poor prognosis	– Excellent prognosis
– Adolescent	– Biphasic incidence (young and > 55 years)	– Old age group	– Old age group	– Young males









Ann Arbor Staging

- Stage I → Involvement of single lymph node or single extra lymphoid site
- Stage II → Involvement of 2 or more lymph node on same side of diaphragm
- Stage III → Involvement of both sides of diaphragm
- Stage IV → Diffuse or disseminated involvement of one or more extra lymphoid organs or tissue with or without associated lymph node involvement



Differences between Hodgkin's lymphoma and Non-hodgkin's lymphoma

Feature	Hodgkin's	Non-hodgkin's
1. Cell derivation	B-cell	90% B 10% T
2. Nodal involvement	Localized, may spread to contiguous nodes	Disseminated nodal Spread
3. Extranodal spread	Uncommon	Common
4. Bone marrow involvement	Uncommon	Common
5. Constitutional symptoms	Common	Uncommon
6. Chromosomal defects	Aneuploidy	Translocations, deletions
7. Spill-over	Never	May spread to blood
8. Prognosis	Better (75-85% cure)	Bad (30-40% cure)

(11.11) NON-HODGKIN'S LYMPHOMA (NHL)

University Exams

Long questions

1. Define and classify Non-Hodgkin's Lymphoma. Discuss Burkite's Lymphoma in detail
2. Define and classify Non-Hodgkin's Lymphoma. Discuss Diffuse large B cell lymphoma (DLBCL) Lymphoma in detail
3. Define and classify Non-Hodgkin's Lymphoma. Discuss Follicular Lymphoma in detail
4. Define and classify Non-Hodgkin's Lymphoma. Discuss hairy cell Leukemia in detail

Short questions

1. Burkitt lymphoma.
2. Microscopy and Immunophenotyping of Burkitt lymphoma
3. Molecular pathogenesis of follicular lymphoma.
4. Morphology of follicular center cell lymphoma.
5. Peripheral and bone marrow biopsy in hairy cell Leukemia
6. Pathogenesis
7. Biopsy and Immunophenotyping of DLBCL

Overview

- Diffuse Large B-cell Lymphoma (DLBCL)
- Follicular Lymphoma
- Burkitt lymphoma
- Hairy Cell Leukemia

} Introduction, Age, Clinical features, Pathogenesis, Biopsy, Immunophenotype

Diffuse Large B-cell Lymphoma (DLBCL)

Introduction →

- Most common type of NHL in world and India
- Worse prognosis amongst all types of NHL.
- Mature B-cell tumour

Age →

60 years

Clinical features →

Presents as Rapidly enlarging mass at a single or multiple nodal or extranodal sites.

Pathogenesis

a) Translocation of BCL2 gene → t (14; 18) translocation
b) Mutations of the BCL6 gene.



Biopsy



Lymph node biopsy →

- Composed of large cleaved cells spread in a diffuse pattern.
- Cytoplasm in these tumour cells is pale and abundant
- Nuclei have prominent 1-2 nucleoli.

Immunophenotype→

Tumor cell wall express →

- CD19
- CD20
- CD10
- BCL6

Follicular Lymphoma

Introduction →

B-cell lymphoma arising from germinal centre.

Age →

Middle-aged adults.

Clinical features →

Presents as painless, generalized lymphadenopathy

Pathogenesis

- t (14:18)
- 30 to 50% of cases will transform to diffuse large B-cell lymphoma (DLBCL).





Biopsy

Lymph node biopsy →

- Follicular (nodular) growth pattern
- composed of 2 principal cell types→

  a) **Centrocytes** (predominant) → small cell with cleaved nuclear contours and scant cytoplasm (Buttock cells).
  b) **Centroblasts** → large cell with open nuclear chromatin, several nucleoli, and modest amount of cytoplasm.

Immunophenotype→

Tumor cells will express →

- CD10
- CD19
- CD20
- BCL6
- Surface Ig
- BCL2 (90% of tumor cells)

Burkitt lymphoma

Introduction →

- Mature B-cell lymphomas
- Very aggressive type of tumour but chemoseristive.
- Most common cause of tumor lysis syndrome

Age and clinical features

3 categories →

1. African (endemic) BL
    - Seen in children or young adults.
    - Most commonly affects mandible (face)
    - All cases will be latently infected with EBV.
2. Sporadic (nonendemic) BL
    - Seen in children or young adults.
    - Mos commonly occurs as an ileocecal or peritoneal mass
    - 15% to 20% of the patients are latently infected by EBV
3. HIV associated aggressive lymphomas
    - 25% of the patients are latently infected by EBV.

Pathogenesis

C-MYC gene → normal location on chromosome 8
↓
translocated to chromosome 14
↓
Fuses with IgG gene
↓
overexpression of MYC protein
↓
Uncontrolled mitosis
↓
Burkitt lymphoma



Biopsy

Lymph node biopsy →

- Loss of architecture.
- Involved tissues show diffuse infiltrate of monotonous lymphoid cells
- Appearance of neoplastic lymphoid cells:
    - Medium-sized cells.
    - Round or oval nuclei with several (2–5) nucleoli.
    - Moderate amount of deeply basophilic cytoplasm,
    - Multiple, small, round lipoid (clear) vacuoles
    - Numerous mitotic figures.

Starry sky pattern →

- Tumor cells undergo apoptosis
- Nuclear remnants of these apoptotic cells are phagocytosed and cleared by benign macrophages.
- These macrophages in the background of lymphoid cells creates "starry sky" appearance









Immunophenotype →

Tumor cells will express →

- CD19
- CD20

- CD10
- Surface IgM
- BCL6

- BCL2 is absent

Hairy Cell Leukemia

Introduction →

- Neoplasm of small mature B cells having abundant cytoplasm with fine hair-like cytoplasmic projections when viewed under the phase-contrast microscope→ hence name hairy cell leukemia
- Chemosensitive
- Overall prognosis is excellent.

Age →

Median age → 55

Clinical features →

- Most common presentation → massive splenomegaly
- Lymphadenopathy is rare.
- Pancytopenia occurs in more than 50% of cases due to marrow infiltration and splenic sequestration.

Pathogenesis

Point mutations in serin/threonine kinase BRAF



Peripheral smear

- Hairy cells are B-cells with reniform nuclei and hair like cytoplasmic projections which is best seen by phase contrast microscopy
- Pancytopenia and tear drop RBC due to bone marrow fibrosis.





Biopsy

Bone marrow biopsy →

- Dry tap due to extensive marrow fibrosis.
- Honeycomb appearance of tumor cell arrangement.
- Fried egg cell appearance of tumor cell due to perinuclear halo.



Immunophenotype

Tumor cells express→

- Pan-B-cell markers CD19, CD20 and surface Ig
- Other distinctive markers → CD11c, CD25, CD103 and annexin A1.
- Annexin A1 is best and most specific marker for hairy cell leukemia.

Features	DLBCL	Follicular Lymphoma	Burkitt's lymphoma	Hairy Cell Leukemia
Introduction				
Age				
Clinical features				
Pathogenesis				
Biopsy				
Immunophenotype				

(11.12) PLASMA CELL DISORDERS → MULTIPLE MYELOMA

University Exams

Long questions

1. A 60-year-old man was admitted with pathological fracture. On examination he was found to be anemic. Laboratory examination showed erythrocyte sedimentation rate (ESR); 100 mm/hr., X-ray-multiple lytic lesions. Peripheral smear showed Increased Rouleaux formation of the red cells and thrombocytopenia. Bone marrow examination confirmed the diagnosis.

    a) Whal is your probable diagnosis?
    b) Whal the other hematological and urinary findings that would be positive in this condition?
    c) Describe the bone marrow findings in this condition.

2. Discuss Morphological features, clinical features, laboratory investigations and prognostic factors of multiple myeloma.

Short questions

1. Revised Multiple Myeloma Diagnostic Criteria
2. Radiological appearance in multiple myeloma.
3. Write the bone marrow findings in multiple myeloma.
4. Morphology of plasma cells in multiple myeloma.
5. M protein.
6. Urinary findings in multiple myeloma.
7. Bence Jones proteins and its demonstration/importance.
8. Describe the investigations in multiple myeloma
9. Types of Plasma Cell Disorders

Overview

- Introduction
- Types
- Multiple myeloma →
    - Definition
    - Etiology
    - Pathogenesis
    - Morphological features →
        a) Osseous lesions → Gross, microscopy
        b) Extraosseous lesions
    - Clinical Features
    - Diagnosis
    - Revised Multiple Myeloma Diagnostic Criteria
    - Prognostic factors

Introduction

Abnormal proliferation of immunoglobulin-producing cells (Plasma cells)
↓
Accumulation of monoclonal immunoglobulin in serum and urine
↓
Plasma cell dyscrasias = paraproteinaemias = dysproteinaemias = monoclonal gammopathies



PCD synthesise excess of →

- Complete immunoglobulins
- Light chains (kappa or lambda) → Bence-Jones proteins
- Heavy chains of a single class (alpha, gamma, or mu)



Types

1. Monoclonal gammopathy of undetermined significance (MGUS)
2. Smoldering multiple myeloma
3. Multiple myeloma
4. Waldenstrom's macroglobulinemia
5. Heavy chain disease

Multiple myeloma

Definition →

- Multifocal malignant proliferation of plasma cells derived from a single clone of cells (i.e. monoclonal).
- Peak incidence → 5th-6th decades
- M > F

Etiology

1. Radiation exposure → For eg. → survivors of nuclear attack in World War-II developed myeloma about 20 years later.
2. Epidemiologic factors → Myeloma has higher incidence in blacks. Certain occupations such as farmers, wood workers and leather workers are more prone.

3. Karyotypic abnormalities →

- t (11;14)
- t (4;14)
- Deletion of 13q

4. Oncogenes-antioncogenes →

- Overexpression of MYC and RAS growth promoting oncogenes
- Mutation in p53 and RB growth-suppressing antioncogene

Pathogenesis

Cell-surface adhesion molecules
↓
Bind myeloma cells to bone marrow stromal cells and ECM proteins.
↓
Triggers adhesion - mediated signalling
↓
Production of cytokines by fibroblasts of marrow → IL-6, VEGF, TGF-**β**, TNF-**α** IL-1, lymphotoxin, macrophage inhibitory factor-1**α** (MIP-1**α**) and receptor activator of nuclear factor-kB (RANK) ligand
↓
IL-6 → proliferation and cell survival of tumor cells IL-1, lymphotoxin, VEGF, MIP-1a, RANK ligand, and TNF → osteoclast-activating factor (OAF) → bony destruction



Morphologic Features

a) Osseous (bone marrow) lesions
b) Extraosseous lesions

a) Osseous (Bone Marrow) Lesions

- Multiple myeloma begins in bone marrow.
- Most affected bones are those with red marrow → skull, spine, ribs and pelvis, long bones of the limbs
- Lesions begin in medullary cavity → erode cancellous bone → destruction of bony cortex.



Gross

Normal bone marrow
↓
Replaced by soft, gelatinous, reddish-grey tumours
↓
Focal or diffuse osteoporosis

Radiographically → punched out, rounded, 1-2 cm sized defects

Microscopy

- Hypercellularity of bone marrow
- Myeloma cells >10% of marrow cellularity
- Form clumps or sheets
- Binucleate and multinucleate cells
- Nucleus of myeloma cell is eccentric but usually lacks cart-wheel chromatin pattern (seen in classical plasma cells)
- Cytoplasm → abundant and basophilic
- Perinuclear clear area (Hof area) corresponding to Golgi apparatus
- Contains vacuolisation







Flame cells →

plasma cells with fiery red cytoplasm.



Mott cells or Morular cells→

Plasma cells with multiple grape like bluish cytoplasmic droplets




Russell bodies →

Globular cytoplasmic inclusions in cytoplasm (endoplasmic reticulum)



Dutcher bodies →

Nuclear inclusions.



Rouleaux formation →

- Due to RBCs sticking by M proteins.
- RBCs are arranged in rows like stacks of coins.













b) Extraosseous Lesions

1. Blood →

    - 50% of patients have atypical plasma cells in blood
    - Normocytic normochromic anemia
    - marked RBC rouleaux formation
    - Hyperviscosity of blood
    - Elevated ESR.

2. Myeloma kidney or myeloma nephrosis →

    - By filtration of light chain proteins (Bence Jones proteins) → precipitated in DCT in combination with Tamm-Horsfall proteins as tubular casts.

3. Myeloma neuropathy →

    - Infiltration of nerve trunk roots by tumour cells
    - Nonspecific polyneuropathy

4. Liver, spleen →

    - Hepatomegaly
    - Splenomegaly

5. Systemic amyloidosis →

    - Generalized amyloidosis (AL amyloid)

Clinical Features

1. Bone pain →

    - most common symptom.
    - Involves back and ribs.
    - Pathological fractures may occur

2. Susceptibility to infections →

    - 2nd most common clinical feature.
    - Common bacterial infections → pneumonias and pyelonephritis.
    - Due to hypogammaglobulinaemia

3. Renal failure →

    - Causes → hypercalcaemia, glomerular deposits of amyloid, hyperuricaemia and infiltration of the kidney by myeloma cells.

4. Anaemia →

    - Marrow replacement by tumour cells → Inhibition of haematopoiesis.

5. Bleeding tendencies →

    - Due to thrombocytopenia, deranged platelet function and interaction of the M component with coagulation factors.

6. Hyperviscosity syndrome →

    - Due to hyperglobulinaemia
    - Headache, fatigue, visual disturbances and hemorrhages.

7. Neurologic symptoms →

    - Due to hyperviscosity, cryoglobulins and amyloid deposits

8. Biochemical abnormalities→

    - Hypercalcaemia → due to destruction of bone
    - Hyperuricaemia → from necrosis of tumour and from uraemia related to renal failure
    - Increased b-2 microglobulins in urine and serum.

9. POEMS → simultaneous manifestations of →

    - Polyneuropathy
    - Organomegaly
    - Endocrinopathy

- Multiple myeloma
- Skin changes

Diagnosis

Classic triad →

1. Marrow plasmacytosis of more than 10%
2. Radiologic evidence of lytic bony lesions
3. Demonstration of serum and/or urine M component.







- Rise in total serum protein due to paraproteinaemia
- Normal serum immunoglobulins (IgG, IgA and IgM) and albumin are depressed.
- Paraproteins → abnormal immunoglobulins circulating in plasma and excreted in urine.
- On serum electrophoresis → paraprotein usually appears as a single narrow homogeneous M-band component in g-globulin region

Most frequent paraprotein →

- IgG → 50%
- IgA → 25%
- IgD → 1%
- Only light chains → 20%

Revised Multiple Myeloma Diagnostic Criteria:

New definition of active multiple myeloma →

Clonal bone marrow plasma cells > 10% or biopsy-proven bony or extramedullary plasmacytoma and any one or more of →

- Related organ or tissue impairment ROTI / CRAB features
- Myeloma-defining events (MDE)

Related organ or tissue impairment ROTI or CRAB Features

1. Hypercalcemia → serum calcium >0.25 mmol/L (>1 mg/dL) higher than upper limit of normal or >2.75 mmol/L (>11 mg/dL)
2. Renal insufficiency → creatinine clearance <40 mL per minute or serum creatinine >177pmol/L (>2mg/dL).
3. Anemia→ Hb value of >20g/L below lowest limit of normal or a Hb value <100g/L
4. Bone lesions → one or more osteolytic lesion on skeletal radiography → CT or PET/CT

Myeloma Defining Events (MDE)

1. 60% or greater clonal plasma cells on bone marrow Examination.
2. Serum involved / uninvolved free light chain ratio of 100 or greater
   (a patient's "involved" free light chain-either kappa or lambda is the one that is above the normal reference range; the "uninvolved" free light chain is the one that is typically In or below the normal range).

3. More than one focal lesion on MRI that is at least 5mm or greater in size.

Prognostic factors

Poor Prognostic factors→

1. Serum B-2 microglobulin > 6 mg/dl (best prognostic marker)
2. C-reactive protein>6 mg/dl.
3. Presence of plasmablastic morphology.
4. Serum LDH raised
5. Plasma cell labelling index >3%
6. Serum creatinine> 2 mg/dl.
7. Cytogenetics showing:

    a) Deletion of 13q9; 119; 17p
    b) Monosomy 13;
    c) Hypodiploidy

(11.13) SPLEENOMEGALY

University Exams

Short questions

1. Causes of splenomegaly/splenic enlargement.
1. Causes for massive splenomegaly.
2. Hypersplenism.
3. Enumerate causes of splenomegaly.
4. Classification of splenomegaly.
5. List the disorders associated with Splenomegaly.
6. Gross and Microscopy in splenomegaly.

Overview

- Definition
- Causes
- Degree → Mild, Moderate, Massive
- Gross
- Microscopy
- Hypersplenism

Definition

Occurs in disorders which increase cellularity vascularity of Spleen

Causes

1. Infections
2. Disorders Of Immunoregulation
3. Disorders Of Immunoregulation
4. Haematolymphoid Malignancies
5. Diseases With Abnormal Erythrocytes
6. Storage Disease
7. Miscellaneous



Normal Spleen



Splenomegaly

1) Infections

a) Malaria
b) Leishmaniasis
c) Typhoid
d) Infectious mononucleosis
e) Bacterial septicaemia
f) Bacterial endocarditis
g) Tuberculosis
h) Viral hepatitis

2) Disorders Of Immunoregulation

a) Rheumatoid arthritis
b) Systemic lupus erythematosus
c) Immune haemolytic anaemias
d) Immune thrombocytopenias

3) Altered Splenic Blood Flow

a) Cirrhosis of liver
b) Portal vein obstruction
c) Splenic vein obstruction
d) Congestive heart failure

4) Haematolymphoid Malignancies

a) Hodgkin disease
b) Non-Hodgkin lymphomas
c) Multiple myeloma
d) leukemias
e) Myeloproliferative neoplasms

5) Diseases With Abnormal Erythrocytes

a) Thalassaemia
b) Sickle cell disease
c) Spherocytosis
d) Ovalocytosis

6) Storage Disease

a) Gaucher's disease
b) Niemann-pick's disease

7) Miscellaneous

a) Amyloidosis
b) Primary and metastatic splenic tumours

Degree of splenomegaly

- Mild enlargement
- Moderate enlargement
- Massive enlargement



Mild

- Up to 500 gm
- Palpable up to 5cm below costal margin

Causes →

- CVC of the spleen in CHF
- Acute malaria
- Typhoid fever
- Bacterial endocarditis
- SLE
- Rheumatoid arthritis
- Thalassaemia minor.

Moderate

- 500-1000 gm
- Palpable up to umbilicus

Causes →

- Hepatitis
- Cirrhosis
- Lymphomas
- Infectious mononucleosis
- Haemolytic anaemia
- Splenic abscesses
- Amyloidosis.

Massive

- More than 1000 gm
- Palpable below umbilicus

Causes →

- CML
- Storage diseases
- Thalassaemia major
- Chronic malaria
- Leishmaniasis
- Portal vein obstruction.

Gross

Grossly→

- Congested, tense and cyanotic
- Sectioned surface → gray tan



Microscopy

- **Red pulp →** enlarged due to marked sinusoidal dilatation
- **Capillarization of sinusoids →** Sinusoids may get converted into capillaries
- White pulp → atrophic.



- Fibrous tissue get deposits of haemosiderin pigment → Gamna-Gandy bodies

Hypersplenism

Altered splenic blood flow or production of antibodies against blood cells
↓
Increased sequestration of RBCs, WBCs or platelets in spleen
↓
excessive removal from circulation
↓
Hyperfunctioning spleen
↓
Hypersplenism

Criteria for hypersplenism

1. Splenomegaly
2. Splenic destruction of one or more of cell types causing → anaemia, leucopenia, thrombocytopenia, or pancytopenia
3. Normal or hyperplastic bone marrow cellularity
4. Splenectomy is followed by improvement in severity of blood cytopenia.

(12.01) BLEEDING DISORDERS (DEFINITION AND CLASSIFICATION)

University Exam

Short questions

1. Define and classify Bleeding disorders
2. Steps of Hemostasis
3. Platelet adhesion
4. Platelet aggregation

Very Short questions

5. Coagulation cascade.
6. Primary and secondary clot

Overview

- Definition
- Hemostasis
- Steps of Hemostasis →

    1. Arteriolar vasoconstriction
    2. Primary hemostasis
    3. Secondary hemostasis
    4. Clot stabilization and resorption

- Classification of Bleeding disorders

Definition

- Also known as Hemorrhagic Diathesis
- Characterized by defective haemostasias with abnormal bleeding.

    - Spontaneous
    - Excessive external or internal bleeding following trivial trauma

Hemostasis

Physiological process → bleeding is stopped after injury → protecting integrity of vascular system after tissue injury.



Steps of Hemostasis

1) Arteriolar vasoconstriction

- Occurs immediately and is transient



2) Primary hemostasis

- Formation of the platelet plug





Injury
↓
Disruption of the endothelium
↓
Exposes subendothelial collagen
↓
Platelet adhesion
↓
Platelet activation
↓
Platelet aggregation/recruitment
↓
Primary hemostatic plug



1. Platelet adhesion → Platelets adhere to collagen in sub endothelium due to presence of receptor (Gp) Ia-IIa and Gp Ib-IX
    - vWF forms a bridge between collagen of endothelium and platelet
2. Platelet activation → 2 types of granules
    a) Alpha -granules
    b) Dense granules
3. Platelet aggregation → Adherence of platelets to one another due to presence of receptor Gp IIb-IIIa

3) Secondary hemostasis

- Deposition of fibrin

Injury
↓
Tissue factor binds and activates factor VII or factor XII
↓
A cascade starts (Intrinsic, extrinsic, common)
↓
Fibrinogen into insoluble fibrin
↓
Fibrin meshwork is formed
↓
Secondary hemostasis





4) Clot stabilization and resorption

Counterregulatory mechanisms (tissue plasminogen activator, t-PA, thrombomodulin) → Clot resorption



Classification of Bleeding disorders

1. Hemorrhagic diathesis due to vascular abnormalities
2. Hemorrhagic diathesis related to platelet abnormalities
3. Disorders of coagulation factors
4. Hemorrhagic diathesis due to fibrinolytic defects
5. Combination of all these as occurs in disseminated intravascular coagulation (DIC)

(12.02) THROMBOCYTOPENIA (ITP, TTP, HUS)

University Exam

Long questions

1. Classify bleeding disorders. Describe the pathogenesis and laboratory findings of idiopathic thrombocytopenic purpura (ITP).
2. Describe and classify bleeding disorders. Describe etiology, hematological features, clinical features and laboratory diagnosis of Thrombotic Thrombocytopenic Purpura (TTP)

Short questions

1. Describe Idiopathic thrombocytopenic purpura (ITP). Write its etiopathogenesis and laboratory findings
2. Describe Thrombotic Thrombocytopenic Purpura (TTP). Write its etiopathogenesis and laboratory findings
3. Describe Hemolytic uremic syndrome (HUS). Write its etiopathogenesis and laboratory findings
4. Megakaryocytic thrombocytopenia and non-megakaryocytic thrombocytopenia
5. Differentiate TTP and HUS

Very Short questions

1. Enumerate the causes of thrombocytosis.
2. Causes of megakaryocytic thrombocytopenia
3. Etiopathogenesis of ITP.
4. Bone marrow findings in ITP.
5. Etiopathogenesis of TTP.
6. Etiopathogenesis of Hemolytic uremic syndrome (HUS).
7. Pentard of TTP
8. Triad of HUS

Overview

- Introduction
- Definition
- Types
- Causes
    - ITP
    - TTP
    - HUS

  } Introduction, Pathogenesis, Clinical features, Laboratory findings

Introduction

Hemorrhagic diatheses due to Platelet Disorders →
- Quantitative → Due to reduction in no. of platelets → thrombocytopenia
- Qualitative → Due to defective platelet functions

Definition

- Reduction in peripheral blood platelet count below lower limit of normal i.e. below 150,000/µl
- Spontaneous bleeding is seen usually when count falls below 20,000 cells/ µl
- Prolonged BT with normal PT and aPTT



Types

1. Megakaryocytic thrombocytopenia→ If thrombocytopenia is associated with compensatory increase in megakaryocytes in bone marrow → seen in peripheral destruction of platelets.
2. Amegakaryocytic thrombocytopenia→ If thrombocytopenia is associated with NO compensatory increase in megakaryocytes in bone marrow → due to defect in bone marrow itself



Causes

Causes

If bone marrow is functional → **Megakaryocyte thrombocytopenia**

- **Immune mediated**
  - Coombs test ⊕
    1. ITP
    2. SLE
    3. B cell cancers
    4. Drugs (Quinidine, Heparin)
    5. Miscellaneous : Dengue
- **Non-Immune mediated**
  - Coombs test ⊖
    1. TTP
    2. HUS
    3. DIC
    4. Giant hemangioma

If bone marrow is having reduced function → **Non-megakaryocytic thrombocytopenia**

1. Bone marrow failure
2. B12 and folate deficiency
3. Leukemia ; disseminated cancer
4. Myelodysplastic syndrome
5. Drugs (cytotoxic drugs, Alcohol)

Immune Thrombocytopenic Purpura (ITP)

Introduction →

- Immunologic destruction of platelets by autoantibodies of class IgG

- Type II hypersensitivity reaction
- Due to peripheral platelet destruction, there is **compensatory increase in megakaryocytes in bone marrow**

Pathogenesis

2 Types→

1. Acute
2. Chronic

1) Acute ITP

- In children (2-6 year)
- M = F
- Following recovery from a viral infection

Viral antigens
↓
Formation of **IgG antibodies against viral antigens**
↓
Antibodies Cross react with platelets
↓
Immunologic destruction of platelets
↓
Thrombocytopenia

2) Chronic ITP

- More commonly in adults
- F > M

Pathogenesis → By formation of anti-platelet autoantibodies

Clinical Features

- Petechial hemorrhages
- Easy bruising mucosal bleeding
- Menorrhagia in women
- Nasal bleeding
- Bleeding from gums
- Melaena
- Haematuria
- Intracranial haemorrhage

Laboratory Findings

1. Platelet count → Reduced
2. BT → Increased
3. Blood film → Occasional platelets
4. Bone marrow → Increased no. of megakaryocytes
5. Anti-platelet IgG antibody → in serum of patients



Peripheral Smear

Thrombotic Thrombocytopenic Purpura (TTP) / Moschcowitz disease

Introduction

Pentad →

1. Microangiopathic hemolytic anemia
2. Thrombocytopenia
3. Neurological symptoms
4. Renal dysfunction
5. Fever

Pathogenesis

Deficiency of enzyme called
ADAMTS.13 (vWF metalloprotease) that degrades vWF
↓
Multimers of vWF accumulate in plasma
↓
Promote platelet thrombus formation
throughout the microcirculation
↓
TTP

Clinical Features

- Petechial hemorrhages
- Easy bruising mucosal bleeding
- Menorrhagia in women
- Nasal bleeding
- Bleeding from gums
- Melaena
- Hematuria
- Intracranial hemorrhage

- Formation of microthrombi →Neurological symptoms and Renal dysfunction and Microangiopathic haemolytic anaemia

Laboratory Findings

1. Platelet count → Reduced
2. BT → Increased
3. Blood film → Occasional platelets
4. Bone marrow → Increased no. of megakaryocytes
5. Biopsy → Typical microthrombi in arterioles, capillaries and venules

Hemolytic uremic syndrome (HUS)

Introduction →

Triad →

1. Microangiopathic hemolytic anemia
2. Renal failure
3. Thrombocytopenia

Pathogenesis

2 Types →

1. **Typical** (epidemic, classical, diarrhea-positive) HUS
2. **Atypical** (non-epidemic, diarrhea-negative) HUS

1. Typical (epidemic, classical, diarrhea-positive) HUS

Acute gastroenteritis caused
by E. coli strain 0157: H7
↓
Shiga-like toxin into circulation
↓
Alters endothelial cell function
↓
Platelet activation and aggregation
↓
Typical HUS

2) Atypical (non-epidemic, diarrhea-negative)

Normally Factor H, Factor I, and CD46 prevent excessive
activation of alternative complement pathway
↓
Deficiency of these regulatory proteins
↓
excessive complement activation
↓
Platelet activation
↓
Thrombosis
↓
Atypical HUS

Laboratory Findings

Peripheral smear of HUS and TTP shows features of microangiopathic hemolytic anemia:

1. Fragmented RBCs (Schistocytes)
2. Helmet cells
3. Burr cells
4. Target cells



	TTP	HUS
ADAMT 13	↓	Normal
CNS involvement	+	Not affected
Renal involvement	+	+++

(12.03) FUNCTIONAL PLATELET DISORDERS (BERNARD SOULIER SYNDROME, GLANZMANN'S SNDROME)

University Exam

Short questions

1. Enumerate functional platelet disorders.
2. Bernard-Soulier syndrome
3. Glanzmann's disease

Overview

- Introduction
- Classification
- Bernard-Soulier syndrome
- Glanzmann's disease

Introduction

- Normal platelet count with defect in function of platelet
- Bleeding time prolonged but platelet count will be normal

Classification

Classification	
Disorders of adhesion	
Inherited Bernard-Soulier syndrome	**Acquired** Uremia Acquired vWD
Disorders of aggregation	
Inherited Glanzmann thrombasthenia	**Acquired** FDP inhibition Dysproteinaemias Drugs-Ticlopidine, Gp IIb/IIIa Inhibitors
Disorders of granule release	
Inherited Oculocutaneous albinism Chediak-Higashi syndrome Isolated dense granule deficiency Gray platelet syndrome (combined a and b granule deficiency)	**Acquired** Cardiopulmonary bypass Myeloproliferative disease Drugs- NSAIDs

Bernard-Soulier syndrome →

- Autosomal recessive disorder
- Due to deficiency or dysfunction of glycoprotein Ib-IX
- Defect in **platelet adhesion**

Glanzmann's disease→

- Autosomal recessive disorder
- Due to deficiency/dysfunction of glycoprotein IIb-IIIa
- Thus there is **defective platelet aggregation**
- At birth present with **increased bleeding from umbilical cord stump**.



(12.04) COAGULATION DISORDERS- HEMOPHILIA AND VON WILLEBRAND'S DISEASE

University Exams

Long questions

1. Define hemophilia. Describe briefly a case of Hemophilia A
2. Define hemophilia. Describe pathogenesis, Clinical features and lab diagnosis of Hemophilia B (Christmas Disease)

Short questions

1. Write a note on hemophilia.
2. Pathogenesis of Hemophilia
3. Von Willebrand's disease (clinical feature and laboratory diagnosis).

Overview

- Classification

1. Classic Hemophilia (Hemophilia A)
2. Christmas Disease (Hemophilia B)
3. Von Willebrand's Disease

} Introduction, Pathogenesis, Laboratory Findings

Classification

1. Classic hemophilia or hemophilia A → due to inherited deficiency of factor VIII
2. Christmas disease or hemophilia B → due to inherited deficiency of factor IX
3. Von Willebrand's disease → due to inherited defect of von Willebrand's factor

1) Classic Hemophilia (Hemophilia A)

Introduction →

- Inherited deficiency of factor VIII
- X linked recessive trait
    - Males → Manifests clinically
    - Females → carriers

Pathogenesis

Quantitative reduction of factor VIII

Functions of factor VIII →
a) Regulates activation of factor X in intrinsic coagulation pathway
b) Factor VIII circulates in blood complexed to another larger protein → von Willebrand's factor (vWF)

Laboratory Findings

a) BT is prolonged
b) PT is normal
c) APTT is prolonged
d) Specific assay for factor VIII shows lowered activity.

2) Christmas Disease (Hemophilia B)

Introduction →

- **Inherited deficiency of factor IX** (Christmas factor or plasma thromboplastin component)
- Hemophilia B is **rarer** than hemophilia A
- **X linked recessive trait**
    - Males → Manifests clinically
    - Females → carriers

Pathogenesis

Quantitative reduction of factor IX

Laboratory Findings

a) BT is prolonged
b) PT is normal
c) APTT is prolonged
d) Specific assay for factor IX shows lowered activity.

3) Von Willebrand's Disease

Introduction →

- Most common hereditary coagulation disorder
- Due to qualitative or quantitative defect in von Willebrand's factor (vWF)

vWF helps in →
1. Platelet adhesion
2. Factor VIII carrier in plasma.



Pathogenesis

Platelet Adhesion defects
• Deficiency of vWF results in defect in the adhesion of platelets to collagen preventing the formation of hemostatic plug. • Leads to mucus and cutaneous bleeding in the form of epistaxis, menorrhagia and GI bleeding.
Coagulation defect
• Reduced half-life of factor VIII leading to its deficiency • Results in hemorrhages and intramuscular hematoma.

Laboratory Findings

a) BT Prolonged
b) Normal platelet count.
c) Reduced plasma vWF concentration
d) Synthesis of factor VIII remains normal but half-life of decreases due to reduced vWF (carrier) levels→ secondary VIII → Intrinsic pathway is affected → aPTT is increased

e) Positive Hess Capillary Resistance Test (Torniquet Test)

Hess Capillary Resistance Test (Torniquet Test)

- Pressure in it between diastolic and systolic for 5 minutes.
- After deflation → number of petechiae appearing in next 5 minutes in 3 cm^2 area over cubital fossa are counted.
- **Presence of more than 20 petechiae is considered a positive test.**

(12.05) DISSEMINATED INTRAVASCULAR COAGULATION (DIC)

University Exams

Long questions

1. Discuss Etiology, Pathogenesis, Clinical Features and Laboratory Findings of Disseminated Intravascular Coagulation (DIC)

Short questions

1. Pathogenesis of DIC

Overview

- Introduction
- Etiology
- Pathogenesis
- Clinical Features
- Laboratory Findings

Introduction

- DIC is a common acquired coagulation disorder resulting due to excessive activation of the coagulation system, due to massive tissue injury or sepsis.
- The normal anticoagulants and fibrinolytic systems are overwhelmed and cannot control the coagulation activation which becomes systemic, resulting in disseminated microvascular thrombi formation.

Massive tissue injury or sepsis
↓
Excessive activation of coagulation system
↓
Normal anticoagulants and fibrinolytic systems are overwhelmed and cannot control coagulation activation
↓
Becomes systemic
↓
Disseminated microvascular thrombi formation
↓
DIC

Also known as
- Defibrination syndrome
- Consumption coagulopathy
- Thrombo-haemorrhagic disorder

Etiology

1. Massive tissue injury → In obstetrical syndromes (e.g. abruptio placentae, amniotic fluid embolism, retained dead fetus), massive trauma, metastatic malignancies, surgery.
2. Infections → Endotoxaemia, Meningococcal septicemia, malaria, aspergillosis.
3. Widespread endothelial damage → In aortic aneurysm, haemolytic-uraemic syndrome, severe burns, acute glomerulonephritis.
4. Carcinoma → Pancreas, Stomach, Prostate, Lung, Acute promyelocytic leukemia

Pathogenesis

Any of above cause
↓
Activation of coagulation pathway
↓
Formation of thrombus → Ischemia and MAHA
↓
Consumption of coagulation factors
↓
Bleeding
Activation of protective fibrinolytic system
↓
Formation of FDP's
↑



Massive tissue destruction
Products of conception
Sepsis
Endothelial injury
↓
Release of tissue factor
Platelet aggregation
↓
Widespread microvascular thrombosis
↓
Activation of plasmin
Vascular occlusion → **Microangiopathic Hemolytic Anemia**
↓
Fibrinolysis
Proteolysis of clotting factors
Ischemic tissue damage
Consumption of clotting factors and platelets
↓
Fibrin split products
↓
Inhibition of thrombin, platelet aggregation, and fibrin polymerization
→ **Bleeding** ←

Clinical Features

1. Bleeding
2. Organ damage due to ischemia caused by thrombosis



Laboratory Findings

1. Platelet count → Low
2. BT, PT, APTT, TT → Prolonged
3. Plasma fibrinogen levels → Reduced
4. Fibrin degradation products (FDPs) → raised
5. Blood film shows MAHA → schistocytes

(13.01) ATHEROSCLEROSIS

University Exams

Long questions

1. Describe etiology, pathogenesis & clinical features of atherosclerosis.
2. Define atherosclerosis. Describe risk factors, pathogenesis, gross and microscopy of atherosclerosis.
3. Define atherosclerosis. Discuss the risk factors, etiopathogenesis and pathology of atherosclerosis. Also briefly write about the sites and clinical significance of atherosclerosis. Describe the morphology (along with neat and labeled diagram) of an atherosclerotic plaque. Mention the Stages of development of atheromatous plaque
4. Enumerate the risk factors of atherosclerosis and discuss the role of lipid and endothelial injury in its pathogenesis.

Short questions

1. Atheroma-pathogenesis.
2. Role of macrophages in atherosclerosis.
3. Morphology of atheroma.
4. Describe gross and microscopic features of atherosclerotic lesion.
5. List the various risk factors and describe the morphology of atherosclerotic plaque.
6. Define Atherosclerosis. Describe its etiopathogenesis.
7. Microscopic appearance of uncomplicated aortic atheromatous plaque

Very Short Answer

1. Most commonly affected arteries in atherosclerosis.
2. Ennumerate modifiable and non-modifiable risk factors of atherosclerosis.

Overview

- Histology of Artery
- Introduction
- Most commonly affected arteries
- Etiology / Risk factors
- Pathogenesis (Reaction-to-Injury Hypothesis)
- Stages of development of atheromatous plaque
- Microscopy
- Clinical features

Histology of Artery

1. Tunica intima
2. Tunica media
3. Tunica adventitia





Introduction

- Thickening and hardening of large and medium-sized muscular arteries due to involvement of tunica intima
- Characterised by fibrofatty plaques or atheromatous plaques that protrude into vessel



Most commonly affected arteries in decreasing order

Mnemonic → **ACP** **D**elhi **T**raffic **IS** **C**ute

A	**A**bdominal aorta
C	**C**oronaries (LAD>RCA>LCX)
P	**P**oplitial
Delhi **T**raffic	**D**escending **t**horasic
IS	**I**nternal carotid
Cute	**C**erebral arteries (circle of willis)

Etiology/ Risk Factors

Etiology or Risk Factors

Modifiable
- Obesity
- Hyperlipemia
- Hypertension
- Diabetes
- Alcohol
- Cigarette smoking
- Physical inactivity
- Lipoprotein (a)
- Trans- unsaturated fat intake
- Infections like Chlamydia, Pneumoniae, CMV and herpes viruses

Non-modifiable
- Old age
- Male gender
- Stress (type A personality)
- Hyperhomocysteinemia

Pathogenesis (Reaction-to-Injury Hypothesis)

1) Endothelial injury

Etiological culprits
↓
Endothelium injury
↓
Endothelium is activated
↓
Increased expression of adhesion molecule- VCAM-1
↓
Increased permeability to endothelium
↓
Lipoproteins accumulate within intima



Chronic endothelial "injury":
- Hyperlipidemia
- Hypertension
- Smoking
- Homocysteine
- Hemodynamic factors
- Toxins
- Viruses
- Immune reactions

2) Migration of leukocytes

Monocytes and T lymphocyte cross endothelial barrier
↓
Accumulate in subendothelial intimal space
↓
Monocytes Become macrophages
↓
Activated macrophages form oxygen free radicals
↓
Cause oxidation of LDL
↓
Oxidized LDL ingested by macrophages
↓
Form foam cells



Endothelial dysfunction
(e.g., increased permeability, leukocyte adhesion),
monocyte adhesion and emigration



Macrophage activation,
smooth muscle recruitment

3) Smooth muscle cell migration and proliferation

Inflammatory cells in subendothelial intimal space
↓
Secrete cytokines (PDGF, TGF-α, FGF)
↓
Migration of smooth muscle cells from media to subendothelial intimal space
↓
Proliferation of smooth muscle cells in intima
↓
Smooth muscle cell also ingest lipid



4) Maturation of plaque

Smooth muscle cells synthesize collegen in intima
↓
Mature fibrofatty atheroma





Stages of development of atheromatous plaque

- Type I (Fatty dot) → due to accumulation of isolated foam cell →not visible.
- Type II (fatty streaks) → Earliest visible lesion
- Type III (intermediate) → small extracellular lipid pool
- Type IV (atheroma) → core of extracellular lipid
- Type V (fibroatheroma or mature atherosclerosis)→ Fully developed lesion
- Type VI (Complicated) → Rupture or thrombus



Microscopy

- Superficial luminal part of fibrous cap→ composed of smooth muscle cells and collagen.
- Cellular area under fibrous cap (shoulder)→comprised by a mixture of macrophages, foam cells, lymphocytes and lipid laden smooth muscle cells
- Deeper central soft core → consists of extracellular lipid material, cholesterol clefts



Clinical Effects

1. Slow luminal narrowing causing **ischaemia and atrophy.**
2. Sudden luminal occlusion causing **infarction necrosis.**
3. Propagation of plaque by formation of **thrombi and emboli.**
4. Formation of **aneurysmal dilatation and eventual rupture.**



(13.02) ANEURYSMS & AORTIC DISSECTION

University Exams

Short questions

1. Syphilitic (Lutic) aortitis.
2. Define Aneurysms. Write its Classification
3. Atherosclerotic aneurysm
4. Enumerate types and causes of aneurysm. Describe gross features of abdominal aortic aneurysm.
5. Mycotic aneurysm.
6. True and False aneurysm
7. Cor bovinum
8. Define Aortic Dissection. Write its Classification
9. Double - barrel aorta
10. Define Aortic Dissection. Write its pathogenesis
11. Morphological features of Aortic Dissection
12. Stanford classification of Aortic Dissection

Overview

Aneurysms

- Introduction
- Clinical features
- Classification →
    1. Based on composition of wall
    2. Based on shape
    3. Based on pathogenetic mechanisms
- Atherosclerotic Aneurysms
- Syphilitic (Luetic) Aneurysms

Aortic Dissection

- Introduction
- Pathogenesis
- Morphological features
- Classification
- Clinical features

Aneurysms

Introduction →

Permanent abnormal dilatation of a blood vessel or heart due to congenital or acquired weakening of vessel wall

↑ Vessel wall stress
↓
Weakening of vessel wall
↓
Outpouching of vessel wall
↓
Abnormal dilation of vessel wall
↓
Aneurysm

Clinical features

1. Thrombosis
2. Thromboembolism
3. Rupture of vessel
4. Compression of neighbouring structures





Classification

1. Based on composition of wall
2. Based on shape
3. Based on pathogenetic mechanisms

1) Based on composition of wall

True aneurysm → composed of all layers of a normal vessel wall.

False aneurysm → having fibrous wall and occur from trauma to vessel

- When wall ruptured → collection of blood outside vessel wall (extravascular hematoma) that is freely communicating with vessel wall (pulsating hematoma)





2) Based on shape

1. Saccular
2. Fusiform
3. Cylindrical
4. Serpentine or varicose
5. Racemose or circoid

4) Based on pathogenetic mechanisms

1. Atherosclerotic (arteriosclerotic) aneurysms → most common type. Wall weakness due to atherosclerosis
2. Syphilitic (luetic) aneurysms → Wall weakness due to tertiary stage of the syphilis.
3. Mycotic aneurysms → Wall weakness due to microbial infection (misnomer)

Atherosclerotic Aneurysms

- Most common type
- Most common site → Abdominal aorta (infrarenal)





Syphilitic (Luetic) Aneurysms

- Develops in tertiary stage of syphilis
- Most common site → Ascending Aorta
- Extend proximally into aortic valve → aortic incompetence

Tertiary syphilis
↓
Inflammatory infiltrate around the vasa vasorum of the adventitia
↓
Endarteritis obliterans
↓
Ischaemic injury to the media
weakening of wall
↓
Syphillitic Anurysm



- Contraction of fibrous scar with in vessel wall → wrinkling of arotic intima → **Tree - barking.**
- **Linear calcification** of ascending aorta.

Tree Bark

appearance

- Aortic valve insufficiency → cardiac hypertrophy → **Cor bovinum/ Cow heart** (increased size of heart)



Aortic Dissection

Introduction →

Blood separates laminar planes of media to form a blood-filled channel within aortic wall





Pathogenesis

Weakened aortic media →

- Hypertensive state (90%)
- Non-hypertensive cases (young persons)

a) Marfan's syndrome (Cystic medial degeneration)
b) Iatrogenic trauma during cardiac catheterisation or coronary bypass surgery.
c) Pregnancy

Morphologic Features

Initial tear
↓
Blood enters into media
↓
Dissecting hematoma spreads along laminar planes between middle and outer third of aortic media
↓
Distal tear → Blood comes out into lumen
↓
Double - barrel aorta



Classification

2 classification schemes→

1. Stanford classification
2. DeBakey Classification



Clinical features

1. **Sudden onset of excruciating pain** → beginning in anterior chest→ radiating to back → moving downward as dissection progresses (confused with MI)
2. Retrograde dissection into aortic root → **Aortic incompetancy**
3. Most common cause of death → **rupture of dissection** into the pericardial, pleural, or peritoneal cavities.

(13.03) HYPERTENSIVE VASCULAR DISEASE

University Exams

Long questions

1. Describe etiology and pathogenesis of Hypertension. Write a note on Vascular Pathology in Hypertension
2. Describe Vascular Pathology in Benign and malignant Hypertension

Short questions

1. Hyaline arteriolosclerosis
2. Hyperplastic arteriolosclerosis
3. Necrotising arteriolitis

Overview

- Introduction
- Types of Hypertension
- Blood Pressure Regulation
- Vascular Pathology in Hypertension →

    1. Hyaline arteriolosclerosis
    2. Hyperplastic arteriolosclerosis
    3. Necrotising arteriolitis

Introduction

- Hypertension → elevation in blood pressure
- Hypertension →

    - Cause end-organ damage
    - Major risk factors for atherosclerosis

	Systolic mm Hg		Diastolic mm Hg
Normal	<120	**and**	<80
Prehypertension	120 - 139	**or**	80-89
Stage I	140- 159	**or**	90-99
Stage II	≥160	**or**	≥100

Types of Hypertension

Essential hypertension →

Affecting Blood Pressure regulating mechanisms

Secondary hypertension →

Secondary to other disease processes → Renal, Endocrine, Cardiovascular, Neurogenic

Malignant hypertension →

Severe hypertension → systolic pressure more than 200 mm Hg, diastolic pressure more than 120 mm Hg

- If untreated, leads to death within 1 to 2 years.

Essential hypertension
Accounts for 90% to 95% of all cases
Secondary hypertension
Renal
Acute glomerulonephritis
Chronic renal disease
Polycystic disease
Renal artery stenosis
Renal vasculitis
Renin-producing tumors
Endocrine
Adrenocortical hyperfunctlon (Cushlng syndrome, primary aldosteronism, congenital adrenal hyperplasia, licorice Ingestion)
Exogenous hormones (glucocorticoids, estrogen [Including Pregnancy-Induced and oral contraceptives], sympathomimetics and tyramine-contalning foods, monoamine oxidase Inhibitors)
Pheochromocytoma
Acromegaly
Hypothyroidism (myxedema)
Hyperthyroidism (thyrotoxicosis)
Pregnancy-induced

Cardiovascular
Coarctation of aorta
Polyarteritis nodosa
Increased intravascular volume
Increased cardiac output
Rigidity of the aorta
Neurologic
Psychogenic
Increased Intracranial pressure
Sleep apnea
Acute stress Including surgery

Blood Pressure Regulation



- **Kidneys and heart** contain cells that sense changes in blood pressure or volume.



Low BP
↓
Kidney will secrete Renin from JG cells

Angiotensinogen
↓
Angiotensin I — Bradykinin PGs/ nitric oxide → vasodilatation
↓ Converting enzyme → inactive peptides
Angiotensin II
↓
Angiotensin receptor → Angiotensin III → inactive fragments
↓
Aldosterone | Vasoconstriction | Sympathetic stimulation

High BP
↓
Heart will secrete ANF
→ **Inhibit Na resorption in DCT**
→ **Vasodilatation**

Vascular Pathology in Hypertension

1) Hyaline Arteriolosclerosis

- In moderate hypertension

Morphologic Features →

- Vascular walls are thickened and Lumina narrowed or obliterated.
- Thickened vessel wall shows structureless, eosinophilic, hyaline material in the intima and media





2) Hyperplastic Arteriolosclerosis

- Characteristic lesion of **malignant hypertension**

Morphologic Features →

- Onion-skin lesion → consists of loosely-placed concentric layers (laminated)of hyperplastic intimal smooth muscle cells like bulb of an onion.



3) Necrotising Arteriolitis

- In malignant hypertension

Morphologic Features →

- Fibrinoid necrosis of vessel wall
- Acute inflammatory infiltrate of neutrophils in adventitia





(13.04) VASCULITIS

University Exams

Short questions

1. Define and classify Vasculitis
2. Buerger's disease.
3. Microscopy of Wegener granulomatosis.
4. Thromboangitis obliterans
5. Temporal Arteritis (Giant cell arteritis)
6. Takayasu Arteritis
7. ANCA associated Vasculitis
8. Ennumerate and describe p-ANCA associated Vasculitis
9. Ennumerate and describe c-ANCA associated Vasculitis
10. Polyarteritis Nodosa (PAN)

Overview

- Definition
- Classification→
  - Based on size of Blood vessels
  - Based on type of inflammation
- Features of important Vasculitis →

1. Temporal Arteritis
2. Takayasu Arteritis
3. PAN
4. Kawasaki's Disease
5. Berger disease
6. Microscopic polyangiitis
7. Churg Strauss Vasculitis
8. Wegener's granulomatosis

} Introduction, Arteries, Morphology, Symptoms

Definition

- Inflammation of vessel wall





Classification

- Based on size of Blood vessels
- Based on type of inflammation

Based on size of Blood vessels

Large Blood Vessel
- Temporal Arteritis
- Takayasu Arteritis

Small Blood Vessel

Medium Blood Vessel
- Polyarteritis Nodosa
- Kawasaki's Disease
- Burger Disease

ANCA Associated

p-ANCA
- Microscopic polyangiitis
- Churg Strauss Vasculitis

c-ANCA
- Wegener's granulomatosis

Non-ANCA Associated
- Hypersensitivity Vasculitis
- H S Purpura
- Behcet's syndrome

Antineutrophilic cytoplasmic antibody (ANCA)

- Autoantibodies directed against primary granules of neutrophil

Cytoplasmic ANCA (c-ANCA)

- Against proteinase 3.
- It is found in Wegner's granulomatosis

Perinuclar ANCA (p-ANCA)

- Against myeloperoxidase.
- It is found in microscopic polyangitis, chrug-strauss syndrome

Based on type of inflammation

Vasculitis granulomatous inflammation	Vasculitis with necrotizing inflammation
• Giant cell arteritis	• Polyarteritis nodosa • Kawasaki disease
• Takayasu's disease	• Microscopic polyangitis
• Wegner's granulomatosis	• Wegner's granulomatosis
• Churg strauss syndrome	• Churg strauss syndrome









Features of important Vasculitis →

1) Temporal arteritis

Introduction →

- Most common form of systemic vasculitis in adults.
- Also known as → Giant cell arteritis



Arteries →

a) Temporal arteries (Most common)
b) Vertebral arteries
c) Ophthalamic arteries

- Segmental involvement
- Nodular thickening

Morphology →

- Granulomatous inflammation
- Foreign body and Langhans type multinulated giant cells

Symptoms →

1. Headache → most common symptom
2. Jaw claudication → most specific symptom
3. Visual symptoms →diplopia, sudden blindness (If ophthalmic artery involved)
4. Arthritis, and polymylagia rheumatica
5. Non-specific constitutional symptoms like fever, fatigue, anorexia and weight loss.

2) Takayasu arteritis

Introduction →

- More common in **Adolescent girls and young women.**

Also known as →
- **Pulseless disease** (Weakening of pulses in upper extremities)
- **Aortic arch syndrome** (strong predilection for aortic arch and its branches)



Arteries →

- Subclavian artery (most common)
- Other → common carotid, abdominal aorta, coeliac, superior mesenteric, renal, vertebral, iliac, pulmonary, and coronary arteries.



Morphology →

Granulomatous inflammation

Symptoms →

1. Marked weakening of pulses in upper extremities → **Pulseless disease**
2. Lower BP and weaker pulse in upper limb than lower limb.
3. Ocular disturbance

3) Classical Polyarteritis nodosa (PAN)

Introduction →

- More common in people with HBV infection

Arteries →

a) Renal artery (Most common)
b) Other arteries (NEVER LUNG)

Morphology →

Fibrinoid necrosis in the vessel wall

Symptoms →

1. Non-specific consititutional symptoms
2. Musculoskeletal symptoms (arthritis, myalgia)
3. Renal artery involvement (major cause of death)
4. Mononeuritis multiplex
5. Other systems involved → GIT, Skin, heart, CNS and genitourinary system.

4) Kawasaki disease

Introduction →

- Seen in children younger than 5 years

Also known as →
- Lymph node syndrome
- Mucocutaneous lymph node syndrome Infant polyarteritis

Arteries →

1. Mucosal artery
2. Coronary artery

Morphology →

Fibrinoid necrosis in the vessel wall

Symptoms →

Mucocutaneous lymph node involvement:
- Erythema of mouth
- Strawberry tongue
- Bilateral conjunctival injection (red eye)
- Beaus lines (transverse grooves on nails),
- Cervical lymphadenopathy



Heart:
- Coronary artery aneurysm
- MI and sudden death
- Myocarditis
- Pericarditis
- Valvular regurgitation

5) Thromboangitis obliterans (Buerger's disease)

Introduction →

- Inflammation of **medium sized arteries** which secondarily extending to **veins and nerves.**
- Occurs **exclusively among heavy cigarrete smoking persons**
- M > F
- Associated with **HLA B-5 and HLA A9**

Arteries →

- Arteries of lower extremities.

Morphology →

- Acute and chronic segmental inflammation of vessels
- Accompanied **thrombosis** in lumen.
- Inflammatory process extends into **contiguous veins and nerves**

Symptoms →

- Intermittent claudication due to ischaemia
- Intense pain affecting the limbs (legs)
- Gangrene of the affected extremities → requiring amputation.

6) Microscopic polyangitis (microscopic polyarteritis)

Introduction →

- Variant of polyarteritis nodosa (PAN)
- But in contrast to classical PAN → microscopic polyangitis involves smaller vessels, i.e. arterioles, venules and capillaries
- Associated with p-ANCA.

Arteries →

1. Renal artery
2. Others

Morphology →

Fibrinoid necrosis in the vessel wall

Symptoms →

1. Kidney → most commonly involved organ → pauci-immune glomomerulonephritis
2. Lung
3. URT (ear, nose, sinuses)
4. Skin (palpable purpura
5. Peripheral nerves (mononeuritis multiplex)

PAN	Microscopic polyangitis
Renal involvement without glomerulonephritis	Renal involvement with glomerulonephritis.
Skin involvement without purpura	Skin involvement with purpura
Lack of pulmonary involvement	Pulmonary involvement seen
Not associated with ANCA	pANCA positive

7) Churg-strauss syndrome

Introduction →

- Allergic granulomatosis and angitis.
- Associated with p-ANCA.

Arteries →

Invoves blood vessels of **lungs, GIT, peripheral nerves, skin, kidney.**

Morphology → Both

1. Granulomatous inflammation (predominant)
2. Necrotizing inflammation

Symptoms →

- Lung involvement is seen in all patients→ asthma
- Allergy →Eosinophilia and raised IgE.
- Other system involved are peripheral nerves (mononeuritis multiplex), skin, heart, kideny, joints, and URT (ear).

8) Wegner's granulomatosis

Introduction→

- Associated with c-**ANCA.**

Arteries →

1. Renal artery
2. Others

Morphology → Both

1. Granulomatous inflammation (predominant)
2. Necrotizing inflammation

Symptoms → Triad →

1. Acute necrotizing granulomas of **upper respiratory tract.**
2. Necrotizing granulomatous vasculitis affecting mainly **lung**
3. **Renal** disease → focal proliferative crescentic glomerulonephritis

Features	Temporal Arteritis	Takayasu Arteritis	PAN	Kawasaki's Disease	Berger disease	Microscopic polyangiitis	Churg Strauss Vasculitis	Wegener's granulomatosis
Intro								
Arteries								
Morphology								
Symptoms								

(13.05) RAYNAUD'S PHENOMENON

University Exams

Short questions

1. Describe pathogenesis and types of Raynaud's phenomenon.
2. Primary and secondary Raynaud phenomenon

Very Short questions

1. Sequence of change in colour in Raynaud's phenomenon.

Overview

- Introduction
- Sequence of change in colour
- Types

Introduction

- Episodic pallor of digits of hands or feet.
- Results from exaggerated vasoconstriction (abnormal spasm) of arteries and arterioles in extremities, particularly fingers and toes, but also occasionally nose, earlobes, or lips.
- Typical sequence of change in colour from white → blue → red



Sequence of change in colour

1. Initially → **white** → because of diminished blood supply.
2. Then → **blue** → because of prolonged lack of oxygen →cyanosis
3. Finally → **red** → blood vessels reopen → flushing phenomenon

Types

1. Primary Raynaud phenomenon
2. Secondary Raynaud phenomenon

Primary Raynaud's	Secondary Raynaud's
More common	Less common
Women under 30 years of age.	Occurs later after 40 years of age
Mild symptoms	Serious symptoms
No known cause	Caused by another underlying medical condition → SLE, scleroderma, Buerger disease,atherosclerosis.
Requires no treatment	Requires more aggressive treatment
Risk factors → Sex, age, climate, and family history	Risk factors → Associated disease, certain occupations causing repetitive trauma, and exposure to certain substances like vinyl chloride

(13.06) TUMOURS OF BLOOD VESSELS

University Exams

Short questions

1. Classify Tumors of blood vessels.
2. Describe Gross, Microscopy and Epidemiological classification of Kaposi sarcoma
3. Hemangioma
4. Morphology of Capillary hemangioma and Cavernous hemangioma

Overview

- Classification
- Benign tumour →Hamangiomas →

    1. Capillary hemangioma
    2. Cavernous hemangioma

- Borderline Tumors → Kaposi's sarcoma →

    - Introduction
    - Most common site
    - Gross
    - Microscopy
    - Epidemiological classification

Classification

Benign Neoplasms, Developmental and Acquired Conditions
Hemangioma • Capillary hemangioma • Cavernous hemangioma **Lymphangioma** • Capillary lymphangioma • Cavernous lymphangioma (cystic hygroma) **Glomus tumor** **Vascular ectasias**
Intermediate-Grade Neoplasms
Kaposi sarcoma Hemangioendothelioma
Malignant Neoplasms
Angiosarcoma Hemgiopericytoma

Benign tumour → Hamangiomas

- Most common benign vascular neoplasms
- Characterized by increased numbers of normal or abnormal vessels filled with blood

2 types →

1. Capillary hemangioma
2. Cavernous hemangioma

1) Capillary hemangioma

- Most common type of vascular tumor
- Occurs in skin, mucus membrane and viscera.
- Strawberry hemangioma or juvenile hemangioma → grows rapidly in first few months and regresses by age of 7 years

Microscopy

- Lobulated unencapsulated
- Aggregates of **closely packed, thin-walled capillaries** which are blood filled and lined by a flattened endothelium.



2) Cavernous hemangioma

- Less common than capillary hemangioma
- Same age and anatomic distribution.
- Involve **deep structures**
- **No tendency to regress** → require surgery.

Microscopy

- Large, cavernous vascular spaces
- Intravascular thrombosis and dystrophic calcification



Borderline Tumors → Kaposi's sarcoma

Introduction →

- Multicentric vascular tumor
- Caused by Human herpes virus-8 (HHV-8) or Kaposi sarcoma associated herpes virus (KSHV).

- Second most common malignancy in HIV (most Common is NHL)

Most common site→

- Most common site for kaposi sarcoma → **Skin of feet** Most common extracutaneous site → **lymph node**
- Most common site in gastrointestinal tract → **Duodenumo**

Gross

Grossly there are three stages →

1. Patch stage (1st stage)
2. Plaque stage (2nd stage)
3. Nodular stage (last **stage)**



Microscopy

Late nodular stage →

- Slit-like vascular spaces containing red blood cells
- Separated by spindle-shaped plump tumour cells (endothelial origin)



Epidemiological classification → 4 forms

Type of Kaposi Sarcoma	Association with HIV	Chief affected organs
Classic/Chronic/ European KS	Absent	Skin plaques and nodules
African/Endemic KS	Absent	No skin lesions; lymphadenopathy present
Transplant / Immunosuppression associated KS	Absent	Lymph nodes, mucosa and visceral organs
Epidemic/AIDS associated KS	Present	Lymph nodes and viscera

(14.01) CONGENITAL HEART DISEASES

University Exams

Short Questions

1. Left-to-right Shunts
2. Morphological features of Ventricular septal defect.
3. Atrial septal defect.
4. Etiopathogenesis of Patent ductus arteriosus
5. Fallot's tetralogy → components and complications.

Overview

- Introduction
- Classification

1. Malposition → Detrocardia
2. Left-to-right shunts →
    - VSD
    - ASD
    - PDA
3. Right-to-left shunts →
    - TOF
    - TGA

Introduction

- Abnormality of heart present from birth.
- 0.5% of newborn children.

Classification

I. Malpositions of Heart	
II. SHUNTS	
A. Left-to-right shunts (Acyanotic or late cyanotic group)	
1. Ventricular septal defect (VSD)	25-30%
2. Atrial septal defect (ASD)	10-15%
3. Patent ductus arteriosus (PDA)	10-20%
B. Right-to-left shunts (Cyanotic group)	
1. Tetralogy of Fallot	6-15%
2. Transposition of great arteries	4-10%
3. Persistent truncus arteriosus	2%
4. Tricuspid atresia and stenosis	1%

III. OBSTRUCTIONS (OBSTRUCTIVE CONGENITAL HEART DISEASE)	
1. Coarctation of aorta	5-7%
2. Aortic stenosis and atresia	4-6%
3. Pulmonary stenosis and atresia	5-7%

Malpositions

- Dextrocardia → apex of heart points to the right side of the chest.

1. Situs inversus → all organs of body are transposed in similar way → heart is in normal position in relation to them.
2. Isolated dextrocardia → associated with major anomalies of the heart



Left-to-right Shunts (Acyanotic or Late Cyanotic Group)

Shunting of blood from left-to-right side of heart
↓
volume overload on right heart
↓
Pulmonary hypertension and RV hypertrophy
↓
Pressure on right side is higher than on left side
↓
Acyanotic / Late cyanotic heart disease



1. Ventricular septal defect (VSD)
2. Atrial septal defect (ASD)
3. Patent ductus arteriosus (PDA)

1) Ventricular Septal Defect (VSD)

- Most common congenital anomaly of heart
- Comprises about 30% of all congenital heart diseases.
- Condition is recognised early in life.

Morphologic Features

VSD
↓
Left-to-right shunt
↓
Increased volume in left side of the heart
↓
Increased pulmonary flow



1. Volume hypertrophy of RV
2. Enlargement and haemodynamic changes in tricuspid and pulmonary valves.
3. Endocardial hypertrophy of RV
4. Pressure hypertrophy of RA
5. Volume hypertrophy of the LA and LV
6. Enlargement and haemodynamic changes in mitral and aortic valves.

2) Atrial Septal Defect (ASD)

- Isolated ASD comprises 10% of congenital heart diseases.

Morphologic Features

1. Volume hypertrophy of RA and RV
2. Enlargement and haemodynamic changes of tricuspid and pulmonary valves.
3. Volume atrophy of LA and LV
4. Small-sized mitral and aortic orifices.



3) Patent Ductus Arteriosus (PDA)

Ductus arteriosus is a normal vascular connection between aorta and pulmonary artery
↓
Normally ductus closes functionally within 1st or 2nd day of life
↓
Due to continued synthesis of PGE2 after birth → Keeps it patent
↓
Its persistence after 3 months of age is abnormal
↓
PDA



Morphologic Features

1. Volume hypertrophy of LA and LV
2. Enlargement and haemodynamic changes of mitral and pulmonary valves.
3. Enlargement of ascending aorta.

Right-to-left Shunts (Cyanotic Group)

Shunting of blood from right side to left side of heart
↓
Entry of poorly-oxygenated blood into systemic circulation
↓
Early cyanosis

1. Tetralogy of Fallot (TOF)
2. Transposition of great arteries (TGA)
3. Persistent truncus arteriosus
4. Tricuspid atresia and stenosis

1) Tetralogy Of Fallot (TOF)

- Tetralogy of Fallot is the most common cyanotic congenital heart disease.

2 two forms of tetralogy →

a) Cyanotic
b) Acyanotic



Morphologic Features

1. VSD
2. Displacement of aorta to right so that it overrides VSD.
3. Pulmonary stenosis → obstruction
4. Right ventricular hypertrophy.

a) Cyanotic Tetralogy

Pulmonary stenosis is
greater and VSD is mild
↓
More resistance to outflow of blood from RV
↓
Right-to-left shunt at ventricular level
↓
Cyanosis



b) Acyanotic tetralogy

VSD is larger and pulmonary
stenosis is mild
↓
Left-to-right shunt
↓
Increased pulmonary flow
↓
Increased volume in left heart
↓
But no cyanosis



2) Transposition of Great Arteries (TGA)

a) Regular transposition
b) Corrected transposition

a) Regular transposition

Aorta which is normally situated to right and posterior with respect to pulmonary trunk → displaced anteriorly and to right.



Transposition of great arteries with
- Aorta arising from RV
- Pulmonary trunk from LV

- So cyanosis from birth.

b) Corrected transposition

Transposition of great arteries with
- Aorta arising from RV
- Pulmonary trunk from LV

Transposition of great veins so that
- Pulmonary veins enter the RA
- Systemic veins drain into LA

- This results in a physiologically corrected circulation.

(14.02) ISCHEMIC HEART DISEASES / CORONARY ARTERY DISEASE

University Exams

Long Questions

1. A 45-year-old man was rushed to the hospital following the sudden onset of an episode of crushing substernal chest pain. He receives advanced life support measures. His course was marked by intractable cardiogenic shock and he died 4 days later. At autopsy, a large transmural anterolateral area of coagulative necrosis was found in the anterolateral wall of the left ventricle.

   a) What is your diagnosis?
   b) What microscopic findings are most likely to be present in this case?
   c) What are the risk factors leading to this condition?
   d) Whal are the complications of this disease?

2. Classify myocardial infarction. Discuss etiopathogenesis of myocardial infarction. Add a note on enzyme changes in myocardial infarction.
3. Write about etiopathogenesis and pathology of myocardial infarction with the diagnostic tests.
4. Describe the morphology/sequential changes seen in myocardial infarction in time. Mention the complications.
5. Classify ischemic heart disease. Describe Etiopathogenesis, types, gross microscopy and clinical features of myocardial infarction.
6. Define Infarct. List the enzymes used in diagnosis of Myocardial infarction and write its gross and microscopic features as per the age of infarct.

Short Questions

1. Pathogenesis of myocardial infarction.
2. Discuss Types of infacts with diagrams
3. Gross assessment of age of myocardial infarction.
4. Microscopic assessment of myocardial infarction.
5. Discuss consequences and complications of myocardial infarction.
6. Serum enzymes in myocardial infarction.
7. Describe laboratory diagnosis of acute myocardial infarction.
8. Types of Angina Pectoris

Very Short Questions

1. Laboratory diagnosis of myocardial infarction.
2. Enzyme markers for myocardial infarction.

Overview

- Introduction
- Coronary artery distribution
- Effects of IHD

Angina Pectoris →

- Definition
- Types

  1. Stable or Classical angina
  2. Unstable or crescendo angina

Myocardial Infarction →

- Definition
- Etiopathogenesis
- Types of infacts
- Location of infacts
- Gross

3. Prinzmetal's variant angina
   - Microscopy
   - Clinical features
   - Diagnosis
   - Complications

Introduction

Arise from Imbalance between myocardial supply and demand for oxygenated blood.



Coronary artery distribution





Effects of IHD



Angina Pectoris

Definition →

- Paroxysmal and recurrent attacks of substernal or precordial chest discomfort caused by transient myocardial ischemia
- Levels of cardiac markers remains unchanged



Types

1. Stable or Classical angina
2. Unstable or crescendo angina
3. Prinzmetal's variant angina

1) Classical (stable) angina

- Most common form of angina
- Caused by reduction of coronary perfusion to a critical level due to coronary atherosclerosis without plaque rupture.
- Chest discomfort →
    - Occurs on exertion
    - Relieved by rest
- Patient complains chest discomfort → heaviness, pressure, squeezing, chocking or smothering.
- It is not described as frank pain
- When patient is asked to localize sensation → he/she will typically place their hand over sternum with clenched fist → **Levine's sign**
- Chest discomfort lasts **2-5 minutes.**
- Crescendo - decrescendo in nature.



2) Unstable angina

- Due to disruption of atherosclerotic plaque with thromosis.

Unstable angina is defined as **chest discomfort** that has at least one of three feature→

1. It occurs at **rest** and lasting **> 10 min.**
2. It is **severe and of new onset** (i.e. within prior 4-6 weeks).
3. It occurs with a **crescendo pattern**



3) Prinzmetal variant angina

- Due to focal spasm in right coronary artery
- Pain occurs **at rest**
- Responds promptly to vasodilators.



Myocardial Infarction

Definition

- Infaction (ischemic necrosis) of myocardium of heart due to decreased blood supply

Etiopathogenesis

- Coronary atherosclerosis
- Superadded changes in coronary atherosclerosis
- Vasospasm



Types of Infarcts

1. **Subendocardial** → occupy inner subendocardial half of the myocardium
2. **Full-thickness or transmural** → involve entire thickness of ventricular wall.





Location of Infarcts

- Depends on which branch of coronary artery is obstructed





Gross

- Necrotic area can be visualized after 2-3 hours by immersion of tissue slices in a solution of triphenyltetrazolium chloride (TTC)
    - Non infarcted myocardium → Brick red color (dehydrogenases enzymes are preserved)
    - Infarcted area → unstained pale zone

Microscopy

0- ½ Hour	→	Normal
½- 4 Hours	→	Waviness
4-12 Hours	→	CN
12-24 Hours	→	CN + few neutrophils
1-3 Days	→	CN + brisk neutrophils
3-7 Days	→	Neutrophils ↓ + Macrophages ↑
7-10 Days	→	Granulation tissue at margin
10-14 Days	→	Granulation tissue well established
2-8 Weeks	→	Collagen
> 2 Months	→	Scar



Changes at 2 months

Changes during the first 24 hours


Changes during the first 48-72 hours


Changes by the end of first week

Clinical features

1. **Pain**: Sudden, severe, crushing and prolonged, substernal or precordial in location, unrelieved by rest or nitroglycerin, radiating to one or both the arms, neck and back.
2. **Indigestion**: Pain is accompanied by epigastric or substernal discomfort interpreted as 'heartburn' with nausea and vomiting.
3. **Shock**: Hypotension,lethargy, cold clammy limbs, peripheral cyanosis, weak pulse
4. **Oliguria**: Urine flow < 20 ml per hour.
5. **Acute pulmonary oedema**: dyspnoea, orthopnoea

Diagnosis

1. ECG changes



NORMAL ECG | ST-SEGMENT ELEVATION | T-WAVE INVERSION | DEEP Q-WAVE

2. Serum cardiac markers





Enzyme	Seen at	Peak	Return to normal	Comment
Myoglobin	2 hours		24 hours	Earliest marker to be detected, not specific
CPK-MB	2-4 hours	24 hours	48-72 hours	Sensitive and specific; Marker of choice for reinfarction
Troponin (T or I)	2-4 hours	48 hours	7-10 days	Most sensitive and specific marker
AST/ SGOT	Within 12 hours	48 hours	4-5 days	Not specific
LDH	24 hours	4-5 days	After 10 days	Only marker raised after 10 days, Not specific



Complications

1. **Arrhythmias** → Most common complication in acute MI, Ooccur due to ischaemic injury or irritation to the conduction system, resulting in abnormal rhythm
2. **Heart failure (Right or left)**
3. **Cardiogenic shock**
4. **Rupture** → most often from the infarcted ventricular wall into the pericardial cavity causing haemopericardium

5. Cardiac aneurysm → occurs in healed infarcts through thin, fibrous, non-elastic scar tissue
6. Pericarditis → Sterile pericarditis appearing on second day is common
7. Postmyocardial infarction syndrome/ Dressler's syndrome → occurs 1 to 6 weeks after attack of MI, characterised by pneumonitis.

(14.03) RHEUMATIC FEVER AND RHEUMATIC HEART DISEASE

University Exams

Long Questions

1. A 12-year-old boy presented with fever, migrating joint pain and palpitation. Child had an upper respiratory infection 3 weeks back. On examination a subcutaneous nodule was observed on the extensor aspect of right elbow. On auscultation heart sounds were weak with tachycardia.

   a) What is your diagnosis?
   b) Describe the morphological features expected in the organ involved.
   c) Add a note on its etiopathogenesis.

2. An 8-year-old boy was brought to clinic by his mother with complaints of shifting joint pain and swelling involving knee joint, elbow joint, etc. and fever for three days. On inquiry it was found that the child had an attack of sore throat. Based on this information:

   a) What is your probable diagnosis?
   b) Enlist the organs, which may get affected.
   c) What is the pathogenesis of this disease?
   d) Enlist the causes of death in this condition.

3. What is etiology of Rheumatic fever? Enumerate clinical manifestations of Rheumatic fever. Describe pathologic changes in heart in cases of acute rheumatic heart disease.
4. Discuss etiopathogenesis and extracardiac lesions of Rheumatic heart disease.
5. Define Rheumatic fever and Rheumatic heart disease. Enumerate (major) Jones criteria. Describe Aschoff nodule formation and its fate. Mention complications of Rheumatic Heart Disease.

Short Questions

1. Classify endocarditis.
2. Etiopathogenesis of rheumatic heart disease.
3. Morphology (pathology) of heart in acute rheumatic carditis.
4. Gross and microscopic appearance of Acute Rheumatic Heart Disease.
5. Microscopy of Aschoff body (Aschoff nodule).
6. Extracardiac manifestations of rheumatic fever.
7. What are modified Jones criteria?
8. Rheumatic endocarditis.
9. Vegetations in Rheumatic carditis.

Very Short Questions

1. MacCallum's patch in Rheumatic endocarditis.
2. List lesions of rheumatic valvulitis.
3. Name extracardiac lesions of rheumatic heart disease.
4. Jones major criteria.
5. Enlist two major and two minor Jones criteria.

Overview

Endocarditis→
- Definition
- Types

Rheumatic fever→
- Introduction
- Etiology
- Pathogenesis
- Revised Jones criteria

1. Rheumatic heart disease → Pancarditis
2. Migratory Polyarthritis
3. Sydenham's chorea
4. Erythema marginatum
5. Subcutaneous nodules

Endocarditis

Definition →

- Inflammation of inner layer of heart → Endocardium
- Involve heart valves (native or prosthetic)



Types

Rheumatic Fever	Infective Endocarditis (Bacterial)	Non-bacterial Thrombotic (Marantic)	Libman Sack's Endocarditis (SLE)
• Small	• Large	• Small (1-5 mm)	• Medium sized (small)
• Firm, Friable (but less than those of NBTE)	• Irregular, Friable (easily detachable)	• Friable (produce emboli)	• Flat, Verrucous
• Along lines of closure	• On upper surface of cusps	• Along lines of closure	• On surface of cusps both surfaces
• Sterile	• Non-Sterile (bacteria)	• Sterile	• Sterile
• Seen in Rheumatic fever	• Caused by micro-organisms	• Seen in hypercoagulabie states	• Seen in SLE

Rheumatic fever

Introduction→

Rheumatic fever (RF) is a **systemic, post-streptococcal, nonsuppurative inflammatory disease** affecting → **heart, joints, central nervous system, skin and subcutaneous tissues.**

Etiology

- Age → children between 5 to 15 years.
- Sex → M = F
- Risk factor → Poor economy and overcrowding

Pathogenesis

- Autoimmune response
- Not caused by bacteria directly effects

Susceptible host
↓
Group A Streptococcus infection
(pharyngitis)
↓
Auto - antibodies formed
↓
Damage to human tissues due to cross-reactivity
between epitopes in bacteria and the host tissue
↓
Molecular mimicry and cross-reactivity



Revised Jones Criteria

Major	Minor	Supportive (GAS Infection)
Carditis	Fever	GAS on Throat swab (Culture)
Arthritis	Arthralgia	Anti-streptolysin O titre (ASOT)
Sydenham's Chorea	↑ PR interval on ECG	Anti-deoxyribonuclease B (Anti-DNase B)
Erythema marginatum	ESR ≥30mm/hr. or CRP ≥30mg/L	
Subcutaneous nodules		

2 Major manifestations + 1 Supportive

OR

1 Major And 2 Minor manifestations + 1 Supportive

1) Rheumatic heart disease → Pancarditis



Rheumatic Valvulitis

- Formation of characteristic vegetations or verrucae



Gross →
- Small
- Multiple
- warty
- Sterile



Valves → Mitral > Aortic > Tricuspid > Pulmonary

Location → along line of closure of leaflets and cusps





Mitral stenosis with and verrucae (view from left atrium)

Deformity → Chronic healed mitral valve in RHD → 'fish mouth' or 'button hole' stenosis



Rheumatic Mural Endocarditis

MacCallum's patch → endocardial surface in the posterior wall of the left atrium just above the posterior leaflet of the mitral valve





Rheumatic Myocarditis

Aschoff bodies →
- Pathogmatic of RHD
- Granulomas

Structure of Aschoff bodies →
Granulomas with Central fibrinoid necrosis

Surrounded by →
- Multinucleate Aschoff cells
- Anitschkow cells (caterpillar cells / owl-eye appearance)
- Lymphocytes
- Plasma cells
- Fibroblast



- Neutrophils (polymorphonuclear cells) are characteristically absent





Rheumatic Pericarditis

- Fibrinous pericarditis
- Two separated surfaces are shaggy due likened to "Bread and butter appearance"



2) Migratory Polyarthritis

- Earliest manifestation (within 2 weeks of sore throat)
- Most common Manifestation (75%)
- Involves larger joints: Knees, ankles, wrists & elbows
- Hot, red, swollen & exquisitely tender
- Migratory in nature
- A dramatic response to even small doses of salicylates

3) Sydenham's chorea

- St. Vitus'dance
- Damage to caudate nucleus
- Prepubertal girls (8-12 yrs)
- A long latency period (1-6 months) between streptococcal pharyngitis & the onset of chorea
- Jerky movements of trunk and the extremities
- Emotional instability

4) Erythema Marginatum

Characteristic rash→

- Erythematous
- Serpiginous
- Macular lesions with pale centers
- Not pruritic
- Occurs primarily on trunk & extremities (not on face) → Skin- bathing suit distribution
- Migrating from place to place.
- No residual scarring occurs.

5) Subcutaneous Nodules

- Firm nodules → Apporox 1 cm in diameter
- Along extensor surfaces of tendons near bony prominences
- Painless and Nontender nodules
- Disappear spontaneously

(14.04) INFECTIVE ENDOCARDITIS

University Exams

Long Questions

1. Describe etiology, pathogenesis, morphology and complications of infective endocarditis.
2. Describe types, pathogenesis and vegetations of infective endocarditis.
3. Differentiate between rheumatic and bacterial endocarditis.
4. What are causes of vegetations in heart? Write in brief on Subacute bacterial endocarditis.

Short Questions

1. Modified Duke criteria for infective endocarditis.
2. Etiopathogenesis of Bacterial endocarditis.
3. Types of Infective endocarditis.
4. Describe various types of cardiac vegetations in endocarditis.
5. Write morphology & complications (cardiac) in bacterial endocarditis.
6. Write Cardiac and Extracardiac complications of infective endocarditis.
7. Morphology of vegetations in infective endocarditis.

Overview

- Introduction
- Types
- Etiology
- Predespositing factors
- Pathogenesis
- Vegetations → Gross,Valves, Location, Deformity
- Complications → Cardiac, Extracardiac
- Modified Duke criteria

Introduction →

- Infection of valvular and mural endocardium caused by different forms of **microorganisms**
- Characterised by typical infected and friable vegetations

Types

1. ABE
2. SABE

Feature	Acute	Subacute
1. Duration	<6 weeks	>6 weeks
2. Most common organisms	Staph. aureus	Strep. viridans
3. Virulence of organisms	Highly virulent	Less virulent
4. Previous condition of valves	Previously normal	Previously damaged
5. Lesion on valves	Invasive, destructive, suppurative	Not invasive or suppurative
6. Clinical features	Features of acute systemic infection	Splenomegaly, clubbing of fingers, petechiae



Etiology



Predisposing factors

- High risk lessions: VSD, MR, prosthetic valves, TOF, PDA, AS
- Moderate risk lesions: TS, TR, PS, and MS
- Low risk: ASD

- **VSD** is most common congenital heart disease involved in infective endocarditis.

Pathogenesis

Conditions producing haemodynamic stress on valves
↓
Cause damage to endothelium
↓
formation of platelet-fibrin thrombi
↓
Get infected from circulating bacteria
↓
Non sterile Vegetations

Vegetations



Gross →
- Large
- Irregular
- Single or multiple
- Typically friable
- Non sterile

Valve → Mitral > aortic > both mitral and aortic valves > valves of right heart

Location →
Upper surfaces of cusps
(Atrial surface of atrioventricular valves and ventricular surface of the semilunar valves)



Deformity →
Most destructive vegetations
Ulceration or perforation of underlying valve leaflet
Myocardial abscesses

Complications



1) Cardiac complications

1. Myocardial abscesses (most common)
2. Abscesses in the valve ring
3. Valvular stenosis or insufficiency
4. Perforation, rupture, and aneurysm of valve leaflets
5. Suppurative pericarditis
6. Cardiac failure

2) Extracardiac complications

- Emboli arising from left side of heart → enter systemic circulation → causes infarcts, abscesses in organs like spleen, kidneys, and brain causing
- Emboli arising from right side of heart → enter pulmonary circulation → pulmonary abscesses



Janeway's spots

- In ABE
- Subcutaneous maculopapular lesions on pulp of fingers
- Painless
- Non-tender

Osler's nodes

- In SABE
- Nodules on finger tips of hands and feet
- Painful
- Tender

Modified Duke criteria

Major

- Blood culture positive for a characteristic organism or persistently positive for an unusual organism
- Echocardiography identification of a valve-related or implant-related mass or abscess, or partial separation of artificial valve
- New valvular regurgitation

Minor

- Predisposing heart lesion or intravenous drug use
- Fever
- Vascular lesions, including arterial petechiae, subungual/splinter hemorrhages, emboli, septic infarcts, mycotic aneurysm, intracranial hemorrhage, Janeway lesions
- Immunological phenomena, including glomerulonephritis, Osler nodes, Roth spots, rheumatoid factor
- Microbiologic evidence, including a single culture positive for an unusual organism
- Echocardiographic findings consistent with but not diagnostic of endocarditis, including worsening or changing of a pre-existent murmur

(14.05) NON-INFECTIVE ENDOCARDITIS (NBTE AND LIBMAN-SACKS ENDOCARDITIS)

University Exams

Short Questions

1. Morphology of vegetations in Libman-sacks Endocarditis
2. Etiopathogenesis and morphology of Non-bacterial Thrombotic Endocarditis

Overview

- Non-bacterial
  Thrombotic Endocarditis (NBTE) / Marantic Endocarditis
  - Introduction
  - Etiology
  - Vegetations → Gross, Valves, Location, Deformity
- Libman-sacks Endocarditis /
  Atypical Verrucous Endocarditis
  - Introduction
  - Etiology
  - Vegetations → Gross, Valves, Location, Deformity

Non-bacterial Thrombotic Endocarditis (NBTE) / Marantic Endocarditis

Introduction →

Involvement of heart valves by sterile thrombotic vegetation

Etiology →

Seen in patients having hypercoagulable state from various etiologies e.g. advanced cancer → mucinous adenocarcinomas, chronic tuberculosis, renal failure and chronic sepsis

Vegetations

Gross →

- Small
- Single or multiple
- More friable than vegetations of RHD
- Sterile

Valves →

Mitral > aortic > tricuspid valve

Location →

Occur along line of closure of leaflets



Deformity →

Organised and healed vegetations appear as fibrous nodules



Libman-sacks Endocarditis / Atypical Verrucous Endocarditis

Introduction →

one of the manifestations of **collagen diseases**

Etiology →

seen in 50% cases of **systemic lupus erythematosus (SLE)**

Vegetations



Gross →

- Medium sized
- Multiple
- Flat
- Granular
- Sterile

Valve →

Mitral > Tricuspid valves

Location →

occur on both surfaces of affected valves, in valve pockets and on adjoining ventricular and atrial endocardium



LSE

Deformity →

do not produce any significant valvular deformity.

(14.06) CARDIOMYOPATHIES

University Exams

Short Questions

1. Cardiomyopathy → definition and types
2. Write etiology, gross and microscopy of Hypertrophic cardiomyopathy
3. Define Dilated cardiomyopathy. Write its etiology and morphology

Overview

- Definition
- Types→

1. Extrinsic cardiomyopathies
2. Intrinsic cardiomyopathies

a) Dilated cardiomyopathy
b) Hypertrophic cardiomyopathy
c) Restrictive cardiomyopathy

} Introduction, Dysfunction, Etiology, Gross, Microscopy

Definition

- Heart muscle disease
- Deterioration of function of myocardium

Types

1. Extrinsic cardiomyopathies → Primary pathology is outside myocardium. Eg. ischemic cardiomyopathy
2. Intrinsic cardiomyopathies → Weakness in muscle of heart that is not due to an identifiable external cause

3 Types →

a) Dilated cardiomyopathy (most common)
b) Hypertrophic cardiomyopathy
c) Restrictive cardiomyopathy (least common)



NORMAL DILATED HYPERTROPHIC RESTRICTIVE

a) Dilated cardiomyopathy

Introduction →

- Most common type of intrinsic cardiomyopathy.
- Characterized by **progressive dilatation of ventricles** (particularly left ventricle) and atria with diminshed pumping activity of heart.
- Associated with concomitant **hypertrophy.**
- Heart is enlarged with dilatation of **all 4 chambers**

Dysfunction →

Systolic dysfunction

Etiology

1. Alcohol
2. Connective tissue disorders (SLE, PAN, RA, dermatomyositis, systemic sclerosis),
3. Metabolic disorders (mucopolysaccharidosis, hemochromatosis Fabrys disease, glycogen storage disease type II)
4. Infections
5. Radiation
6. Thyrotoxicosis
7. Neuromuscular disorders (Friedreich's ataxia, myotonic dystrophy, muscular dystrophy)
8. Deficiency of thiamine, selenium, carnitine, calcium, phosphate or magnesium.

Gross

- The heart is enlarged
- Increased in weight
- Prominent dilatation of all 4 chambers → Globular appearance
- Thickening of ventricular walls even if present is masked by ventricular dilatation

Microscopy

- Hypertrophy of some myocardial fibres and atrophy of others.
- Increased length and width of myocytes
- Interstitial and endocardial fibrosis.





b) Hypertrophic cardiomyopathy

Introduction →

- Also known as Hypertrophic obstructive cardiomyopathy (HOCM)
- Massive myocardial hypertrophy without ventricular dilatation

Dysfunction →

- Because of hypertrophy ventricular filling during diastole is impaired → Diastolic dysfunction
- Systolic function is usually preserved.

Etiology

- Autosomal dominant
- HCM is caused by mutations in any one of several genes that encode sarcomeric protein
- β-myosin heavy chain (β-MHC) > cardiac TnT > α-tropomyosin > myosin-binding protein C (MYBP-C)

Gross



- The hypertrophy of myocardium is typically **asymmetrical**
- Affects **interventricular septum** more than free walls of ventricles
- Most prominent hypertrophy being in subaortic region (outflow area of left ventricle) → left ventricle outflow obstruction
- On cross-section → ventricular cavity is compressed into → Banana-like configration
- Atrial cavity (especially right atrium) may be dilated



Microscopy



- Myocardial Cell disorganisation in ventricular septum
- The bundles of myocardial fibres are irregularly and haphazardly arranged rather than usual parallel pattern and are separated by bands of interstitial fibrous tissue → Myocardial disarray
- Individual muscle cells show hypertrophy and large prominent nuclei.

c) Restrictive cardiomyopathy

Introduction →

- Primary decrease in ventricular compliance ie. ventricles become stiff.
- Ventricles are normal in size without any dilatation of cavity.
- Atrial cavities are dilated.

Dysfunction →

Resulting in impaired ventricular filling during diastole because heart cannot relax normally during diastole → **Diastolic dysfunction**

Etiology

1. Amyloidosis (most common)
2. Radiation
3. Myocardial fibrosis after open heart surgery.
4. Sarcoidosis
5. Hemochromatosis
6. Carcinoid syndrome
7. Fabry's disease
8. Loefflers syndrome
9. Scleroderma,
10. Glycogen depositon

Gross

- Diffuse or patchy, rigid, pearly-white thickening of the mural endocardium

Microscopy

- Fibroelastosis→ Proliferation of collagen and elastic tissue → comprising thickened endocardium.
- Devoid of inflammatory cells.





DILATED HYPERTROPHIC RESTRICTIVE



Features	Dilated cardiomyopathy	Hypertrophic cardiomyopathy	Restrictive cardiomyopathy
Introduction			
Dysfunction			
Etiology			
Gross			
Microscopy			

(14.07) PERICARDITIS

University Exams

Short Questions

1. Define Pericarditis. Write its types
2. Types of Acute Pericarditis

Overview

- Definition
- Causes
- Types

1. Acute Pericarditis →

   a) Serous pericarditis
   b) Fibrinous or serofibrinous pericarditis
   c) Purulent pericarditis
   d) Haemorrhagic pericarditis

2. Chronic Pericarditis →

   a) Tuberculous pericarditis
   b) Chronic adhesive pericarditis
   c) Chronic constrictive pericarditis
   d) Pericardial plaques (milk spots, soldiers' spots)

Definition

Inflammation of pericardium



Causes

1. Infections → Viruses (Echovirus, coxasackie), pyogenic bacteria, tuberculosis, fungi.
2. Immune mediated → RF, SLE, scleroderma, postcardiotomy, post MI (Dressler syndrome).
3. Other → MI, uremia, neoplasia, radiation, trauma

Types

Acute Pericarditis	Chronic Pericarditis
a) Serous pericarditis b) Fibrinous or serofibrinous pericarditis c) Purulent or fibrinopurulent pericarditis d) Haemorrhagic pericarditis	a) Tuberculous pericarditis b) Chronic adhesive pericarditis c) Chronic constrictive pericarditis d) Pericardial plaques (milk spots, soldiers' spots)

1) Acute Pericarditis

Characterized by accumulation of fluid in pericardial space, i.e**. pericardial effusion.**

a) Serous pericarditis → serous exudate.

Causes →
- Acute MI
- Dressier syndrome
- uremia
- SLE
- RF
- Trauma

b) Fibrinous or serofibrinous pericarditis → most common type

Cause → same as serous

c)Purulent pericarditis → Accumulation of pus

Causes →
- Pyogenic organism

d) Hemorrhagic pericarditis → blood mixed with fibrinous effusion

Causes →
- Neoplastic involvement (most common)
- TB
- Uremia
- Cardiac surgery
- Pateints with bleeding disorders

2) Chronic pericarditis

a) Tubercular pericarditis →

Accumulation of **caseous material**

Causes → TB

b) Chronic Adhesive Pericarditis →

Stage of organisation and healing by formation of fibrous adhesions in the pericardium following preceding fibrinous, suppurative or haemorrhagic pericarditis.

- Function of the heart is not compromised

c) Chronic Constrictive Pericarditis →

Dense fibrous or fibrocalcific thickening of pericardium → mechanical **interference with function of heart → reduced cardiac output.**

Causes →
- Tuberculous pericarditis
- Purulent pericarditis
- Haemopericardium

d) Pericardial Plaques (Milk Spots, Soldiers' Spots) →

Opaque, white, shining and well circumscribed areas of organisation with fibrosis in pericardium

Cause →
- Healing of preceding pericarditis

(14.08) COR PULMONALE

University Exams

Short Questions

1. Define Cor pulmonale. Write its 3 causes.
2. Types of Cor pulmonale
3. Pathogenesis of Cor pulmonale

Overview

- Definition
- Pathogenesis
- Types
- Causes
- Morphology

Definition

- Cor pulmonale (cor = heart; pulmonale = lung)
- Also known as pulmonary heart disease
- Disease of right side of heart resulting from disorders of lungs
- Characterised by right ventricular dilatation or hypertrophy, or both.

Pathogenesis

CHRONIC LUNG DISEASES, PULMONARY VASCULAR DISEASE
↓
Pulmonary resistance
↓
Pulmonary hypertension
↓
Pressure overload on RV
↓
RVH, RHF
↓
(COR PULMONALE)





Types

- Acute cor pulmonale → following massive pulmonary embolism resulting in sudden dilatation of the pulmonary trunk, conus and right ventricle.
- Chronic cor pulmonale (more common)→ preceded by chronic pulmonary hypertension

Causes

Following chronic lung diseases → cause chronic pulmonary hypertension → cor pulmonale

1. Chronic emphysema
2. Chronic bronchitis
3. Pulmonary tuberculosis
4. Pneumoconiosis
5. Cystic fibrosis
6. Hyperventilation in marked obesity (Pickwickian syndrome)
7. Multiple organised pulmonary emboli.

Morphology

- In acute cor pulmonale → Dilatation of RV and sometimes of RA
- In chronic cor pulmonale → Increase in thickness of RV wall with RV Dilatation



In acute cor pulmonale



In chronic cor pulmonale

(15.01) NORMAL ANATOMY AND HISTOLOGY

Overview

- Respiratory tract
- Pulmonary Lobes
- Pulmonary Lobules
- Histology
- Alveolar septa

Respiratory tract





Pulmonary Lobes



Pulmonary lobule

- A cluster of three to five terminal bronchioles, each with its appended acinus,













Histology

- Respiratory tract is lined by pseudostratified columnar epithelium
- Alveoli (alveolar sac) are lined by

    1. Type 1 → membranous pneumocytes (95%)
    2. Type 2 → granular pneumocytes (5%) → secrete surfactant

Alveolar septa

- Alveolar walls or alveolar septa are the sites of exchange between the blood and air



(15.02) ACUTE RESPIRATORY DISTRESS SYNDROME (ARDS)

University Exams

Short Questions

1. What is hyaline membrane disease? What is the basic defect in it.
2. Etiopathogenesis of Adult respiratory distress syndrome (ARDS)
3. Gross and microscopy of Hyaline membrane disease.

Overview

- Introduction
- Types →
  1. Neonatal type
  2. Adult type

  } Etiology, Pathogenesis, Clinical features
- Gross
- Microscopy

Introduction

- Severe life-threatening form of progressive respiratory insufficiency which involves pulmonary tissues diffusely
- Formation of hyaline membrane in the alveoli → Hyaline membrane disease (HMD)

Types

2 forms:

1. Neonatal type
2. Adult type → Also known as → Shock lung/ Stiff lung/ Diffuse alveolar damage/ Acute lung injury

1) Neonatal ARDS

Etiology

1. Preterm infants
2. Infants born to diabetic mothers
3. Delivery by caesarean section
4. Infants born to mothers with previous premature infants
5. Excessive sedation of the mother causing depression in respiration of the infant
6. Birth asphyxia from various causes such as coils of umbilical cord around the neck
7. Male preponderance (1.5 to 2 times) over female babies due to early maturation of female lungs
8. Finally, many cases of neonatal ARDS remain idiopathic.

Pathogenesis

Prematurity / Neonatal hypoxia
↓
Deficiency of pulmonary surfactant (normally synthesised by type II alveolar cells)
↓
Increased alveolar surface tension
↓
Atelectasis
↓
Hypoventilation
↓
Ischaemic necrosis of alveolocapillary wall
↓
exudation of plasma proteins
↓
Formation of hyaline membrane on alveolar surface

Clinical Features

1. Dyspnoea within a few hours after birth with
2. Tachypnoea
3. Hypoxia and cyanosis
4. In severe cases death within a few hours.

2) Adult ARDS

Etiology

Infection
Sepsis
Diffuse pulmonary infections
Viral, Mycoplasma, and Pneumocystis pneumonia; miliary tuberculosis
Gastric aspiration
Physical/Injury
Mechanical trauma, including head injuries
Pulmonary contusions
Near-drowning
Fractures with fat embolism
Burns
Ionizing radiation
Inhaled Irritants
Oxygen toxicity
Smoke
Irritant gases and chemicals

Chemical Injury
Heroin or methadone overdose
Acetylsalicylic acid
Barbiturate overdose
Paraquat
Hematologic Conditions
Transfusion associated lung injury (TRALI)
Disseminated intravascular coagulation
Pancreatitis
Uremia
Cardiopulmonary Bypass
Hypersensitivity Reactions
Organic solvents
Drugs

Pathogenesis

Etiological agent
↓
Activated pulmonary macrophages
↓
Release proinflamatory cytokines such as IL8, IL1, and TNF
↓
Number of neutrophils in alveoli is increased.
↓
Neutrophils on activation release products which cause active tissue injury e.g. proteases, platelet activating factor, oxidants and leukotrienes.
↓
Increased vascular permeability
↓
Formation of hyaline membranes



Clinical Features

1. Sudden and severe respiratory distress
2. Tachypnoea
3. Tachycardia
4. Cyanosis
5. Severe hypoxaemia

	Neonatal type	Adult type
Etiology		
Pathogenesis		
Clinical features		

Gross

Lungs in ARDS →

- Normal in size
- Characteristically stiff
- Congested
- Heavy
- Airless so that they sink in water



Microscopy

1. Collapsed alveoli (atelectasis) alternating with dilated alveoli.
2. Necrosis of alveolar epithelial cells→ formation of characteristic eosinophilic hyaline membranes lining respiratory bronchioles, alveolar ducts and alveoli.
3. Interstitial oedema



(15.03) COPD – CHRONIC BRONCHITIS

University Exams

Long Questions

1. What are chronic obstructive pulmonary diseases? Write the etiology, pathogenesis and morphology of Chronic Bronchitis.
2. A 40-year-old smoker male presented with history of chronic cough with profuse expectoration, occasional hemoptysis and also clubbing with coarse crepitation in right lung base:

    a) What is your probable diagnosis?
    b) Discuss the etiopathogenesis, gross and microscopy of target organ involved.

Short Questions

1. Pathogenesis and clinical features of Chronic Bronchitis
2. Reid index.

Overview

- COPD →
    - Definition
    - Types
- Chronic Bronchitis →
    - Introduction
    - Location
    - Etiology
    - Pathogenesis
    - Types
    - Gross
    - Microscopy
    - Clinical features

Chronic Obstructive Pulmonary Disease (COPD)

Definition →

- Group of pathological conditions in which → chronic, partial or complete, obstruction to the airflow at any level from trachea to the smallest airways → functional disability of the lungs

Types

1. Chronic bronchitis
2. Emphysema
3. Bronchial asthma
4. Bronchiectasis



Chronic Bronchitis



Introduction →

Persistent cough with expectoration on most days for at least three months of year for two or more consecutive years.

Location →

Trachea and bronchi

Etiology →

Smoking, Air pollutants

Pathogenesis →

Smoking
↓
Irritates respiratory mucosa and submucosa
↓
Increase in number and size of submucosal gland of trachea and bronchi
↓
over secretion of mucus
↓
Mucous plug
↓
Cough, Dypsnea, Cynosis
↓
Chronic Bronchitis

Gross

- Bronchial wall → thickened, hyperaemic and oedematous
- Lumina of bronchi → contain mucus plugs

Microscopy

1. Increased Reid index (normal 04)
    - Reid index is ratio between thickness of the submucosal mucus glands (i.e. hypertrophy and hyperplasia) in cartilage-containing large airways to that of total bronchial wall
2. Goblet cells increased
3. Squamous metaplasia and dysplasia.
4. Chronic inflammatory cell infiltrate



Clinical Features

1. **Persistent cough** with copious expectoration of long duration
2. **Dyspnoea**
3. Patients are called **'blue bloaters'** due to cyanosis
4. Recurrent **respiratory infections**
5. Features of **right heart failure (cor pulmonale)**
6. Chest X-ray → **enlarged heart with prominent vessels.**

(15.04) COPD – EMPHYSEMA

University Exams

Long Questions

1. What is emphysema? Write the types of emphysema? Describe etiology and pathogenesis of emphysema.

Short Questions

1. Define and classify emphysema.
2. Describe gross and microscopic features of emphysema
3. Describe briefly etiopathogenesis of emphysema.
4. Etiopathogenesis of Panacinar emphysema
5. Very Short Questions
6. List differences between emphysema and chronic bronchitis.
7. Enumerate Types of Emphysema.

Overview

- Introduction
- Location
- Etiology
- Pathogenesis
- Types
- Gross
- Microscopy
- Clinical features

Introduction →

Combination of permanent dilatation of air spaces distal to the terminal bronchioles and the destruction of the walls of dilated air spaces.



Location →

Acinus

Etiology →

Smoking, Air pollutants

Pathogenesis

Normally → Alveolar wall → Balance between →
- Proteases (elastase) → destruction of alveolar wall
- Antiproteases (antielastase / α1-antitrypsin) → prevent this damage

Protease-antiprotease hypothesis
↓
Smoking
↓
Decreases antiprotease
Increases protease
↓
Protease-antiprotease Imbalance
↓
Alveolar wall (Acinus) destruction
↓
Emphysema

Types



1) Centriacinar (centrilobular) emphysema

- Most common type of emphysema seen clinically
- Involves **Respiratory bronchioles (central part of Ainus)**
- **Upper lobe**
- Occurs predominantly in heavy smokers



2) Distal acinar (Paraseptal) emphysema

- Involves distal part of the acinus
- Upper half of lungs.
- Common cause of spontaneous pneumothorax in young adults.



3) Panacinar (Panlobular) emphysema

- Acini are uniformly enlarged from the level of the respiratory bronchiole to the alveoli.
- Lower zones (base)
- Associated with α1-antitrypsin deficiency



4) Irregular emphysema (Para-cicatricial emphysema)

- Most common type of emphysema histologically (Clinically silent)
- Acinus is involved irregularly
- Associated with scarring



Gross

- Lungs → voluminous
- Subpleural bullae and blebs
    - Bullae → rupture of adjacent air spaces



- Blebs → rupture of alveoli directly into subpleural interstitial tissue



Microscopy

1. Acini → distended
2. Their walls stretched and thin
3. Ruptured alveolar walls and spurs of broken septa
4. Inflammatory changes are usually absent

Clinical Features

1. Severe exertional **dyspnoea**
2. Obvious use of accessory muscles of respiration → Chest is barrel-shaped and hyperresonant
3. Cough with scanty mucoid sputum occurs late after dyspnoea
4. Recurrent respiratory infections (not frequent)
5. Patients are called '**pink puffers**' as they remain well oxygenated
6. Features of right heart failure (cor pulmonale)



	Predominant Bronchitis	Predominant Emphysema
Age (yr.)	40-45	50-75
Dyspnea	Mild; late	Severe; early
Cough	Early; copious sputum	Late; scanty sputum
Infections	Common	Occasional
Respiratory insufficiency	Repeated	Terminal
Cor pulmonale	Common	Rare; terminal
Airway resistance	Increased	Normal or slightly increased
Elastic recoil	Normal	Low
Chest radiograph	Prominent vessels; large heart	Hyperinflation; small heart

Appearance	Blue bloater	Pink puffer



(15.05) COPD – ASTHMA

University Exams

Long Questions

1. Define obstructive disease of lung. Enumerate chronic obstructive pulmonary disease. Classify bronchial asthma. Discuss pathogenesis of atopic bronchial asthma. Add a note on the sputum findings in bronchial asthma.

Short Questions

1. Etiopathogenesis of Bronchial asthma
2. Describe gross and sputum microscopic features of Bronchial asthma
3. List differences between Extrinsic asthma and Intrinsic asthma

Overview

- Introduction
- Location
- Etiology
- Pathogenesis
- Types
- Gross
- Microscopy
- Clinical features

Introduction →

Asthma is a disease of airways that is characterised by
increased responsiveness of tracheobronchial tree to a variety of stimuli
↓
Widespread spasmodic narrowing of air passages
↓
Relieved spontaneously or by therapy





Location →

Trachea and Bronchi

Etiology →

Allergens

Pathogenesis

Asthma is an example of **type I hypersensitivity**

During first contact of the host with the antigen
↓
Antigen is captured by antigen presenting cells
↓
Presented to T cell which differentiates into TH2 cell
↓
TH2 cell releases IL-3, IL-4 and IL-5
↓
IL-4 causes activation of B cell
↓
Activated B cells differentiate to form IgE-secreting plasma cells
↓
IgE antibodies formed
↓
Bind to the Fc receptors present on mast cells
↓
Mast cells are now fully sensitised for the next event
↓
During the second contact with the same antigen (shocking dose)
↓
IgE antibodies on the surface of mast cells-basophils are activated
↓
Mast cell lysis
↓
Degranulation of mast cells-basophils
↓
Leukotrienes B4 and D4
↓
Bronchospasm
↓
Asthma



Types

Features	Extrinsic asthma	Intrinsic asthma
Pathogenesis	Type 1 hyper-sensitivity reaction due to exposure to an extrinsic antigen	Initiated by non-immune mechanisms with intrinsic body stimuli
Age on presentation	Child	Adult
Family history	Present	Absent
Prior allergic reaction/ allergen exposure	Positive history of rhinitis, urticaria, eczema	Absent
Serum IgE levels	Increased	Normal
Skin test	Positive	Negative
Associated COPD	Rare	Usually present
Examples	Atopic/allergic asthma, Occupational asthma, Allergic bronchopulmonary aspergillosis	Drugs (most commonly aspirin), viral infections, cold exposure, exercise induced asthma

Gross

- Lungs → **overdistended** due to over-inflation.
- Cut surface → occlusion of the bronchi and bronchioles by **viscid mucus plugs.**

Microscopy → Sputum



- Curschmann's spirals → Mucus plugs contain normal or degenerated respiratory epithelium forming twisted strips
- Charcot-Leyden crystals → Sputum contains numerous eosinophils and diamond-shaped crystals
- Creola bodies → Ciliated columnar cells sloughed from bronchial mucosa.

Clinical features

Triad →

1. Paroxysms of dyspnoea
2. Cough
3. Wheezing

(15.06) COPD – BRONCHIECTESIS

University Exams

Long Questions

1. Define bronchiectasis. Discuss the etiopathogenesis, gross and microscopy and clinical features of bronchiectasis.

Short Questions

1. Kartagener's syndrome
2. Define Bronchiectasis and write its Morphology
3. Describe morphology and complications of bronchiectasis.
4. Etiopathogenesis and Types of Bronchiectasis.

Overview

- Introduction
- Location
- Etiology
- Pathogenesis
- Types
- Gross
- Microscopy
- Clinical features
- Kartagener's syndrome

Introduction →

Abnormal and irreversible dilatation of the bronchi and bronchioles developing secondary to inflammatory weakening of the bronchial walls.

Location →

Bronchi and bronchioles

Etiology →

Obstruction → Infection → Inflammation



Pathogenesis

Endobronchial obstruction
↓
Resorption of air distal to obstruction
↓
Atelectasis
↓
Retention of secretions
↓
Promoting
bacterial growth→ Infection
↓
Inflammation
↓
weakening of bronchial walls
↓
Abnormal and irreversible dilatation of bronchi and bronchioles
↓
Bronchiectasis



Types

- **Cylindrical:** the most common type characterised by tube-like bronchial dilatation.
- **Fusiform**: having spindle-shaped bronchial dilatation.
- **Saccular**: having rounded sac-like bronchial distension.
- **Varicose**: having irregular bronchial enlargements.



Gross

- Cut surface → characteristic **honey-combed appearance**



Microscopy

1. Bronchi and bronchioles → **dilated**
2. Bronchial epithelium → **ulcerated or sloughed**
3. Bronchial wall → infiltration by **acute and chronic inflammatory cells**
4. In bronchial wall → **destruction of normal muscle and elastic tissue with replacement by fibrosis**

Clinical features

- Persistent cough with expectoration of copious amounts of foul-smelling, purulent sputum.

Kartagener's syndrome

Triad→

1. Bronchiectasis
2. Sinusitis
3. Situs invertus (including dextrocardia)



Feature	Chronic bronchitis	Emphysema	Bronchial asthma	Bronchiectasis
Introduction				
Location				
Etiology				
Pathogenesis				
Types				
Gross				
Microscopy				
Clinical features				

(15.07) PNEUMOCONIOSIS → ASBESTOSIS, SILICOSIS, BAGASSOSIS

University Exams

Long Questions

1. Define and classify pneumoconiosis. Discuss the pathogenesis, morphology and complications of asbestosis.

Short Questions

1. What is pneumoconiosis? Enlist pneumoconiosis.
2. Pathogenesis and Morphology of Silicosis.
3. Name different lesions associated with asbestosis.
4. Asbestos body.
5. Mention three tumors caused by asbestos.
6. Bagassosis.
7. Coal Workers' Pneumoconiosis

Overview

- Introduction
- Pathogenesis
- List of Pnemoconioses

1. Coal Workers' Pneumoconiosis
2. Silicosis
3. Asbestosis
4. Byssinosis
5. Bagassosis

} Introduction, Pathogenesis, Morphology, Clinical features

Introduction

- Non-neoplastic lung responses to inhaled aerosols or dust → including mineral dusts, organic dusts, fumes and vapors.
- **Caplan syndrome** → Coexistence of rheumatoid arthritis with pneumocconiosis.

Pathogenesis

- Pneumoconiosis development depends on

a) Amount of dust retained
b) Size of particle: most dangerous particle size are between 0.5 to 3 um because they can reach terminal alveoli.
c) Highly soluble particles will induce more toxicity
d) Additional effects of other irritants (e.g. cigarette smoking)

List of Pnemoconioses

Disease	Exposure source
Silicosis	Silica dust
Anthracosis	Coal dust
Asbestosis	Asbestos dust
Byssinosis	Cotton fibre
Bagassosis	Molasses (sugarcane)- Thermoactinomyces sacchari
Berylliosis	Beryllium
Farmer's Lung	Mouldy hay - Micropolyspora faeni
Siderosis	Iron dust
Stannosis	Tin dust
Bird fancier's lung	Avian/ bird droppings
Compost lung	Compost - Aspergillus

1) Coal Workers' Pneumoconiosis

Introduction →

- Parenchymal lung disease caused by inhalation of carbon particles (anthracotic pigment / coal mine dust).
- Not associated with risk of cancer or tuberculosis.

Pathogenesis →

Inhaled carbon pigment
↓
Engulfed by macrophages present in alveoli (Dust cells)
↓
Accumulate in connective tissue
↓
Carried by lymphatics to regional lymph nodes
↓
CWP

Morphology

Stages →

1. Pulmonary asymptomatic anthracosis
2. Simple CWP with little/no pulmonary dysfunction
3. Complicated CWP or progressive massive fibrosis—PMF

1) Pulmonary asymptomatic anthracosis

- Harmless in lungs of coal miners, persons residing in urban areas and tobacco smokers.



2) Simple CWP with little/no pulmonary dysfunction.

Gross:
- Coal macule: nonpalpable lesion (1–2 mm in diameter)
- Coal nodule: Larger (<2 cm) palpable, round or irregular and firm lesions.
- Both are mainly seen in the upper lobes

Microscopy:
- Coal macule: focal collections of many carbon-laden macrophages
- Coal nodule: carbon-laden macrophages and collagen and reticulin fibers.

3) Complicated CWP or progressive massive fibrosis—PMF

Gross →
- dark, multiple, black scars (> 2 cms)

Microscopy→
- carbon pigment, carbon-laden macrophages, dense collagen and disrupted lung's architecture.

Clinical features

PMF develops →
1. Pulmonary dysfunction
2. Pulmonary hypertension
3. Cor pulmonale

2) Silicosis

Introduction →

- Parenchymal lung disease associated with inhalation of crystalline silicon dioxide (silica).
- Most common chronic occupational disease in world.
- Involves upper lobes of the lung.
- Crystalline silicon dioxide → found in earth's crust

Susceptible individuals → Sandblasters, stone cutting, polishing and sharpening of metals, ceramic manufacturing, foundry work, tunneling through rock with high quartz content

Forms of silica → 2 forms

1. Crystalline (More toxic) → quartz, cristobalite, and tridymite
2. Amorphous forms.

Pathogenesis

Inhalation of silica particles
↓
Reach lung
↓
Interact with alveolar epithelial cells
↓
Ingested by alveolar macrophages
↓
Activates macrophages to release of mediators
(IL-1, TNF, fibronectin, oxygen-derived free radicals, fibrogenic cytokines)
↓
Stimulate fibroblast proliferation and collagen formation
↓
Silicosis

Morphology

2 forms →

1. Simple form
2. Progressive massive fibrosis

Gross:
Simple nodular silicosis →

- Tiny, barely palpable, discrete pale to blackened (2–4 mm) in the upper zones.
- Hilar lymph nodes → Periphery of lymph node → thin rim of calcification and radiographically → eggshell calcification



Progressive massive fibrosis →

- Expansion and coalescence of lesions
- Nodular masses > 2 cm diameter in a background of simple silicosis.



Microscopy

- Silicotic nodule → Whorled appearance →
    - Center → concentric layers of hyalinized acellular collagen surrounded by a dense cellular capsule of more condensed collagen.
    - Periphery → Aggregates of macrophages, lymphocytes and fibroblasts.



- Polarized light microscopy → birefringent silica particles in the center of silicotic nodule.

Clinical features

Progressive massive fibrosis →

1. Shortness of breath.
2. Increased susceptibility to lung infections, such as Mycobacterium tuberculosis, atypical mycobacteria and fungi.
3. Crystalline silica → carcinogenic (Increased risk of cancer)

- Silica: Increased risk of cancer of lung and tuberculosis.

3) Asbestosis and Asbestos Related Diseases

Introduction →

Asbestos-related diseases →
1. Asbestosis
2. Localized fibrous plaques
3. Pleural effusions
4. Carcinoma of lung
5. Malignant mesotheliomas
6. Laryngeal carcinomas.

Asbestosis →
- Interstitial fibrosis of the lung caused by exposure to asbestos dust.
- Does not include asbestos-induced pleural diseases and carcinoma of lung that are found in asbestos-exposed workers.

Asbestos Types → 2 forms
- Asbestos is a silica combine with magnesium, iron, calcium and aluminum
- Used in roofs, pipes, break lining

1. Serpentine form →

    - Chrysotile → used in industry
    - Long, curly, flexible structure
    - More soluble than amphiboles
    - Less readily carried to lungs and even if trapped in lungs → fragment into short particles → ingested by macrophages.

2. Amphibole form →

    - Crocidolite and amosite
    - Straight, rigid, brittle fibers that may align themselves in the airstream and remain stable in the lung.
    - Neither fragment nor soluble
    - Reach deeper into lungs → penetrate epithelial cells and reach the interstitium.
    - More pathogenic than chrysotiles

Pathogenesis

1. Fibrogenic Effect
2. Oncogenic Effect

1) Fibrogenic Effect

Both amphibole and serpentine forms → fibrogenic

Inhaled asbestos fibers
↓
Reach air spaces
↓
Macrophages ingest and clear fibers
↓
Activates macrophages
↓
Release chemotactic factors and fibrogenic mediators
↓
Interstitial fibrosis

2) Oncogenic Effect

Asbestos can act both as a tumor initiator and as promoter.

1. Reactive free radicals: produced by asbestos fibers → oncogenic.
2. Potentially toxic chemicals: Inhaled toxic chemical are adsorbed onto the asbestos fibers → oncogenic. (Eg. tadsorption of carcinogens in tobacco smoke onto asbestos fibers → increases the risk of lung carcinoma in asbestos workers.

Morphology

Gross →

- Sites: lower lobes and subpleura
- Fibrosis destroy normal architecture of lung → dilated airspaces (cystic spaces) surrounded by thick fibrous walls → honeycombed appearance



Microscopy →
Asbestos bodies



- Hallmark of asbestosis
- Consist of translucent center of asbestos fibers coated with a film of proteins rich in iron.
- Golden brown, fusiform or beaded rods.
- Coating is thickest at ends of fibers → dumb-bell appearance.
- Form when macrophages attempt to phagocytose asbestos fibers → iron derived from phagocyte ferritin.

Ferruginous body → Iron protein complex coating other inorganic particles, such as talc, mica and glass in the lung (not asbestos fiber).

Clinical Features

1. Dyspnea
2. Productive cough
3. Chest X-rays show irregular linear densities in both lower lobes.
4. As the disease progresses, a honeycomb pattern develops → may progress to respiratory failure, cor pulmonale, and death.

4) Byssinosis

Introduction →

Inhalation of cotton fiber dust over **> 10** years

Clinical features →

- Chronic cough
- Progressive Dyspnoea
- Ending in chronic bronchitis
- Affects about 10% of spinning workers

5) Bagassosis

Introduction →

Inhalation of bagasse (sugar cane) dust

Clinical features →

- Cough
- Breathlessness
- Haemoptysis
- Untreated may lead to diffuse fibrosis, emphysema and bronchiectasis

Feature	Coal Workers' Pneumoconiosis	Silicosis	Asbestosis	Byssinosis	Bagassosis
Introduction					
Pathogenesis					
Morphology					
Clinical features					

(15.08) PNEUMONIAS

University Exams

Long Questions

1. Describe stages of lobar pneumonia with gross and microscopy of each stage.
2. Classify pneumonia. Discuss in detail etiopathogenesis, pathology and complications of lobar pneumonia.

Short Questions

1. Write gross and microscopic appearance of various Stages of lobar pneumonia.
2. Difference between Lobar & Broncho pneumonia.
3. Etiopathogenesis of Lobar pneumonia.
4. Enumerate the stages in the evolution of lobar pneumonia. List the complications.
5. Morphology of Viral pneumonias.

Very Short Questions

1. Mention stages of lobar pneumonia.
2. Red hepatization – Gross and microscopy
3. Grey hepatization - Gross and microscopy
4. Microscopic appearance of lung in bronchopneumonia.
5. Pneumocystis carinii pneumonia.

Overview

- Definition
- Routes
- Pathogenesis
- Classification
    - Based on anatomic region of lung parenchyma
    - Based on etiology and pathogenesis
    - Based on clinical settings in which infection occurred

1. Bacterial Pneumonia
    a) Lobar
    b) Lobular
2. Viral Pneumonia
3. Fungal Pneumonia

} Introduction, Etiology, Gross, Microscopy

Definition

- Inflammation of lung parenchyma distal to terminal bronchioles (consisting of respiratory bronchiole, alveolar ducts, alveolar sacs and alveoli).
- **Consolidation of lung**



Routes

1. Inhalation → microbes present in the air.
2. Aspiration → organisms from nasopharynx or oropharynx.
3. Haematogenous spread → from a distant focus of infection.
4. Direct spread → from an adjoining site of infection.

Pathogenesis

Normal lung is free of bacteria because of presence of a number of lung **defense mechanisms** →

1. Nasopharyngeal filtering action
2. Mucociliary action of lower respiratory airways
3. Cough reflex
4. Presence of phagocytosing alveolar macrophages and immunoglobulins



Failure of these defense mechanisms

1. Altered consciousness → Oropharyngeal contents may be aspirated e.g. in coma, cranial trauma, seizures, cerebrovascular acci dents, drug overdose, alcoholism etc.
2. Depressed cough and glottic reflexes → e.g. in old age, pain from trauma or thoraco abdominal surgery, neuromuscular disease, kyphoscoliosis, endotracheal intubation and tracheostomy.
3. Impaired mucociliary transport → eg. Cigarette smoking, immotile cilia syndrome, inhalation of hot or corrosive gases
4. Impaired alveolar macrophage function → e.g. cigarette smoke, hypoxia, starvation, anaemia, pulmonary oedema and viral respiratory infections.
5. Endobronchial obstruction → from tumour, foreign body, cystic fibrosis and chronic bronchitis.
6. Immunocompromised states → congenital and acquired immunodeficiencies

Classification

Based on anatomic region of lung parenchyma

1. Lobar pneumonia
2. Bronchopneumonia (or Lobular pneumonia)
3. Interstitial pneumonia

Based on etiology and pathogenesis

1. Bacterial pneumonia
2. Viral pneumonia
3. Pneumonias from other etiologies

Based on clinical settings in which infection occurred

1. Community-acquire pneumonia
2. Nosocomial (Hospital acquired pneumonia)
3. Ventilator-associated pneumonia

Classification

1. Bacterial Pneumonia →
    - Lobar
    - Lobular (Bronchopnemonia)
2. Viral pneumonia → Intertitial / atypical pneumonia
3. Fungal pnemonia

1) Bacterial Pnemonia

a) Lobar pneumonia
b) Lobular pneumonia (Bronchopnemonia)





a) Lobar Pneumonia



Introduction →

- Acute bacterial infection of a part of a lobe, entire lobe, or even two lobes of one or both lungs
- Age → adults

Etiology

1. Streptococcus pneumonia → 90% of all lobar pneumonias
2. Staphylococcus aureus
3. Beta haemolytic streptococci
4. Gram-negative aerobic bacteria → Haemophilus influenzae, Klebsiella pneumonia (Friedlander's bacillus), Pseudomonas, Proteus and Escherichia coli, H. influenzae

Pathologic phases

1. Stage of congestion (initial phase)
2. Red hepatisation (early consolidation)
3. Grey hepatisation (late consolidation)
4. Resolution

1) Stage of Congestion: Initial Phase
Gross →
Enlarged, heavy, dark red and congested.

Microscopy →
- Dilatation and congestion of capillaries in alveolar walls.
- Pale eosinophilic oedema fluid in air spaces.
- Few RBCs and neutrophils in intra-alveolar fluid.
- Numerous bacteria demonstrated in alveolar fluid by Gram's staining



Red Hepatisation: Early Consolidation

Gross →
Cut surface → Red, firm and consolidated, airless, dry, granular → liver-like consistency

Microscopy →
- Oedema fluid of preceding stage is replaced by strands of fibrin.
- Marked cellular exudate of neutrophils and extravasation of RBCs.
- Alveolar septa are less prominent due to cellular exudation



Grey Hepatisation: Late Consolidation

Gross →
Cut surface → dry, granular and grey , liver-like consistency → liver-like consistency

Microscopy →
- Fibrin strands → dense and more numerous.
- Neutrophils and RBCs decreased
- Macrophages begin to appear
- Cellular exudate is separated from the septal walls by a thin clear space.





Resolution
Gross →
Solid fibrinous constituent is liquefied by enzymatic action→ restoring normal aeration in affected lobe.

Microscopy →
- Macrophages→ predominant cells
- Fragmented strands of fibrin in alveolar spaces
- Removal of cellular exudate from air spaces
- Restoration of normal parenchyma with aeration

b) Lobular Pneumonia / Bronchopneumonia



Intoduction →

- Infection of terminal bronchioles that extends into surrounding alveoli resulting in patchy consolidation of the lung.
- Age → Extremes of life (i.e. in infancy and old age)

Etiology

- Staphylococci
- Streptococci
- Pneumococci
- Klebsiella pneumoniae
- Haemophilus influenzae
- Gram negative bacilli like Pseudomonas and coliform bacteria

Gross

- Patchy areas of consolidation
- Mostly → bilaterally
- More common in → lower zones
- Cut surface → dry, granular, firm, red or grey in colour

Microscopy

- Acute bronchiolitis.
- Suppurative exudate → consisting chiefly of neutrophils in peribronchiolar alveoli.
- Thickening of alveolar septa by congested capillaries and leucocytic infiltration.
- Less involved alveoli contain oedema fluid.



Feature	Lobar Pneumonia	Bronchopneumonia
1. Definition	Acute bacterial infection of a part of a lobe of one or both lungs, or the entire lobe/s	Acute bacterial infection of the terminal bronchioles extending into adjoining alveoli
2. Age group	More common in adults	Commoner at extremes of age- infants and old age
3. Predisposing factors	More often affects healthy individuals	Pre-existing diseases e.g. chronic debility, terminal illness, flu, measles
4. Common etiologic agents	Pneumococci, Klebsiella pneumoniae, staphylococci, streptococci	Staphylococci, streptococci, Pseudomonas, Haemophilus influenzae
5. Pathologic features	Typical case passes through stages of congestion (1-2 days), early (2-4 days) and late consolidation (4-8 days), followed by resolution (1-3 weeks)	Patchy consolidation with central granularity, alveolar exudation, thickened septa
7. Prognosis	Better response to treatment, resolution common, prognosis good	Response to treatment variable, organisation may occur, prognosis poor
8. Complications	Less common; pleural effusion, empyema, lung abscess, organisation	Bronchiectasis may occur; other complications same as for lobar pneumonia

2) Viral / Interstitial / Primary Atypical Pneumonia

Introduction →

Patchy inflammatory changes → confined to interstitial tissue of lungs → without any alveolar exudate.

Also called as →

- Interstitial pneumonia → due to interstitial location of inflammation
- Primary atypical pneumonia → atypicality is absence of alveolar exudate (which is commonly present in other pneumonias)

Etiology

1. Respiratory syncytial virus (RSV) → most common

2. Others → Mycoplasma pneumonia, influenza and para influenza viruses, adenoviruses, rhinoviruses, coxsackieviruses and cytomegaloviruses (CMV).



Gross

Patchy to massive and widespread consolidation of one or both the lungs.
Lungs → heavy, congested and subcrepitant.

Microscopy

1. Hallmark → interstitial inflammatory reaction comprised by lymphocytes, macrophages and some plasma cells.
2. Thickening of alveolar walls due to congestion, oedema and interstitial inflammation
3. Lack of alveolar exudate
4. Necrotising bronchiolitis → foci of necrosis of bronchiolar epithelium





3) Fungal pnemonia

Introduction →

Almost 100% cases of **HIV/ AIDS** develop opportunistic infection during course of disease → most commonly Pneumocystis pneumonia.

Etiology

P. jirovecii → causes pneumonia in immunosuppressed people.

Gross

Lung → **consolidated, dry and grey.**

Microscopy

- Interstitial pneumonitis
- Thickening of alveolar walls
- No significant inflammatory exudate is seen in the air spaces.
- Alveolar lumina contain pink frothy fluid having organisms
- By Grocott's methenamine-silver (GMS) stain→ numerous tiny black dotlike organism P. jirovecii are demonstrable in the frothy fluid





Feature	Bacterial Pneumonia		Viral Pneumonia	Fungal Pneumonia
	Lobar	**Lobular**		
Introduction				
Etiology				
Gross				
Microscopy				

(15.09) PULMONARY TUBERCULOSIS

University Exams

Long Questions

1. Discuss Pathogenesis of Pulmonary Tuberculosis. Write its Clinical Manifestations and add a note on Laboratory Diagnosis
2. Write Mode of Transmission, Risk Factors, Pathogenesis and Clinical Manifestations of Pulmonary Tuberculosis

Short Questions

1. Risk Factors of Pulmonary tuberculosis.
2. Pulmonary and Extra-pulmonary Clinical Manifestations Risk Factors
3. Acid fastness
4. Type of Tubercles in Pulmonary tuberculosis.

Overview

- Introduction
- Morphology
- Acid fastness
- Classification
- Source of Infection
- Mode of Transmission
- Risk Factors
- Pathogenesis
- Tubercles types
- Clinical Manifestations → Pulmonary and Extra-pulmonary
- Specimen collection
- Laboratory Diagnosis

Introduction

- Robert Koch (1882) isolated tubercle bacillus and proved its causative role in tuberculosis
- Mycobacteria comprise of several species which produce various infections ranging from pulmonary, extrapulmonary, and cutaneous infections.

Morphology

- Weakly gram-positive
- Non-motile
- Non-sporing
- Non-capsulated
- Straight or slightly curved rod-shaped bacteria
- Obligate aerobes





Acid fastness

- Unique property of Mycobacteria → 'acid-fast', i.e. they resist decolorization by dilute mineral acids
- Acid fastness → due to presence of high content of mycolic acids in cell wall
- Important identification feature in diagnostic laboratories.

Classification

- M. tuberculosis complex: causes tuberculosis (TB) in man→
    - M. tuberculosis → most common species to cause TB in man
    - M. bovis → bovine tubercle bacillus
    - Others→ M. africanum, M. microti, M. caprae, M. canetti, M. suricattae, M. mungi
- M. leprae (Hansen's bacilli): causes leprosy
- Nontuberculous mycobacteria (NTM):
    - **Saprophytic** e.g. M. gordonae
    - **Commensal** eg. M. smegmatis in urine
    - **Opportunistic** e.g. M. kansasii.

Source of Infection

1. Human → e.g. pulmonary tuberculosis
2. Bovine → e.g. consumption of unpasteurized infected milk


Human

Bovine

Mode of Transmission

1. Airborne → Mainly transmitted by inhalation of aerosols (dry droplet nuclei <5 µm size), generated while coughing, sneezing, or speaking of infected patients.
2. Inoculation: Direct skin contact with an infected person
3. Ingestion: Swallowing of sputum (in infants) or consumption of unpasteurized (infected) milk.

Inoculation

Risk Factors

1. Sputum positive patients → transmit more efficiently than sputum negative patients
2. Bacillary load → 10^4 bacilli/mL in sputum is required for an effective transmission
3. Overcrowding → in poorly ventilated rooms.
4. Low cell mediated immunity→ eg. HIV- infected people
5. Other comorbid conditions → post-transplantation, gastrectomy, chronic renal failure, diabetes, IV drug abuse, smoking, etc.
6. Age: Elderly people are at increased risk due to waning immunity and underlying comorbidity

Pathogenesis

Inhalation of Droplet nuclei containing tubercle bacilli from infectious patients
↓
Mycobacterial surface lipoarabinomannan (LAM) binds macrophages
↓
Internalization of bacilli
↓
Phagocytosis by macrophages
↓
Macrophage try to destroy it
↓
Survival inside macrophages (due to bacterial cell wall LAM which impairs phagosomelysosome fusion)
↓
Macrophage fail to digest and degrade antigen
↓
Macrophages present antigen to TH1 cells lymphocytes
↓
TH1 cells get activated
↓
Release lymphokines
↓
IL-1, IL-2, interferon-Y, TNF-α



Release of Cytokines

- Interferon-γ → activates macrophages and transform it into epitheloid cells
- IL-1 and IL-2 → stimulate proliferation of more T cells.
- TNF-α → promotes fibroblast proliferation



Tubercles types

Hard tubercles:

- Tubercles are initially hard → composed of a central zone containing epithelioid and giant cells and a peripheral zone of lymphocytes and fibroblasts.

Soft tubercles:

- Later central part of the lesion undergoes caseous necrosis
- Growth of M. tuberculosis is inhibited within this necrotic environment because of low oxygen tension and low pH.



Clinical Manifestations

1. Pulmonary (60–90% of all cases of TB) →

    a) Primary type
    b) Secondary (Post primary) type

2. Extra-pulmonary (10–40% of all cases of TB)



Pulmonary Tuberculosis (EPTB)

Features	Primary PTB	Secondary PTB
Results due to	Initial exogenous infection with tubercle bacilli	➢ Exogenous reinfection ➢ Endogenous- reactivation of latent primary lesion
Age group affected	Children	Adults
Parts of lungs commonly affected	Sub pleural lesion affecting, upper part of lower lobe and lower part of upper lobe	Apical and posterior segments of the upper lobes (high oxygen tension)
Lesions formed at the initial sites	Fibrotic nodular lesions are formed (Ghon focus)	Hematogenous seedling in the apex of lungs called Simon's focus Reactivated Simon focus with central caseation (Assmann focus)
Lymph node	Ghon focus with associated hilar lymphadenopathy is common (called as primary complex)	Lymph node involvement is unusual
Clinical feature	It may be asymptomatic or may present with fever, productive cough (with or without hemoptysis) and occasionally chest pain, night sweating, weight loss	Lesions undergoing necrosis and tissue destruction, leading to cavity formation. Symptoms are similar, but more pronounced.



Extrapulmonary Tuberculosis (EPTB)

- Results from hematogenous dissemination of tubercle bacilli to various organs.
- Virtually all organ systems may be affected

Sites commonly involved (in order of frequency) are:

Tuberculous lymphadenitis:
- Most common form
- Most common sites are posterior cervical and supraclavicular lymph nodes.
- Presents as painless swelling in neck region

Pleural tuberculosis:
- Presents as pleural effusion.

Tuberculous lymphadenitis

Pleural tuberculosis

Tuberculosis of the upper airways:
- Involving larynx, pharynx, and epiglottis.
- Hoarseness, dysphonia, and chronic productive cough

Tuberculosis of the upper airways

Genitourinary tuberculosis

Genitourinary tuberculosis:
- Renal tuberculosis
- Genital tuberculosis:
    - In females→ fallopian tubes and endometrium → infertility.
    - In males→ epididymis most common site

Skeletal tuberculosis:
- Weight-bearing joints, such as spine (Pott's disease), hips and knees commonly affected.

Tuberculosis of CNS:
- Common in children.
- Tuberculous meningitis and tuberculoma are common


Skeletal tuberculosis

Tuberculosis of CNS

Gastrointestinal tuberculosis:
- Terminal ileum and cecum → most common involved.
- Route of spread may be due to swallowing of sputum with direct seeding, hematogenous spread, or ingestion of cow's milk infected with M. bovis


Gastrointestinal tuberculosis

Tuberculous pericarditis

Tuberculous pericarditis:
- Occurs as a direct extension from adjacent lymph nodes or following hematogenous spread

Tuberculous skin lesions
- Scrofuloderma:
  Skin condition caused by tuberculous involvement of skin by direct extension, usually from underlying tuberculous lymphadenitis
- Lupus vulgaris:
  Apple jelly nodules are formed over the face in females.

Scrofuloderma

Lupus vulgaris

Miliary or disseminated tuberculosis:
- More common in HIV-infected people
- Hematogenous spread of tubercle bacilli results in formation of yellowish 1–2 mm size granulomatous lesions resembling millet seeds (thus termed as miliary) in various organs.



Specimen collection

In pulmonary TB: Sputum (2 specimens—spot and early morning)

- Early morning sputum specimen should be collected in empty stomach, after rinsing mouth well
- Sputum should be at least 3-5 mL in quantity
- Thick and purulent (yellowish mucus).

Extrapulmonary specimens vary depending on site involved

Laboratory Diagnosis

1) Direct microscopy by acid-fast staining:

- The acid-fast stain was discovered by Ehrlich
- Subsequently modified by Ziehl and Neelsen

Interpretation

- Positive result: M. tuberculosis appears as long slender, beaded, less uniformly stained red colored acid-fast bacilli (AFB)
- Negative result: At least 100 oil immersion fields should be examined for 10–15 minutes before giving a negative report



2) Fluorescence Staining



Auramine-phenol solution as primary stain
↓
0.5% acid alcohol as decolorizer
↓
0.1% potassium permanganate as counter stain.
↓
slide is examined under fluorescent LED microscope
↓
Bacilli appear brilliant yellow against dark background

3) Culture Methods

Gold standard method of diagnosis of TB.

Advantages:
- More sensitive than microscopy with detection limit of 10–100 viable bacilli
- Indicates viability
- Drug susceptibility testing can be performed.

- Conventional Solid Media → Lowenstein-Jensen Medium
- Conventional liquid media → Middlebrook 7H9 medium.
- Automated Liquid Culture → BACTEC MGIT (Mycobacteria growth indicator tube)

4) Molecular Methods

- Polymerase Chain reaction (PCR)
- Automated real Time PCR → CBNAAT, TrueNat

Advantages →
- Take less time than culture
- More sensitive than culture.
- Detect the genes coding for drug resistance
- Used for epidemiological typing of strains.

(15.10) LUNG ABSCESS

University Exams

Short Questions

1. Describe etiology and morphology of Lung Abscess.
2. Discuss Causative organisms and clinical features of Lung abscess.

Very Short Questions

1. 3 etiologies of lung abscess.
2. Enumerate complications of lung abscess.

Overview

- Definition
- Causative organisms
- Causes
- Clinical features
- Complications

Definition

- An infection marked by localized suppurative necrosis of lung tissue.
- Characterized by accumulation of pus accompanied by the destruction of lung tissue.



Causative organisms

1. Anaerobes (Most common)
2. Gram-negative Bacterias
3. Staphylococci
4. Streptococci

Causes

1. Aspiration of infective material

- Most common cause of lung abscess → aspiration of oropharyngeal secretions.
- Most common site → right lung (more vertical right bronchus)

2. Antecedent primary bacterial infection
3. Septic emboli from infected thrombi or right sided endocarditis
4. Obstructive tumors (10 to 15% of abscesses)
5. Direct traumatic punctures or spread of infection from adjacent organs.

Clinical features

1. Cough
2. Fever
3. Large amounts of foul-smelling purulent or sanguineous sputum.
4. Chest pain

Complications

1. Extension into the pleural cavity
2. Hemorrhage
3. Septic embolization
4. Secondary amyloidosis

(15.11) LUNG CARCINOMA

University Exams

Long Questions

1. Classify lung tumors. Describe the etiology, pathology and clinical manifestations of bronchogenic carcinoma.
2. Classify lung tumors. Write important risk factors involved. Add a note on paraneoplastic syndrome.
3. Discuss classification of Lung Tumors. Discuss its etiopathogenesis, morphology and clinical features
4. Discuss etiopathogenesis of carcinoma lung and write the morphology of small cell carcinoma.

Short Questions

1. Etiological and clinicopathological features of bronchogenic carcinoma.
2. Histological Types and Clinical types of lung carcinoma
3. Etiopathogenesis of carcinoma lung.
4. Clinical presentation and morphology of small cell carcinoma of lung.
5. Gross and microscopic features of Oat cell carcinoma of lung.
6. Morphology of Adenocarcinoma of lung.
7. Gross and microscopic features of Squamous cell carcinoma of lung.
8. Enumerate paraneoplastic syndromes with appropriate examples.

Very Short Questions

1. Ennumerate non-small cell tumors of lung.
2. Histopathological variants of carcinoma of lung.
3. Microscopy of bronchioloalveolar carcinoma.
4. Spread of bronchogenic carcinoma.
5. List paraneoplastic syndromes associated with lung cancer.
6. Hormones elaborated by carcinoma lung.
7. Horner syndrome

Overview

- Introduction
- Age
- Gender
- Classification → Histological Types and Clinical types
- Etiology
- Molecular pathogenesis
- Gross

  1. Squamous cell carcinoma
  2. Adenocarcinoma
  3. Small cell carcinoma
  4. Large cell carcinoma

  } Introduction, Location, Microscopy, Marker

- Clinical features
- Spread
- Paraneoplastic syndrome
- Horner syndrome
- Prognosis

Introduction →

- Most common cause of cancer related death.
- Also known as Bronchogenic carcinoma

Age →

40 and 70 years

Gender →

- M > F
- There is a recent increase in females due to increased smoking among females.

Classification

- Histological Types
- Clinical types

Histological Types

1. Squamous cell or epidermoid carcinoma
2. Adenocarcinoma
3. Large cell carcinoma
4. Small cell carcinoma
5. Combined squamous cell carcinoma and adenocarcinoma (adenosquamous carcinoma).

Clinical types

1. Small cell carcinomas → SCC (20-25%)
2. Non-small cell carcinomas → NSCC (70-75%) (includes squamous cell carcinoma, adenocarcinoma, and large cell carcinoma)
3. Combined/ mixed patterns → 5-10%

Etiology



1. Smoking → 80% of the lung cancer occurs in active smokers
2. Other Factors →

    a) Radiation exposure
    b) Atmospheric pollution
    c) Occupational causes
    d) Dietary factors
    e) Chronic scarring

Molecular Pathogenesis

Squamous Cell Carcinoma →

- Inactivation of tumor suppressor gene → CDKN2A gene, P53 gene, RB gene
- Activation of oncogenes → FGFR1 and MET

Adenocarcinoma →

- Activation of Oncogenes → EGFR, BRAF, ERBB2/ HER2, ALK, MET, RET, KRAS gene

Small Cell Carcinoma →
- Inactivation of tumor suppressor gene → Loss-of-function mutations involving TP53, RB
- Activation of oncogenes → MYC family

Sequential epithelial changes in the respiratory tract in the form of →
1. Hyperplasia
2. Squamous metaplasia
3. Dysplasia
4. Carcinoma in situ
5. Carcinoma





Gross

1. Squamous cell carcinoma
2. Adenocarcinoma

3. Small cell carcinoma
4. Large cell carcinoma

1) Squamous cell carcinoma

Introduction →

- Most common type of lung cancer in India
- Most common type of lung cancer in men
- Most common type of lung cancer in smokers

Location →

Centrally (hilar)

Microscopy →

- Characterized by presence of keratinization and intracellular bridges
- Keratinization in form of squamous pearls or individual cells with markedly eosinophilic dense cytoplasm.



Marker →

Cytokeratin

2) Adenocarcinoma

Introduction →

- Most common type of lung cancer in World
- Most common type of lung cancer in women
- Most common type of lung cancer in Non – smokers

- Adenocarcinomas grow more slowly than squamous cell carcinomas but tends to metastatize widely and earlier

Location →

Peripheral carcinoma

Microscopy →

Various patterns →
1. Acinar → predominance of glandular structure
2. Papillary→ pronounced macropapillary configuration
3. Bronchioalveolar / Lepidic →

    - cuboidal to tall columnar and mucus-secreting epithelial cells growing along the existing alveoli
    - Bronchioloalveolar carcinoma consists mucin secreting bronchiolar cells and clara cells

4. Solid with mucin production

Marker →

CK 7



3) Small cell carcinoma (Oat-cell carcinoma)

Introduction →

- Have a strong relationship to smoking
- Most aggressive
- Most malignant lung tumor
- Worst prognosis
- Most common site of metastasis is brain
- Most common type of lung cancer responsive to chemotherapy and radiotherapy

Location →

Centrally (hilar)

Microscopy →

Tumour cells →

- Scant cytoplasm
- **Ill** defined cell borders
- Smaller than small resting lymphocytes
- High mitotic count
- Finely granular nuclear chromatin (salt and pepper pattern)
- Absent or incospicuous nucleoli
- Extensive Necrosis



Marker →

Neuroendocrine markers → chromogranin, Synaptophysin

4) Large cell carcinoma

Introduction →

- Large cell carcinoma is an undifferentiated malignant epithelial tumor of lung

Location →

Peripheral carcinoma

Microscopy →

- Characterized by cells with large nuclei, prominent nucleoli and a moderate amount of cytoplasm.



	Squamous cell carcinoma	Adenocarcinoma	Small cell carcinoma	Large cell carcinoma
Introduction				
Location				
Microscopy				
Marker				

Squamous cell carcinoma	Adenocarcinoma	Small cell cancer or oat cancer	Large cell cancer
• **NIC type of lung cancer in smokers.**	• Overall MC type of cancer.	• Associated with smoking	
• **NIC type in males.**	• NIC type in non-smokers and females.	• Commoner in smokers.	
• **Usually central in location (arise from the segmental bronchi)**	• Usually peripheral in location (arise from terminal bronchiole)	• Central in location	• Peripheral in location.
• **Shows highest frequency of p53 mutation.**	• Associated with K-ras mutation.	• Immunohistochemistry shows high expression of Bcl-2 gene in majority of tumors.	
• Intercellular bridges or junction is very specific. • **Histologically, the tumor has presence of keratinization.**	• Glandular pattern of growth of the tumor is seen. • Cells are positive for mucin and thyroid transcription factor-1 (TTF-1)	• Cells have scanty cytoplasm, small nucleoli, granular chromatin (salt and pepper pattern) • Azzopardl effect (Basophilic staining of vascular walls) is frequently present • Electron microscopy shows presence of neurosecretory granules chromogranin, synaptophysis and leu-7.	• Cells have large nuclei, prominent nucleoli and a moderate amount of cytoplasm.

Clinical Features

- MC symptoms → Cough
- Other symptoms are- Dyspnea, chest pain and hemoptysis (<50% only)
- Malignant Pleural effusion
- Clubbing and hypertrophic pulmonary osteoarthropathy (Resolve with excision of lung cancer)
- Hoarseness of voice (Recurrent laryngeal nerve involvement)
- Dysphagia (Esophagus Compression)
- SVC obstruction (MC caused by Small cell cancer)
- Small cell carcinoma- may develop Lambert-Eaton syndrome, a type of myopathy

Spread

1. Direct spread → Extension of cancer located at apex of lung into thoracic cage → Involve brachial plexus and the sympathetic chain causing pain and sensory disturbances, so called Pancoast's syndrome
2. Lymphatic spread → Initially→ hilar lymph nodes are affected. Later → mediastinal, cervical, supraclavicular and para-aortic lymph nodes.
3. Haematogenous spread →liver, adrenals, bones, pancreas, brain, opposite lung, kidneys and thyroid

Paraneoplastic syndromes in lung carcinoma

Hormones or hormone-like factors elaborated include →

1. ADH →Causing SIADH
2. ACTH → Causing cushing syndrome
3. PTH (parathormone related peptide → Causing hypercalcemia
4. Gonadotropins →causing gynaecomastia

Others
1. Lambert eaton syndrome
2. Acanthosis nigricons
3. Hypertrophic pulmonary oseoarthropathy (clubbing & periosteal reaction).

Small cell cancer	Squamous cell 'ca'	Large cell 'ca'	Adenocarcinoma
• Excess ADH. secretion causes SIADH and hyponatremia • Ectopic ACTH causes hypokalemia and Cushing syndrome • Myasthenic EatonLambert syndrome • Retinal blindness • Cerebellar • degeneration	• Ectopic PTH' and PTH related peptide production causes Hypercalcemia and hypophosphatemia • Clubbing - in all non small cell 'Ca'	• Gynecomastia	• Hypertrophic pulmonary osteoarthropathy • DIC • Migratory thrombophlebitis (Trousseau's syndrome)

Horner syndrome

- Caused due to compression of sympathetic nerve plexus by an apical tumor called as Pancoast tumor
- It is usually an **adenocarcinoma**.

Punjabi MEAL

Punjabi - **P**tosis
M - **M**iosis
E - **E**nophthalmos
A - **A**nhidrosis
L - **L**oss of ciliospinal reflex (in this reflex, pinching of the skin on the nape of neck causes dilatation of the pupil of the sameside)

Prognosis

Five years survival of each cancer:

- Squamous cell cancer → 35-50% (Best prognosis)
- Adenocarcinoma → 25-45%
- Adeno squamous → 20-35%
- Undifferentiated → 15-25%
- Small cell carcinoma → 0-5% (bad prognosis)

Remember

- Overall, most common type → Adenocarcinoma
- Most common type in India → Squamous cell carcinoma
- Most common type in non-smokers → Adenocarcinoma
- Most common type in smokers → Squamous cell carcinoma
- Most common type in females → Adenocarcinoma
- Most common type in males → Squamous cell carcinoma

M.C Variety of lung Cain India	→	Squamous Cell Ca
M.C Variety in world	→	Adenocarcinoma
M.C Variety in young Patients	→	Adenocarcinoma
Histological Variety that cavitates	→	Squamous cell Ca
M.C histological variety to metastasize	→	Small cell Ca
Ca which metastasize to opposite lung	→	Adenocarcinoma
Gynaecomastia is seen in which type of Ca	→	Large Cell Ca
Thrombophlebitis is seen with which Ca	→	Adenocarcinoma
Most common site for metastasis from ca lung	→	Brain
Lung Ca with worst prognosis	→	Small cell Ca
Variety most responsive to chemotherapy	→	Small cell Ca
Variety most responsive to radiotherapy	→	Small cell Ca
M.C Endocrine Gland to be involved by metastasis	→	Adrenal
Variety with best prognosis	→	Squamous cell Ca
Variety that is central in location	→	Squamous cell & small cell Ca
Variety that is peripheral in location	→	Adenocarcinoma
Most common variety with Cushing's syndrome	→	Small cell Ca
Most common variety with Hypercalcemia d/t PTH like substance Secretion	→	Squamous cell Ca

(16.01) BARRET ESOPHAGUS

University Exams

Short Questions

1. Reflux esophagitis.
2. Pathogenesis, Gross and microscopy Barrett's esophagus.

Overview

- Definition
- Incidence
- Age
- Gender
- Incidence of malignancy
- Gross
- Microscopy

Definition

Reflux of Gastric juice
↓
Reflux oesophagitis
↓
Replacement of stratified squamous epithelium of the lower oesophagus by columnar epithelium
↓
Columnar metaplasia (Intestinal metaplasia)
↓
Shifting of the oesophago-gastric junction upwards
↓
Barret oesophagus





Incidence→

Occur in ~10% of patients with symptomatic GERD

Age→

40 and 60 years of age.

Gender→

M > F

Incidence of malignancy

Barrett's oesophagus is a premalignant condition

Normal oesophagus
(stratified squamous epithelium)
↓
Barrett's epithelium (columnar metaplasia)
↓
Low grade Dysplasia (LGD)
↓
Highgrade Dysplasia (HGD)
↓
Carcinoma in situ
↓
Oesophageal adenocarcinoma

- Barrett's esophagus is single most important risk factor for adenocarcinoma of esophagus.
- Thereforesurveillance endoscopic biopsies are advised.



Gross

Affected area is **red and velvety.**

Microscopy

- Replacement of squamous epithelium by metaplastic columnar cells with goblet cells→ Intestinal metaplasia
- Dysplasia (low grade or high grade) may be seen in the metaplastic epithelium.



(16.02) ACHALASIA (CARDIOSPASM)

University Exams

Long Questions

1. **Define Achalasia. Discuss its Pathogenesis, Types and Symptoms**
2. Define Achalasia. Discuss its Triad, Morphologic features, Investigations and Complications

Short Questions

1. **Screening test and diagnostic test of Achalasia**
2. Etiopathogenesis of Achalasia

Overview

- Normal physiology
- Pathogenesis
- Types
- Symptoms
- Triad
- Morphologic features
- Investigations → Screening test and diagnostic test
- Complications



Normal

Normal physiology

- **Normally→**LES is closed
- **But during swallowing** → LES is relaxed due to →

    1. Release of nitric oxide
    2. Interruption of normal cholinergic signaling





Peristalsis

Pathogenesis





Types

1. Primary achalasia→ Congenital Myenteric plexusdamage
2. Secondary achalasia → Chagas' disease (sleeping sickness) → Trypanosoma cruzi infection → Destruction of myenteric plexus

Symptoms

1. Dysphagia → for both solids and liquids (in initial stages more for liquids than for solids)
2. Regurgitation
3. Weight loss
4. Recurrent respiratory tract infection due to spillage of liquids can also occur.
5. Retrosternal discomfort

Triad

1. Failure of relaxation of LES with swallowing
2. Increased resting tone of lower esophageal sphincter (LES)
3. Aperistalsis

Morphologic Features

1. Dilatation above the short-contracted terminal segment of the oesophagus
2. Muscularis propria is hypertrophied as a result of obstruction, or thinned out due to dilatation.
3. Secondary oesophagitis supervene and cause oesophageal ulceration and haematemesis

Investigations

Screening test
Diagnostic test

Screening test

Cholecystokinin (CCK) test:

- Normally→ CCK causes a fall in sphincter pressure (because of relaxant effect of inhibitory neurotransmitters like VIP and nitric oxide)
- Achalasia → CCK causes paradoxical increase in LES tone (loss of inhibitory neurons).



Diagnostic test

1. Barium swallow
2. Oesophagoscopy
3. Oesophageal manometry

1) Barium swallow

- Cucumber oesophagus / Bird beak' / rat tailed' appearance → Uniformly dilated oesophagus above, with a smooth tapering segment below
- Megaesophagus or sigmoid esophagus→ diffuse esophageal dilatation

2) Oesophagoscopy

- Dilated sac containing stagnant food and fluid due to stasis which splashes out with each heart beat and with each respiratory movement.
- LES→ closed→ **rosette appearance**

3) Oesophageal manometry

- Diagnostic procedure of choice
    - Hypertensive LOS
    - Does not relax on swallowing.
    - Aperistalsis in body of oesophagus



Complications

Precancerous condition→Leads toSquamous **cell carcinoma**

(16.03) ESOPHAGEAL LACERATIONS

University Exams

Very Short Questions

1. Pathogenesis and Mallory Weiss syndrome.
2. Types of Esophageal lacerations
3. Boerhave syndrome

Overview

- Mallory-weiss Syndrome→
  - Definition
  - Location
  - Pathogenesis
  - Age
  - Gender
  - Risk factor
- Boerhave syndrome →
  - Definition
  - Location

Mallory-weiss Syndrome

Definition →

Longitudinal Partial thickness tearin mucosa of gastroesophageal junction

Location →

At gastroesophageal junction

Pathogenesis

Bout of forceful retching (vomiting, coughing or straining)
↓
Longitudinal lacerations of mucosa of gastroesophageal junction
↓
Partial thickness Tear (extends through mucosa and submucosa, but not through muscular layer)
↓
Mallory-Weiss Syndrome



Age→

Older than **60 years.**

Gender →

M > F

Risk factor →

Majority of patients are **alcoholics**

Boerhave syndrome

Definition →

Tear is full thickness penetrating complete through all layers of esophageal wall with spillage of contents into mediastinum

Location →

Left posterolateral esophagus (3-5 cm above gastroesophageal junction).



(16.04) HIATUS HERNIA

University Exams

Short Questions

1. Define Hiatus Hernia. Write its etiology and types.
2. Morphologic Features of Sliding and Rolling Hiatus Hernia.

Overview

- Definition
- Etiology
- Types
- Morphologic Features
- Symptoms

Definition →

Herniation or protrusion of part of stomach through oesophageal hiatus of diaphragm.



Etiology

Basic defect →failure of muscle fibres of diaphragm that form margin of oesophageal hiatus.

1. Congenitally short oesophagus
2. Acquired→

    a) Degeneration of muscle due to ageing.
    b) Increased intra-abdominal pressure (pregnancy, abdominal tumours)
    c) Increase in fatty tissue in obese people causing decreased muscular elasticity of diaphragm

Types

1. Sliding or oesophago-gastrichernia
2. Rolling or para-oesophageal hernia
3. Mixed or transitional hernia

Morphologic Features

1) Sliding or oesophago-gastrichernia

Most common (85% of cases)



Sliding up on both sides of oesophagus
↓
Herniated part of stomach appears as supradiaphragmatic bell
↓
Cardio-oesophageal junction also moves upwards in thorax

2) Rolling or para-oesophageal hernia

10% of cases.



Cardiac end of stomach rolls up para-oesophageally
↓
Producing an intrathoracic sac
↓
Cardio-oesophageal junction is normal

3) Mixed or transitional hernia

- 5% cases
- **Combination of sliding and rolling hiatus hernia**



Symptoms

1. Heartburn (retrosternal burning sensation)
2. Regurgitation of gastric juice into mouth

(16.05) ESOPHAGEAL TUMOURS

University Exams

Long Questions

1. **Classify oesophageal carcinoma. Describe pathogenesis, gross and microscopic features of same. Discuss spread of oesophageal carcinoma.**

Short Questions

1. **Route of spread of carcinoma of Oesophagus**
2. **Enumerate morphological types of oesophageal carcinoma.**
3. **Describe morphology of oesophageal carcinoma.**
4. Describe gross and microscopic appearance of Squamous cell Carcinoma of Oesophagus

Overview

- Introduction
- Classification→

  1. Squamous cell Carcinoma
  2. Adenocarcinoma

  } Incidence, Age, Gender, Site, Risk factors, Gross, Microscopy

- Clinical features
- Spread
- Prognosis
- Overall, most common site

Introduction

- Carcinoma of oesophagus is **diagnosed late**
- Site → **Lower 1/3**

Classification

1. Squamous cell Carcinoma (epidermoid)
2. Adenocarcinoma
3. Adenosquamous Carcinoma

1) Squamous Cell (Epidermoid) Carcinoma

Incidence →

- Comprises 90% of primary oesophageal cancers.
- Most common type of esophageal carcinoma worldwide and in India

Age →

6th to 7th decades of life

Gender →

M > F

Site →

Middle 1/3> Lower 1/3> Upper 1/3

Risk factors

1. Alcohol
2. Cigarette smoking
3. Mucosal damage from physical agents →Hot tea, Lye ingestion, Radiation-induced strictures, Chronic achlasia
4. Ingested carcinogens →Nitrates, Smoked opiates, Fungal toxins in pickled vegetables
5. Plummer –Vinson / Paterson Kelly syndrome (Esophageal Web + glossitis + Iron deficiency).
6. Tylosis plamaris et plantaris (Congenital hyperkeratosis and pitting of palms and soles)
7. Dietary deficiencies of molybednum, Zinc, Vitamin A
8. Celiac sprue
9. Human papillomavirus (HPV) infection

Gross

3 patterns →

1. Polypoid fungating type (most common) → cauliflower-like friable mass protruding into the lumen.
2. Ulcerating type→ necrotic ulcer with everted edges
3. Diffuse infiltrating type → Annular, stenosing narrowing of lumen due to infiltration into wall of oesophagus.



Microscopy

- Tumour cell are arranged in sheets
- Tumour cells are large with hyperchromatic nuclei and high N/C ratio
- **Intracellular and extracellular keratinization** is seen
- **Keratin Pearls**commonly seen.



2) Adenocarcinoma

Incidence →

- Constitutes 10% of primary oesophageal cancer.
- Most common type of esophageal carcinoma in USA.

Age →

4th to 5th decades

Gender →

M > F

Site →

Lower 1/3>Middle 1/3

Risk factor

1. GERD leading o Barrett esophagus→ Single most important riskfactor
2. Obesity
3. Scleroderma
4. Diet deficient in fruits and vegetables
5. Diet rich in animal fat and cholesterol.

Gross

Nodular, elevated mass in lower oesophagus.

Microscopy

- Tumor cells are arranged in **glandular pattern**
- Tumour cells produce **mucin and form glands** with intestinal-type morphology
- Sometimes tumors are composed of diffusely infiltrative **signet-ring cells**





3) Adenosquamous Carcinoma

- Admixture of adenocarcinoma and squamous cell carcinoma.

Clinical features

Progressive dysphagia (more for solids as compared to liquids)
Weight loss
Chest pain and vomiting

Spread

- Local spread→ Into respiratory tree, aorta, mediastinumand pericardium.
- Lymphatic spread→ dependent on the anatomic site of primary tumor.

  a) Cancer in upper 1/3rd of the esophagus: metastasis to cervical lymph nodes.
  b) Cancer in middle 1/3rd of the esophagus: metastasis to paratracheal, mediastinal and tracheobronchial lymph nodes.
  c) Cancer in lower 1/3rd of the esophagus: metastasis to gastric and celiac lymph nodes.

- Hematogenous spread →to liver, lungs, adrenals, kidneys and bones.

Prognosis

- Dismal
- With standard methods of therapy (surgical resection and/or irradiation) → 70% of the patients die within one year of diagnosis.
- Five-year survival rate is 5-10%.

Overall, most common site

Squamous cell carcinoma →
Middle 1/3> Lower 1/3> Upper 1/3

Adenocarcinoma →
Lower 1/3>Middle 1/3

Overall esophageal carcinoma→
Lower 1/3



(16.06) PYLORIC STENOSIS

University Exams

Short Questions

1. What is Pyloric stenosis. Write its Types and Clinical features.

Overview

- Introduction
- Types
- Clinical features

Introduction

Hypertrophy as well as hyperplasia of the circular layer of muscularis in pyloric sphincter→ Narrowing of pyloric lumen



Types

1. **Infantile form**→predominantly in male children as a congenital defect
2. **Adult form** → due to inflammatory fibrosis or invasion by tumours.

Clinical Features

1. **Vomiting** →Projectile and occasionally contains bile or blood.
2. **Visible peristalsis** →noticed from left to right side of upper abdomen.
3. **Palpable lump** → better felt after an episode of vomiting.
4. **Loss of weight**

(16.07) GASTRITIS

University Exams

Long Questions

1. **What is the pathogenesis of Helicobacter pylori gastritis? Add a note on morphology and complications of it.**

Short Questions

1. **Describe classification of Gastritis. Write etiologic agents, pathogenesis and morphology of acute gastritis.**
2. **Stress gastritis**
3. **Cushing ulcer**
4. **Curling ulcer**
5. Describe Pathogenesis, Microscopy and Clinical features of type A and type B Chronic Gastritis

Overview

- Introduction
- Types

    1. Acute Gastritis→

        - Introduction
        - Etiology
        - Pathogenesis
        - Gross
        - Microscopy

        - Stress gastritis → Cushing ulcer, Curling ulcer

    2. Chronic gastritis→

        a) Type A
        b) Type B

        } Introduction, Site, Pathogenesis
        Microscopy, Clinical features

Introduction

Inflammation of gastric mucosa

Types

1. Acute
2. Chronic

1) Acute gastritis

Introduction →

- Acute mucosal injury
- Transientinjury
- Associated with neutrophilic infiltration.

Etiology

1. Heavy used of NSAIDs
2. Ischemia and shock
3. Alcohol
4. Smoking
5. Uremia
6. Nasogastric intubation
7. Severe stress (Trauma, burn, surgery)

Pathogenesis

When gastric mucosa protective mechanism is defective

Defensive Forces	Damaging Forces
• Surface mucus secretion. • Bicarbonate secretion into mucus. • Mucosal blood flow. • Surface membrane transport. • Epithelial regenerative capacity. • Elaboration of prostaglandins	• Increased acid production with back diffusion. • Decreased bicarbonate or mucin production. • Direct mucosal damage by alcohol and smoking. • Chronic use of nonsteroidal anti-inflammatory drugs (NSAIDs) reduces bicarbonate production and inhibits cytoprotective action of prostaglandins (prostaglandins inhibit acid production, promote mucin synthesis, and increase perfusion).

Gross

Gastric mucosa is oedematous with abundant **mucus and haemorrhagic spots.**



Microscopy

1. Oedema
2. Infiltration of lamina propria by neutrophils
3. Mucosa is sloughed off and there are haemorrhages onsurface.



Stress-related mucosal disease or Stress gastritis

Develops in association with severe stress→ severe trauma, extensive burns, intracranial disease, major surgery and any severe physiologic stress.

Curling Ulcers	Cushing Ulcers
• It is associated with burns. • Most common site is Duodenum > Stomach. • Multiple shallow and superficial erosions with less risk of perforation. • ↑ No synthetase and Endothelin → Vasoconstriction → Local ischemia	• It is associated with intracranial disease • Most common site is Cardia of Stomach. • Solitary deep ulcers with high risk of perforation. • Direct vagal stimulation → increased gastric secretion

2) Chronic gastritis

Chronic mucosal inflammatory changes→ mucosal atrophy and intestinal metaplasia in absence of erosions.



Types

a) Type A gastritis (Autoimmune gastritis)→ 10 % cases
b) Type B gastritis (H. pylori-related) → 90 % cases

a) Type A gastritis (Autoimmune gastritis)

Introduction →

- Also called **autoimmune gastritis** due topresence of circulating antibodies

Site

- Bodymucosa

Pathogenesis

Auto antibodies against parietal cells
↓
Depletion of parietal cells
↙ ↘
Gastric atrophy → Intestinal metaplasia → Decrease intrinsic f. → Pernicious anaemia
↓
- Hypoachlorhydria / Achlorhydria
↓
- Antral endocrine cell hyperplasia (G-cell hyperplasia)
- Hypergastrinaemia



Microscopy

- Rugalfolds are lost.
- Diffuse mucosal damage of acid-producing parietal cells in body and fundus.

Clinical Features

- Vitamin B12 deficiency can also manifest →
  - Pernicious anemia
  - Atrophic glossitis
  - Malabsorption
  - Peripheral neuropathy
  - Spinal cord lesions
  - Cerebral dysfunction
- Associated with **other autoimmune diseases** such as Hashimoto's thyroiditis and Addison's disease.

b) Type B gastritis (H. pylori-related)

Introduction →

- Also called **hypersecretory gastritis** due to excessive secretion of acid

Site

- Antral mucosa

Pathogenesis

H. pylori
↓
Irritates gastric mucosa
↓
Increased Gastric acid secreation
↓
Peptic ulcer



Microscopy

Initially →

- Intraepithelial neutrophils
- Subepithelial (superficial lamina propria) plasma cells.
- **H. pylori,** a spiral shaped bacteriais identified on the epithelial layer on luminal surface and does not invademucosa



Later →

- Mucosa atrophic
- Loss of parietal and chief cells.
- Accompanied by lymphoid aggregates with germinal centers.
- These lymphoid aggregates represent an induced form of mucosa-associated lymphoid tissue or MALT→ transform into lymphoma.



Special stains for H. pylori

Warthin Starry silver staining

Noninvasive Tests→Urea breath test

Patient drinks asolution of urea labeled with non-radioactive 13C
↓
If H. Pylori present
↓
Urease produced
↓
Urea is hydrolyzed
↓
Labeled CO2Produced
↓
Blows into a tube

Labeled CO2 Detected in breath samples by Breath analyser (mass spectroscopy)



Clinical features

1. Epigastric pain
2. Vomiting
3. Hematemesis
4. Malena

	Type A	Type B
Introduction		
Site		
Pathogenesis		
Microscopy		
Clinical features		

	Type A → Autoimmune	Type B → H. Pylori-Associated
Location	Body	Antrum
Inflammatory Infiltrate	Lymphocytes, macrophages	Neutrophils, subepithelial plasma cells
Acid production	Decreased	Increased to slightly decreased
Gastrin	increased	Normal to decreased
Serology	Antibodies to parietal cells (HMC-ATPase, Intrinsic factor)	Antibodies to H. pylori
Sequelae	Atrophy, pernicious anemia	Peptic ulcer, adenocarcinoma, MALToma

(16.08) PEPTIC ULCER

University Exams

Long Questions

1. Define peptic ulcer. Discuss the pathogenesis and morphology of peptic ulcer.
2. Describe etiology, gross, microscopy and clinical features of gastric ulcer. List the complications of gastric ulcer.
3. Describe etiopathogenesis, gross, microscopy and clinical features of benign duodenal ulcer.

Short Questions

1. Describe gross and microscopic features of peptic ulcer.
2. Enumerate various risk factors and describe pathogenesis of chronic peptic ulcer.
3. Role of H. pylori in gastric ulcer.
4. Acid peptic disease.
5. Etiology of peptic ulcer.
6. Differences between Gastric and Duodenal ulcer
7. Write briefly about clinical features and complications of peptic ulcer.

Very Short Questions

1. Macroscopy of benign and malignant gastric ulcer.
2. List complications of peptic ulcer.

Overview

- Definition
- Risk factors
- Etiology
- Types
- Site
- Gross
- Microscopy
- Clinical features
- Complications
- Differences between Gastric and Duodenal ulcer
- Investigations

Definition

- Ulcers → a breach in the mucosa of alimentry tract that extends through submucosa into mucularis mucasea or deep.
- Peptic ulcers → areas of degeneration and necrosis of gastrointestinal mucosa exposed to acid-peptic secretions.

Risk factors

1. Psychological stress
2. Physiological stress →
    - **Curling's ulcers** → Extensive burns (in the posterior aspect of first part of the duodenum).

- **Cushing's ulcers** → Intracranial lesions (developing from hyperacidity following excessive vagal stimulation).
- Shock
- Severe trauma
- Septicaemia
- Drug intake (e.g. aspirin, steroids, butazolidine, indomethacin).
- Local irritants (e.g. alcohol, smoking, coff ee etc).

Etiology

- Imbalance between →Mucosal Damaging factors/ Aggressive factors and Defense mechanisms



Increased Damage (Aggressive factors)

1. H. pylori infection → most common cause of PUD. It causes →
   a) Increased secretion of gastric acid
   b) Reduced duodenal secretion of bicarbonate.
2. H. pylori infection is associated with→
   - 90% of duodenal ulcers
   - 65% of gastric ulcers.
3. NSAIDs → It causes
   a) direct chemical irritation of mucosa
   b) suppresses mucosal prostaglandin synthesis
   c) reduces the bicarbonate secretion.
4. Smoking → Impairs blood flow to mucosa
5. Alcohol, radiation therapy and chemotherapy → cause direct injury to mucosal cells.
6. Ingestion of chemicals →cause direct injury.
7. Gastric hyperacidity
8. High-dose corticosteroids→ suppress prostaglandin synthesis and impair healing.

Impaired Defense

1. Ischemia→ Decreased oxygen delivery (e.g. at high altitudes and shock)
2. Shock
3. Delayed gastric emptying
4. Host factors →Reduced mucin synthesis in elderly increased susceptibility to gastritis.

Types

1. Gastric ulcer
2. Duodenal ulcers



Site

1. Duodenal ulcers (most common)→ first part of duodenum (anterior wall >posterior wall)
2. Gastric ulcers (2nd most common)→ lesser curvature in region of pyloric antrum (posterior wall > anterior wall)
3. Gastroesophageal junction of esophagus.
4. Anastomotic site → gastrojejunostomy
5. At metaplastic or heterotopic gastric mucosa→E.g. Meckel diverticulum within an ileum having ectopic gastric mucos



Gross

- Solitary ulcers.
- A sharply punched-out defect with overhanging mucosal borders and smooth, clean ulcer bases→ appear like a spoke wheel



Microscopy

4 histological zones →

1. **Necrotic zone→** in floor of ulcer
2. **Superficial exudative zone →** Composed of coagulative necrosis and inflammatory infiltrate
3. **Granulation tissue zone →** composed of nonspecific inflammatory infiltrate and proliferating capillaries.
4. **Zone of cicatrisation →** composed of densefibro collagenic scar



Clinical Features

1. Pain
   Gastric ulcer →

   - Epigastric pain occurs immediately after food
   - Never occurs at night.

   Duodenal ulcer →

   - Pain is relieved by food.
   - severe, occurs late at night (hunger pain)

2. Deep tenderness→

   - Gastric ulcer → midline of the epigastrium
   - Duodenal ulcer → right hypochondrium

3. Vomiting→

   - Gastric ulcer → Common → Vomiting relieves pain
   - Duodenal ulcer → rare

4. Haematemesis→

   - Gastric ulcer → Common
   - Duodenal ulcer → rare

5. Melaena→

   - Gastric ulcer → Rare
   - Duodenal ulcer → Common

6. Choice of Food →

   - Gastric ulcer →Choice bland diet, avoid spices
   - Duodenal ulcer →No particular choice

7. Appetite→

   - Gastric ulcer → good appetite but are afraid to eat
   - Duodenal ulcer → very good appetite

8. Weight→

   - Gastric ulcer → weight loss
   - Duodenal ulcer → weight gain (due to frequent ingestion of milk to avoid pain)

Complications

1) Haemorrhage/ Bleeding

- Most frequent complication.
- More common in posterior wall duodenal ulcers
- Severe bleeding may cause ‘coffee ground’ vomitus or melaena.
- Chronic blood loss may result in iron deficiency anaemia.



2) Perforation

- **Second most common complication** of peptic ulcer disease.
- More common in **anterior wall duodenal ulcers**
- Acute abdominal emergency.
- **Most common cause of death** in peptic ulcer

3) Malignant transformation

- Gastric ulcers→ <1% transform into carcinoma
- Duodenal ulcer → never turns malignant

Differences between Gastric and Duodenal ulcer

Features	Duodenal ulcer	Gastric ulcer
Site	1st part of duodenum	Along lesser curvature
Incidence	More common	Less common
Age	25 - 50 yrs, M>F	Beyond 6th decade, M>F
Pain after food intake	Relieved	Aggravated
Clinical features	Night pain and melena more common	No night pain, hematemesis more common
	No vomiting/no weight loss	Vomiting common/weight loss is present
Complications	No malignant change	Malignant change present (though rarely)

Investigations



- **Screening test:** Urea breath test (radiolabeled urea is broken down to radiolabeled CO2 by urease enzyme which is detected by breath analyzer, thus suggesting presence of H. pylori infection)
- **Gold standard:** Staining of H. pylori with **silver stain or Warthin starry stain**
- **Most specific investigation:** Culture of bacteria (done on Skirrow's medium)

(16.09) GASTRIC CARCINOMA

University Exams

Long Questions

1. A man developed gradual loss of weight, abdominal pain, anorexia, vomiting. A mass was detected in epigastric region. Stools were positive for occult blood. There is a firm lymph nodal mass in the left supraclavicular region and another nodule in the periumbilical region:

    a) What is your probable diagnosis?
    b) What is the etiopathogenesis and pathology of organ involved?

2. Classify gastric carcinoma. Describe etiopathogenesis, gross and microscopic features of same. Discuss the spread of gastric carcinoma.
3. Describe pathogenesis, gross & microscopic picture of carcinoma of stomach. Enumerate its complications.

Short Questions

1. Route of spread of carcinoma of stomach.
2. Enumerate Histologic types of gastric carcinoma.
3. Classification of Gastric carcinoma based on Depth of invasion
4. Describe gross and microscopic appearance of invasive Gastric carcinoma.
5. Early and Advance gastric carcinoma.

Very Short Questions

1. Risk factors for carcinoma stomach.
2. Linitis plastica.
3. Signet ring cell Gastric carcinoma

Overview

- Introduction
- Epidemiology
- Etiology
- Site
- Classification → Based on→

    1. Depth of invasion
    2. Macroscopic growth pattern
    3. Histologic subtype (Lauren classification)

- Clinical Features
- Complications
- Spread

Introduction →

- Adenocarcinoma ismost common malignancy of stomach
- Comprising >90% of all gastric cancers

Epidemiology →

- Age →4th to 6th decades of life
- Gender → M>F
- Geographical factors → More common in Japan
- Racial factors → Incidence is higher in Blacks, American Indians, Chinese

Etiology

- Factors that Increases risk
- Factors that decreases risk

Factors that Increases risk

Environmental factors	Genetic factors	Host factors
• H. pylori infection • Nitrites in diet • Nutritional (vitamin C, E) deficiency • Smoking	• Family history of gastric cancer • Blood group A • Hereditary non polyosis colon cancer syndrome (HNPCC) • Familial gastric cancer syndrome (due to E-cadherin mutation)	• Chronic gastritis • Intestinal metaplasia • Partial gastrectomy • Gastric adenoma • Barrett's esophagus • Menetrier disease

- H. pylori increases the risk of gastric carcinoma by 5 to 6-fold.

Chronic H. pylori infection
↓
Mucosal inflammation
↓
Decreases secretion of acid (hypochlorhydria) and pepsin
↓
Favors bacterial growth
↓
Continuation of chronic inflammation
↓
Progresses to mucosal atrophy
↓
Intestinal metaplasia
↓
Dysplasia
↓
Carcinoma

Factors that decreases risk

1. Aspirin
2. Diet rich in fresh fruit and vegetables
3. Vitamin A and C
4. Calcium, selenium, zinc and iron.

Site

- Pylorus and antrum> Body>Cardia> fundus
- Lesser curvature>Greater curvature
- Most favored site→lesser curvature of antropyloric region.



Classification

Based on→

1. Depth of invasion
2. Macroscopic growth pattern
3. Histologic subtype (Lauren classification)

1) Based on Depth of invasion



a) Early gastric carcinoma (EGC)→

- Cancer limited to the **mucosa and submucosa**
- Prognosis → good
- 5-year survival rate →93-99%

b) Advanced gastric carcinoma (AGC)→

- When carcinoma extended below the submucosa into muscularis propria or beyond
- Prognosis → poor
- 5-year survival rate → 5-15%

2) Based on macroscopic growth pattern

Early gastric carcinoma (EGC)→ 3 patterns

a) Type I (exophytic/polypoid/fungating)
b) Type II (flat or depressed) →

- Elevated (type IIa)
- Flat (type IIb)
- Depressed (type IIc)

c) Type III (excavated)→ Shallow or deeply erosive crater in wall ofstomach.



Advanced gastric carcinoma (AGC)→ 4 patterns

1. Polypoid
2. Fungating
3. Ulceratuve
4. Infilterative→Scirrhous carcinoma (Linitis plastica)→

- Stomach wall is thickened due to extensive desmoplasia
- Givingappearance as 'leather bottle stomach' or 'linitis plastica
- Lumen of the stomach is reduced.
- Worst prognosis.



3) Based on Histologic subtype (Lauren classification)

a) Intestinal type →

- Composed of neoplastic intestinal glands resmbling those of colonic adenocarcinoma.

b) Diffuse type→

- Composed of discohesive tumor cells which do not form glands
- Tumor cells contain abundant mucin which expand the cytoplasm and push the nucleus to the peripherya creating signet-ring cellappearance



Intestinal type



Diffuse type

Clinical Features

1. Persistent abdominal pain
2. Gastric distension and vomiting
3. Loss of weight (cachexia)
4. Loss of appetite (anorexia)
5. Anaemia, weakness, malaise.

Complication

Most common complication of gastric cancer →**haemorrhage** (in the form of haematemesis and/or melaena)

Spread

1. Local/direct spread→

- Spreads intomuscularis and serosa of stomach→ Invade into duodenum, pancreas, liver, colon and retroperitoneum.

2. Lymphatic spread →

    - Spreads to both regional and distant lymph nodes.
    - Pattern of lymph node involvement depends on the site of tumor

        - Supraclavicular sentinel→ Virchow node → may be the first clinical manifestation of an occult neoplasm → Troisier's sign
        - Left axillary lymph node → Irish node
        - Periumbilical lymph nodes →Sister Mary Joseph nodule

3. Transcoelomic spread→

    - Metastasis to the ovaries is called Krukenberg tumor.

4. Blood spread→

    - Spread viaportal vein into liver.
    - Other sites are lungs and bones.

(16.10) MECKEL'S DIVERTICULUM

University Exams

Short Questions

1. Discuss Meckel's diverticulum.
2. Embryologic origin and Morphology of Meckel's diverticulum.

Overview

- Introduction
- Site
- Embryologic origin
- **Rule of 2s**
- Morphology
- Complications

Introduction

- Meckel's diverticulum is the most common congenital anomaly of the gastrointestinal tract (2% of population)
- Meckel's diverticulum is a true diverticulum→an outpouching containing all the layers of intestinal wall in their normal orientation
- (True diverticulum →blind pouch leading off alimentary tract, lined by mucosa and including all layers of the bowel wall →mucosa, submucosa, and muscularis propria.)



Site

- Meckel's diverticulum commonly situated on →antimesenteric border of ileum, about 1-meter above ileocaecal valve.



Embryologic origin

- Embryologic origin of Meckel's diverticulum is from incomplete obliteration or persistence of vitello intestinal duct (connecting yolk sac and gut lumen)→ leaving a solitary out pouching within 85 cm of the ileocecal valve.





Rule of 2s

- Occur in about 2% of the population.
- Present within 2 feet ofileocecal valve.
- 2 inches long (varies from 2 to 15 cm).
- 2 times common in males than in females.
- Most often symptomatic by age 2 years

Morphology

- Always lined by small intestinal type of epithelium
- Rarely it may contain islands of gastric mucosa and ectopic pancreatic tissue.

Complications

1. Severe hemorrhage→Develops if the diverticulum contains acid-secreting gastric epithelium with peptic ulceration→painless bleeding per rectum.
2. Meckel's diverticulitis→ Develops secondary to peptic ulcerations or obstruction of the orifice of the diverticula→symptoms appear similar to acute appendicitis.
3. Intestinal obstruction →Results from intussusception, volvulus, adhesions or presence of a tumor or ectopic or enteroliths.

(16.11) INTUSSUSCEPTION

University Exams

Short Questions

1. What is Intussusception. Write its etiology and complications.

Overview

- Introduction
- Etiology
- Age
- Site
- Complications

Introduction

- Intussusception → Telescoping of a segment of intestine into segment below due to peristalsis.
- Intussusceptum→ Telescoped segment
- Intussuscipiens→ Lower receiving segment



Etiology

- In children → enlargement of the lymphoid tissue in the terminal ileum
- In adults→foreign bodies and tumours.

Age

More common in infants and young children

Site

Ileocaecal (portion of ileum invaginates into ascending colon)>Ileo-ileal>colo-colic



Complications

1. Intestinal obstruction
2. Infarction
3. Gangrene
4. Perforation
5. Peritonitis

(16.12) INFLAMMATORY BOWEL DISEASE (CROHN'S DISEASE, ULCERATIVE COLITIS)

University Exams

Long Questions

1. Classify Inflammatory Bowel disease. Discuss in details ulcerative colitis.
2. What are Inflammatory bowel diseases? Discuss in detail the etiopathogenesis, pathology and complications of Crohn's disease.
3. Mention inflammatory diseases of large intestine. Describe etiopathogenesis, gross, microscopy, clinical features and complications of ulcerative colitis.
4. Enumerate difference between Crohn's disease & ulcerative colitis.

Short Questions

1. Enumerate gross and microscopic differences of Crohn's disease and Ulcerative colitis.
2. Gross and microscopy of Crohn's disease.
3. Gross and microscopy of ulcerative colitis.

Overview

- Introduction
- Epidemiology
- Types

1. Crohn's disease
2. Ulcerative coli

} Introduction, Site, Gross, Microscopy, Clinical features, Complications, Mnemonics

Introduction

- Immune-mediated chronic intestinal inflammatory condition.
- Results from inappropriate mucosal immune activation.

Epidemiology

Age →
- Both UC and CD → 15 and 30 years of age.
- A second peak → 60 and 80 years.

Gender → M : F Ratio in →
- CD →1.8:1
- UC →1:1

Rural- Urban Prevalence→ urban >rural

Socioeconomic classes→ High>Low

Risk factors →
- Cigarette smoking
- OCP

Types

1. Crohn's Disease (CD)
2. Ulcerative Colitis (UC)

1) Crohn's Disease

Introduction →

- Chronic multifocal relapsing and remitting, progressive inflammatory bowel disease of unknown cause that can involve any portion ofgastrointestinal tract

Site

- Most common→terminal ileum>Ileocecal valve>cecum
- Can involve any portion of GIT



Gross

1. **Skip areas** → Multiple, well demarcated segmental bowel involvement with intervening uninvolved areas
2. Intestinal wall →
    - Fibrotic thickening and rubbery.
    - Lumen is narrowed
    - Appears like a hosepipe→ produces a characteristic radiological 'string sign'





3. Mucosa shows '**serpiginous ulcers.**
4. Intervening surviving mucosa is swollen giving '**cobblestone appearance'.**





5. Deep fissuring into bowel wall
6. Fissures can penetrate deeply through bowel wall → fistula or sinus tract formation
7. Bowel adhesions and serositis.
8. Strictures
9. Serosa studded with minute granulomas
10. Creeping fat→ When disease shows extensive transmural involvement →fat encircle around the antimesenteric serosal surface →



Microscopy

1. Patchy ulceration of mucosa→ Deep fissures
2. Transmural inflammatory cell infiltrate consisting of chronic inflammatory cells (lymphocytes, plasma cells and macrophages)
3. Non-caseating granulomas
4. Widening of submucosa due to oedema

Clinical Features

Intestinal Manifestations →

- Periods of active disease interrupted by asymptomatic periods.
- Intermittent attacks of mild diarrhea, fever and abdominal pain.
- Acute right lower quadrant pain
- Bloody diarrhea that may mimic acute appendicitis or bowel perforation.

Extraintestinal Manifestations →

- Uveitis
- Migratory polyarthritis
- Sacroiliitis,
- Ankylosing spondylitis
- Erythema nodosum
- Clubbing offinger

Complications

1. Iron-deficiency anemia
2. Malabsorption
3. Stricture formation
4. Fistula formation→between loops of intestine and surrounding structures such as urinary bladder, vagina and abdominal or perianal skin.
5. Perforation and hemorrhage.
6. Systemic amyloidosis→ rare

Mnemonic

SISTER IN CCF

- **S** - **S**kip lesions
- **I** - **I**eum is the most commonly involved
- **S** - **S**tring sign on radiology
- **T** - **T**ransmural inflammation
- **E** - **E**xtensive fibrosis
- **R** - **R**ectum usually spared
- **I** - **I**ntermittent attacks with colicky abdominal pain
- **N** - **N**oncaseating granuloma
- **C** - **C**obblestone appearance on mucosa
- **C** - **C**reeping fat on serosa
- **F** - **F**istula may develop

2) Ulcerative Colitis

Introduction →

- Severe, chronic crypt destructive, ulcerating inflammatory bowel disease of unknown cause.
- Inflammation involves only mucosa and submucosa of intestinal wall.
- Associated with exacerbations and remissions of bloody diarrhea.

Site

- Involves rectum and extends proximally for a variable distance to involve part or the entire colon.

Region Involved	Terminology
Entire colon	Pancolitis
Only the left-side of colon (without extension beyond transverse colon)	Left-sided disease
Limited to the rectum	Ulcerative proctitis
Disease process extends from rectum proximally towards the splenic flexure	Ulcerative proctosigmoiditis
Mild mucosal inflammation of the distal ileum	Backwash ileitis

Gross

1. **Continuous involvement** without any uninvolved skip areas
    - Barium enema shows: Garden hose appearance and Pipe stem appearance
2. Mucosa shows **superficial ulcers** (not penetrating muscular layer)
3. Intervening intact mucosa may form inflammatory '**pseudopolyps.**'







Microscopy

1. Superficial mucosal ulcerations
2. Inflammatory infiltrates of lymphocytes, plasma cells and macrophages.
3. Crypt distortion, cryptitis, crypt abscesses.
4. Basal plasmacytosis



Clinical features

Intestinal Manifestations →

- Exacerbations and remissions.
- Presents with attacks of bloody diarrhea with mucoid material, lower abdominal pain and cramps.

Extraintestinal Manifestations →

- Musculoskeletal → Migratory polyarthritis, Sacroiliitis, Ankylosing spondylitis.
- Eye → Uveitis
- Skin lesions →Erythema nodosum, Pyoderma gangrenosum.
- Liver disease: Primary sclerosing cholangitis and pericholangitis.
- DVT

Complications

1. Toxic megacolon→ In fulminant cases→inflammation damage muscularis propria →disturb neuromuscular function → colonic dilation → megacolonmay → perforation
2. Electrolyte imbalances.
3. Intestinal hemorrhage
4. Development of colorectal carcinoma→ Rare

Mnemonic

COLD TIP

- **C** - **C**ryptitis, crypt abscess
- **O** - **O**riginates in the rectum
- **L** - **L**imited to colonic mucosa and submucosa
- **D** - **D**istortion of crypts and depletion of mucin
- **T** - **T**oxic megacolon
- **I** - **I**ncreased risk of colonic cancer compared to Crohn
- **P** - **P**seudopylps.

	CD	UC
Introduction		
Site		
Gross		
Microscopy		
Clinical features		
Complications		
Mnemonics		

Feature	Crohn Disease	Ulcerative Colitis
Macroscopic		
Bowel region	Ileum ± colon	Colon only
Distribution	Skip lesions	Diffuse
Stricture	Yes	Rare
Wall appearance	Thick	Thin
Microscopic		
Inflammation	Transmural	Limited to mucosa
Pseudopolyps	Moderate	Marked
Ulcers	Deep, knife-like	Superficial, broad-based
Lymphoid reaction	Marked	Moderate
Fibrosis	Marked	Mild to none
Serositis	Marked	Mild to none
Granulomas	Yes (-35%)	No
Fistulae/sinuses	Yes	No

Clinical		
Perianal fistula	Yes (in colonic disease)	No
Fat/vitamin malabsorption	Yes	No
Malignant potential	With colonic involvement	Yes
Recurrence after surgery	Common	No
Toxic megacolon	No	Yes

(16.13) INFECTIOUS ENTEROCOLITIS (TYPHOID, TB, AMOEBIC ULCER)

University Exams

Long Questions

1. Enumerate Micro-organisms causing infective enterocolitis. Describe Pathogenesis, gross, microscopy and clinical features of intestinal tuberculosis.
2. Describe etiopathogenesis, gross, microscopic appearance and complications of enteric fever.

Short Questions

1. Gross, microscopy and complications of colon in amoebic colitis.
2. Difference between typhoid and tuberculous ulcers of intestines.
3. Intestinal tuberculosis → gross, microscopy and complications.

Very Short Questions

1. Typhoid ulcer of intestine
2. Amebic ulcer of intestine
3. Tuberculous ulcer of intestine

Overview

- Micro-organisms causing infective enterocolitis
- Important causes of infective enterocolitis

1. Typhoid Fever
2. Amebiasis
3. Intestinal Tuberculosis

} Causative agent, MOT, Incubation period, Pathogenesis, Morphology, Clinical features, Complications

Micro-organisms Causing Infective Enterocolitis

A. Bacterial Enterocolitis

1. **Entero-invasive Bacteria**

   i. **Tuberculosis**
   ii. **Salmonella**
   iii. **Campylobacter Jejuni**
   iv. **Shigella**
   v. **Escherichia Coli**
   vi. **Yersinia Enterocolitica**

2. **Enterotoxin-producing Bacteria**

   i. **Vibrio Cholerae**

B. Viral Enterocolitis

C. Fungal Enterocolitis

i. Candidiasis
ii. Mucormycosis

D. Protozoal and Metazoal Infestations

iii. Giardia Lamblia
iv. Entamoeba Histolytica
v. Balantidium Coli
vi. Taenia Solium
vii. Ascaris Lumbricoides
viii. Ancylostoma Duodenale
ix. Strongyloides Stercoralis

Important causes of infective enterocolitis

1. Typhoid Fever
2. Amebiasis
3. Intestinal Tuberculosis

1) Typhoid Fever

Causative agent →

Gram negative, motile bacilli →Salmonella typhi and Salmonella paratyphi.

MOT →

1. Ingestion of contaminated food and water.
2. Direct spread → Rare → by finger-to-mouth contact with feces, urine, or other secretions.

Incubation period →

7–14 days.

Pathogenesis

Salmonella
↓
Ingested through contaminated food or water
↓
Able to survive in gastric acid of stomach
↓
Reach mucosa of small intestine and penetrate ileal mucosa
↓
Reach lamina propria and submucosa
↓
Localized to Peyer's patches
↓
Cause inflammation, plateau-like elevations of Peyer's patches, necrosis → oval typhoid ulcers

Phagocytosed by macrophages
↓
Salmonella multiply within macrophages
↓
Carried to mesenteric lymph node via lymphatics
↓
Via thoracic duct
↓
Enter the bloodstream →transient bacteremia
↓
Seeded in RE system (liver, gallbladder, spleen, bone marrow)
↓
Disease clinically manifests



Morphology

- Site→ Most commonly involved → terminal ileum
- Typhoid ulcers →
    - Oval ulcers oriented along the long axis of bowel → Longitudinal ulcers (tuberculous ulcers of small intestine are transverse).
    - Base of ulcer → black due to sloughed mucosa.
    - Margins → Slightly raised



Clinical Features

Enteric fever →
- Anorexia
- Abdominal pain
- Bloating
- Nausea, vomiting
- Diarrhea.
- Fever →Continuous rise in temperature (step-ladder fever).
- Rose spots →small erythematous maculopapular lesions on skin that fade on pressure appear on chest and abdomen
- Spleenomegaly
- Hepatomegaly

Complications

Intestinal complications:
- Perforations of ulcer
- Hemorrhage from ulcer.

Extraintestinal complications:
- Encephalopathy
- Meningitis
- Seizures
- Endocarditis
- Myocarditis
- Pneumonia
- Cholecystitis

2) Amebiasis

Causative agent→

protozoa → Entamoeba histolytica

Exists in 3 forms→
1. Trophozoite
2. Pre-cystic stage
3. Cyst



MOT →

- Feco–oral route through ingestion of materials contaminated with human feces containing E. histolytica.
- Cyst is the infective form

Incubation period →

8–10 days

Pathogenesis

Ingestion of water and food containing cysts
↓
Cyst reaches caecum
↓
Excystation
↓
Produces metacystic trophozoites
↓
Lodge in submucous tissue of large intestine
↓
Flask-shaped ulcers
↓
Blood and mucus in faeces → Amoebic dysentery
↓
Some trophozoites enter into portal vein
↓
Carried away to liver
↓
Amoebic hepatitis (Extraintestinal amoebiasis)





Morphology



- Site→ mainly involves colon
- Amebic ulcers →
    - Flask-shaped ulcer with a narrow/ bottle neck and broad base resembling a flask
    - Base of ulcer → gray and necrotic
    - Trophozoites can be detected on surface of ulcer and in exudate.

Clinical Features

Intestinal amoebiasis → (Amoebic dysentery) →

- Ulcerative lesions in caecum
- Abdominal pain
- Profuse bloody diarrhoea
- Weight loss

Complications

Amebic Liver Abscess (Hepatic amoebiasis) →

- Posterio superior surface of right lobe of liver.
- Pus → red-brown anchovy sauce appearance.

Spread of amebic liver abscess →

- Local spread: expand and rupture through capsule of liver → directly spread into peritoneum, diaphragm, pleural cavity, lungs, or pericardium.
- Hematogenous spread: Rare → to brain and kidneys



3) Intestinal Tuberculosis

Causative agent →

Grampositive, acid fast bacilli → Mycobacterium tuberculosis, Mycobacterium bovis, Mycobacterium hominis.

MOT →

- Ingestion of cow's unpasteurized milk infected with Mycobacterium bovis.
- Swallowing of sputum in patients with active pulmonary tuberculosis.

Morphology

Site → Most common site →terminal ileum and ileocecal junction.

Three main forms of intestinal tuberculosis →
1. Primary
2. Secondary
3. Hyperplastic cecal tuberculosis



1) Primary Intestinal Tuberculosis

- Predominant changes are in regional mesenteric lymph nodes without any significant intestinal lesion
- Lymph nodes →enlarged, matted and caseous (tabes mesenterica) → may heal by fibrosis and calcification.

2) Secondary Intestinal Tuberculosis

Lesions of intestine are prominent than that of regional lymph nodes.



Intestinal ulcers→
- Large ulcers
- Transverse to long axis of bowel.
- Surface of ulcers may be coated with caseous material.
- Serosal surface may show visible tubercles
- Healed lesions may show transverse fibrous strictures and intestinal obstruction.



Transverse



Longitudinal

3) Hyperplastic Cecal Tuberculosis

- Variant of secondary tuberculosis.
- Lesions → cecum and ascending colon
- Thickening of wall and mucosal ulceration.
- Clinically →lesion present as a palpable and may be mistaken for carcinoma.



	Typhoid Fever	Amebiasis	Intestinal Tuberculosis
Causative agent			
MOT			
Incubation period			
Pathogenesis			
Morphology			
Clinical features			
Complications			

(16.14) MALABSORPTION SYNDROME (CELIAC SPRUE, TOPICAL SPRUE, WHIPPLE'S DISEASE)

University Exams

Short Questions

1. What is Malabsorption syndrome. Discuss pathogenesis and morphology of Celiac sprue.
2. Morphology of Tropical sprue.
3. Whipple's Disease → pathogenesis, morphology and clinical features

Overview

- Definition
- Classification
- Clinical Features
- **Important Types of MAS →**

1. Coeliac Sprue
2. Tropical Sprue
3. Whipple's Disease

} Introduction, Pathogenesis, Associations, Morphology, Treatment

Definition

Characterised by impaired intestinal absorption of nutrients → fat, proteins, carbohydrates, vitamins and minerals.

Classification

- Primary MAS→ due to primary deficiency of absorptive mucosal surface and associated enzymes
- Secondary MAS→ mucosal changes result secondary to other factors such as diseases, surgery, trauma and drugs.

I. Primary Malabsorption

1. Coeliac Sprue
2. Tropical Sprue
3. Whipple's Disease
4. Collagenous Sprue
5. Disaccharidase Deficiency
6. Allergic and Eosinophilic Gastroenteritis

II. Secondary Malabsorption

1. Impaired Digestion

   a) Mucosaldamage E.G. In Tuberculosis, Crohn's Disease, Lymphoma,
   b) Amyloidosis, Radiation Injury, Systemic Sclerosis
   c) Hepatic And Pancreatic Insufficiency
   d) Resection Of Bowel

2. Impaired Absorption

   a) Short Or Stagnant Bowel (Blind Loop Syndrome) From Surgery Or Disease Resulting In Abnormal Proliferation Of Microbial Flora
   b) Acute Infectious Enteritis
   c) Parasitoses E.G. Giardia, Strongyloides, Hookworms

3. Impaired Transport

   a) Lymphatic Obstruction E.G. In Lymphoma Tuberculosis, Lymphangiectasia
   b) Abetalipoproteinaemia

Clinical Features

1. Steatorrhoea → excessive fecal fat (pale, bulky, greasy, foul-smelling stools)
2. Chronic diarrhoea
3. Abdominal distension and Flatulence
4. Anorexia
5. Weight loss
6. Muscle wasting
7. Dehydration
8. Specific malnutrition deficiencies depending upon cause →

   - Anemia and mucositis (pyridoxine, folate, or B12)
   - Bleeding (vitamin K)
   - Osteopenia and tetany (calcium, magnesium, vitamin D)
   - Peripheral neuropathy (vitamin B12)

Important Types of MAS

1. Coeliac Sprue (Non-tropical Sprue, Gluten-Sensitive Enteropathy, Idiopathic Steatorrhoea)
2. Tropical Sprue (Environmental enteropathy)
3. Whipple's Disease (Intestinal Lipodystrophy)

1) Coeliac Sprue

Introduction →

- Most important cause of primary MASin temperate climates.
- Also known as →

  - Non-tropical Sprue
  - Gluten-Sensitive Enteropathy
  - Idiopathic Steatorrhoea

2 forms →

1. Childhood form →seen in infants and children →coeliac disease.
2. Adult form →seen in adolescents and early adult life →Idiopathic steatorrhoea.

Pathogenesis

- It is an immune-mediated diarrheal disorder

In Genetically predisposed individuals
↓
Gluten-containing foods
("BROW" - Barley, Rye, Oat and Wheat)
↓
Gliadin polypeptide
(resistant to digestive enzymes)
↓
Gluten-mediated delayed type hypersensitivity
Gliadine induces epithelial IL-15 expression
↓
CD4 T cell activation
↓
Cytokine-mediated epithelial
damage and Enterocyte apoptosis



Associations

- Immune disorders like type 1 diabetes, thyroid it is, and Sjogren syndrome, IgA Nephropathy
- Neurological disorders Like ataxia, autism, depression, epilepsy, Down syndrome, and Turner syndrome.

There is increased risk of:

- Enteropathy-associated T-cell lymphoma.
- Small intestinal adenocarcinoma.
- Squamous cell carcinoma of esophagus.

Morphology

- There may be partial villous atrophy or Subtotal villous atrophy
    - Partial villous atrophy → replacement of normal villous pattern by convolutions
    - Subtotal villous atrophy → flat mucosal surface.

- Lamina propria shows increased number of plasma cells and lymphocytes
- Intraepithelial CD8 T cells.
- Severity is greatest in more proximal intestine.



NORMAL VILLOUS | PARTIAL VILLOUS ATROPHY | SUBTOTAL VILLOUS ATROPHY

Remember

- Most Sensitive serologic test → IgA transglutaminase (tTg assay)
- Most Specific serologic test → IgA Endomysial antibodies (IgA EMA)

Treatment

- Elimi nation of gluten from the diet.

2) Tropical sprue

Introduction →

- Also known as → Environmental enteropathy
- Exclusively seen in people inhabiting or visiting tropical climates such as Caribbean countries, South India, Sri Lanka and Hong Kong.

Pathogenesis

Enterotoxin production by E. coli
↓
Intestinal injury

Morphology

- Similar to celiac disease →Partial villous atrophy or Subtotal villous atrophy
- Distal small bowel is most severely affected.
- but distal small bowel is most severely affected.
- partial villous atrophy or Subtotal villous atrophy

Treatment

Relieved by →
  - Removal of the patient from tropical area
  - Oral administration of antibiotics

- Gluten-free diet has no role in improvement.

3) Whipple's Disease

Introduction→

Also known as →Intestinal Lipodystrophy

Pathogenesis

Caused by Gram-positive actinomycete→ Tropheryma whippleie

Associations

- Migratory polyarthritis
- Neurological disturbances
- Focal hyperpigmentation of the skin.

Morphology

- Small intestine mucosa will show following hallmark features:
    - Macrophages in lamina propria showing PAS positive, diastase resistant granule
    - Rod -shaped bacilli within macrophages
- Similarly laden macrophages are present in lymphatics, lymph nodes, joints, and the brain.
- Active inflammation is largely absent.



Treatment

- Oral antibiotic therapy

	Coeliac Sprue	Tropical Sprue	Whipple's Disease
Introduction			
Pathogenesis			
Associations			
Morphology			
Treatment			

(16.15) CARCINOID TUMOURS

University Exams

Short Questions

1. Enlist the symptoms of carcinoid syndrome
2. Gross and microscopic features of carcinoid tumor of GIT.
3. Type of Carcinoid tumor of GIT.

Overview

- Introduction
- Age
- Types
- Gross
- Microscopy
- IHC
- Clinical Features
- Prognosis

Introduction →

- Tumours originating from diffusely distributed endocrine cells (argentaffin cells =Kulchitsky cells =entero chromaffin cells)→ Also known as Argentaffinoma
- Belonging to APUD cell system → also called as apudomas

Age →

Peak incidence is in the sixth decade.

Types

Based on embryologic derivation of tissues where tumour is located→

- Midgut carcinoids (most common → 60-80%) →seen in terminal ileum and appendix
- Hindgut carcinoids (10-20%) →occurring in rectum and colon
- Foregut carcinoids (10%)→located in the stomach, duodenum and oesophagus
- Other → bronchus, trachea, gallbladder, and Meckel's diverticulum.

Gross

- Small, button-like submucosal elevations with intact or ulcerated overlying mucosa.
- Cut section →yellow tan or brown (due to presence of Lippofuscin pigment)
- Firm (due to intense desmoplastic response)



Microscopy

- Well-differentiated neuroendocrine carcinomas.
- Tumor cells arranged in sheets→ palisading of peripheral cells

Tumor cells →

- Small, monotonous
- Uniform cohesive cells
- Poorly-defined cell boundaries
- Scant granular cytoplasm
- Oval, stippled nuclei.
- Mitotic figures → rare.





IHC

Tumor cells are positive for neuroendocrine markers →

- Chromogranin
- Synaptophysin

Clinical Features

Carcinoid tumours that metastasise to liver→ 5-HT and5-HIAA production→Carcinoid syndrome

1. Intermittent attacks of flushing of skin of face
2. Episodes of watery diarrhoea
3. Abdominal pain
4. Attacks of dyspnoea due to bronchospasm
5. Right-sided heart failure due to involvement of tricuspid and pulmonary valves and endocardium

Dietary tryptophan
↓
Hydroxylated to 5-hydroxy tryptophan
↓
Decarboxylated to 5-HT
↓
Oxidised to 5-HIAA by the monoamine oxidase in the liver cells
↓
Excreted in the urine



- Midgut carcinoids → rich decarboxylating enzymes → produce large quantities of 5-HT and 5-HIA →high frequency of carcinoid syndrome
- Foregut and hindgut carcinoids → lack decarboxylating enzymes → less associated with carcinoid syndrome.

Prognosis

Most important prognostic factor for gastrointestinal carcinoid is primary site of the tumor →

- Foregut tumors → rarely metastasize →cured by resection.
- Midgut carcinoids → aggressive
- Hindgut tumors → found only incidentally.
- Appendiceal carcinoids → found attip and benign
- Colonic carcinoids →large and metastasize.

(16.16) APPENDICITIS

University Exams

Short Questions

1. Write gross, microscopy and laboratory findings in a case of Acute appendicitis.
2. Clinical features and complications of acute appendicitis.
3. McBurney's sign in Acute appendicitis.

Overview

- Introduction
- Age
- Etiopathogenesis
- Causes
- Gross
- Microscopy
- Symptoms
- Signs
- Laboratory findings
- Complications

Introduction →

- Acute inflammation of appendix
- Most common acute abdominal condition confronting surgeon.

Age →

- More commonly in older children and young adults
- Uncommon at extremes of age.

Etiopathogenesis

Obstruction of lumen from various etiologic factors
↓
Increased intraluminal pressure
↓
Presses upon blood vessels
↓
Produce ischaemic injury
↓
Favours bacterial proliferation
↓
Acute appendicitis

Causes

A. Obstructive:

1. **Faecolith**
2. **Calculi**
3. **Foreign body**
4. **Tumour**
5. **Worms (Enterobius vermicularis)**
6. **Diffuse lymphoid**
7. hyperplasia

B. non-obstructive:

1. **Haematogenous spread of generalised infection**
2. **Vascular occlusion**
3. Inappropriate diet lacking roughage.

Gross

- Early acute appendicitis→organ is swollen and serosa shows hyperaemia.
- Well-developed appendicitis→serosa is coated with fibrinopurulent exudate and engorged vessels on surface.
- Advanced cases → acute gangrenous appendicitis→necrosis and ulcerations of mucosa which extend through wall→ surface is coated with greenish-black gangrenous necrosis



Microscopy

Histological criterion →neutrophilic infiltration of muscularis.

- Congestion
- Oedema of appendiceal wall.
- Mucosa sloughed off
- Neutrophilic abscesses in wall
- Impacted foreign body, faecolith, or concretion may be seen in the lumen



Symptoms

1. Colicky pain, initially around umbilicus but later localised to right iliac fossa
2. Nausea and vomiting
3. Pyrexia of mild grade
4. Abdominal tenderness
5. Increased pulse rate

Signs

- McBurney's sign→ A classic physical finding is deep tenderness located at McBurney's point
- McBurney's point→ two-thirds of distance from umbilicus to right anterior superior iliac spine

McBurney's sign



McBurney's point

Laboratory findings

1. Raised WBC count
2. Neutrophilic leucocytosis with toxic granules in neutrophils
3. Microscopic hematuria

Complications

1. Peritonitis → due to perforated appendix
2. Appendix abscess → due to rupture of an appendix
3. Adhesions → fibrous adhesions to the greater omentum, small intestine and other abdominal structures.
4. Portal pylephlebitis → Spread of infection into mesenteric veins
5. Mucocele→Distension of distal appendix by mucus following recovery

(16.17) HIRSCHSPRUNG'S DISEASE

University Exams

Short Questions

1. Write Pathogenesis and Clinical Features of Hirschsprung disease.
2. Discuss Screening test, Gross and Microscopy of Hirschsprung disease.

Overview

- Introduction
- Gender
- Incidence
- Site
- Pathogenesis
- Clinical Features
- Screening test
- Gross
- Microscopy

Introduction →

- Also known as congenital aganglionic megacolon
- Results from arrested migration of neural crest cells into gut resulting in formation of aganglionic segment lacking peristaltic contractions.

Gender →

M > F

Incidence →

1 in 5000 live births

Site →

- Rectum →always affected
- Proximal involvement →variable

Pathogenesis

Heterozygous mutations in RET tyrosine kinase receptor
↓
Arrested migration of neural crest cells into gut
↓
Affected segment lacks both Meissner and Auerbach plexuses
↓
Formation of aganglionic segment
↓
Spastic contraction with Peristaltic contractions absent
↓
Functional obstruction
↓

Progressive dilation & hypertrophy of
unaffected proximal colon
↓
Hirschsprung disease

Clinical Features

- Neonatal failure to pass meconlum
- Abdominal distention
- Severely distended megacolon (up to 20 cm in diameter)

Screening test→

Anorectal manometry.

Gross →

Typical case of Hirschsprung's disease
shows 2 segments→

- A distal narrow segment that is aganglionic
- A dilated proximal segment that contains normal number of ganglion cells



Microscopy →

Rectal biopsy →

- Complete absence of submucosal and myenteric ganglion cells
- Submucosal and myenteric nerve hypertrophy in aganglionic rectum



(16.18) COLORECTAL CARCINOMA

University Exams

Long Questions

1. Classify tumors of large intestine. Discuss etiopathogenesis, pathology and staging of carcinoma of the colon. List the precancerous conditions of carcinoma colon.
2. A 72-year-old male presented with changes in the bowel habit, tarry stool, and loss of weight, fatigue and weakness of 6 months duration. After an endoscopic biopsy of colon, left-sided hemicolectomy was done. What is the probable diagnosis? Describe the etiopathogenesis and morphology of the target organ involved.
3. Discuss in detail the pathogenesis of colorectal carcinoma, emphasizing the morphologic and molecular changes. Write briefly about the morphology of the same.
4. Describe etiology, gross, microscopic findings, clinical features, complications and spread of carcinoma colon.

Short Questions

1. Etiopathogenesis of colonic carcinoma.
2. Genetic Basis of Colorectal Carcinogenesis
3. Adenoma-carcinoma sequence in carcinoma colon.
4. Differences between carcinoma of colon on right and left side.
5. Astler-Coller staging of colonic carcinoma.

Overview

- Introduction
- Age
- Gender
- Etiology
- Genetic Basis of Colorectal Carcinogenesis
- Sites
- Gross
- Microscopy
- Clinical Features
- Complications
- Diagnosis
- Staging
- Prognosis

Introduction →

Comprises 98% of all malignant tumours of large intestine.

Age →

Average age of patients is about 60 years.

Gender →

M > F

Etiology

1. **Geographic variations →**
   - More common in North America and Northern Europe than in South America, Africa and Asia.

2. **Dietary factors→**
   - A low intake of vegetable fibre-diet → low stool bulk
   - Consumption of large amounts of fatty foods
   - Excessive consumption of refined carbohydrates → remain in contact with colonic mucosa for prolonged duration→ changes bacterial flora of bowel → production of carcinogenic substances.

3. **Adenoma-carcinoma sequence →**
   - Colonic adenocarcinoma evolves from pre-existing adenomas
   - Hyperplasia → dysplasia → carcinoma in situ → invasive carcinoma.

4. **Hereditary non-polyposis colonic cancer (HNPCC or Lynch syndrome) →**
   - Autosomal dominant condition associated with multiple primary cancers at different sites (endometrium, ovary, colorectal cancer)
   - Due to germline mutations in mismatch repair genes, human mutL homolog abbreviated as hMLH2 located on chromosome 2 and hMLH1 on chromosome 3 resulting in DNA instability.
   - In HNPCC, colorectal cancer is seen in at least two generations of first-degree relatives, occurs at a relatively early age (< 50 years), located more often in proximal colon, and has better prognosis than sporadic colon cancer cases.

Genetic Basis of Colorectal Carcinogenesis
APC mutation / β-catenin mechanism

- Loss of APC gene located on the long arm of chromosome 5 (5q) in 80% cases of sporadic colon cancer. Since the function of APC gene is linked to β-catenin, loss of APC gene results in translocation of β-catenin to the nucleus where it activates transcription of other genes, mainly MYC and cyclin D1, both of which stimulate cell proliferation.
- Point mutation in K-RAS gene follows loss of APC gene and is seen in 10 to 50% cases
- Deletion of DCC gene located on long arm of chromosome 18 i.e. 18q in 60-70% cases
- Loss of p53 gene seen in 70-80% cases



Sites

- Rectum (60%)
- Sigmoid and descending colon (25%)
- Caecum and ileocaecal valve (10%)
- Ascending colon, hepatic and splenic flexures (5%)
- Transverse colon (Uncommon)



Gross

Right-sided colonic growths →

- fungating polypoid carcinoma → tend to be large, cauliflower-like, soft and friable masses projecting into the lumen
- Liquid nature of contents in ascending colon leaving space for luminal growth on right side

Left-sided colonic growth →

- carcinomatous ulcers→ napkin-ring configuration i.e. they encircle bowel wall circumferentially with increased fibrous tissue forming annular ring, and have central ulceration on surface
- Contents in left colon are more solid permitting spread of growth into the bowel wall.



Microscopy

- % of colorectal carcinomas →adenocarcinomas
- Tumors show glands of variable size and configuration separated by moderate amount of stroma.
- Lumen of the glands →filled with inspissated eosinophilic mucus and nuclear and cellular debris
- Mucinous adenocarcinomas →secrete abundant mucin and accumulate within the intestinal wall.
- Signet-ring carcinoma → consists of signet-ring cells



Moderately differentiated adenocarcinoma | Mucin-secreting adenocarcinoma | Signet ring adenocarcinoma

Clinical Features

1. Occult bleeding (melaena)
2. Change in bowel habits, more often in left-sided growth
3. Loss of weight (cachexia)
4. Loss of appetite (anorexia)
5. Anaemia, weakness, malaise.

Complications

1. Obstruction
2. Haemorrhage
3. Perforation
4. Secondary infection

Diagnosis

1. Stool test for occult blood
2. PR examination
3. Proctoscopy
4. Radiographic contrast studies and CT scan
5. Tumour markers→ carcinoembryonic antigen (CEA)

    - Elevated in 100% cases of metastatic colorectal cancers.
    - Have prognostic significance only and is not diagnostic of colorectal cancer because it is positive in other cancers too such as of lungs, breast, ovary, urinary bladder and prostate.

Staging

3 staging systems are in use →

- Dukes' ABC staging (modified Duke's includes stage D as well).
- Astler-Coller staging which is a further modification of Duke's staging and is most widely used.
- TNM staging described by American Joint Committee is also used.

Duke's system as modified by Astler-Coller

Stage		TNM	5-year Survival	Pathologic Features
A	**I**	T_1 N0 M0	>90%	Cancer confined to mucosa only
B_1	**II**	T_2 N0 M0	85%	Cancer extends into submucosa
B_2	**II**	T_3 N0 M0	70-80%	Cancer extends into muscularis±serosa
C_1	**III**	Tx N_1 M0	35-65%	Cancer involves muscularis+regional lymph nodes
C_2	**III**	Tx N_1 M0	20-35%	Cancer extends into serosa+regional lymph nodes
D	**IV**	Tx Nx M_1	5%	Cancer with distant metastases



Prognosis

Depends upon →
1. Extent of bowel involvement
2. Presence or absence of metastases
3. Histologic grade of tumour
4. Location of tumour

(17.01) JAUNDICE

University Exams

Long Questions

1. Define jaundice. Describe Laboratory diagnosis of jaundice following Infective hepatitis.
2. Define jaundice. Describe the role of laboratory in diagnosis of obstructive jaundice.
3. Define and classify jaundice. Discuss laboratory findings which differentiate them.
4. Discuss pathophysiology of Jaundice. Give an account of laboratory tests for its differential diagnosis.
5. Enumerate types of jaundice and write utility of liver function tests in it.

Short Questions

1. Classification of jaundice.
2. Laboratory investigations in jaundice.
3. Describe investigation in Hepatocellular jaundice.
4. Describe investigation in Haemolytic jaundice.
5. Urine in obstructive jaundice.
6. Tabulate differences in liver function tests in hemolytic, hepatocellular and obstructive jaundice.
7. Intrahepatic versus Extrahepatic Cholestasis

Very Short Questions

1. Tabulate differences between prehepatic, hepatic and post hepatic jaundice.
2. Tabulate differences between Unconjugated and Conjugated Hyperbilirubinaemia
3. Importance of urine examination in jaundice patients.
4. Bilirubin metabolism.

Overview

- Introduction
- Definition
- Normal bilirubin metabolism → Synthesis, Hepatic Phase, Intestinal Phase
- Types of bilirubin
- Classification of jaundice →

    1. Prehepatic jaundice → Haemolytic J.
    2. Hepatic jaundice→ Hepatocellular J.
    3. Post hepatic (extrahepatic) jaundice → Obstructive J.

        - Unconjugated Hyperbilirubinaemia
        - Conjugated Hyperbilirubinaemia

- Intrahepatic versus Extrahepatic Cholestasis

Introduction

- Jaundice is not a disease but rather a sign that can occur in many different diseases.

Elevated levels of bilirubin in blood → hyperbilirubinaemia
↓
Bilirubin pigment has high affinity for elastic tissue (skin or sclerae)
↓
Yellow pigmentation of skin or sclerae
↓
Jaundice or Icterus

Definition

- Normal serum bilirubin concentration → 0.3-1.3 mg/dl (80% → unconjugated)
- Latent jaundice → Total serum bilirubin → normal to 2 mg/dl
- Clinically evident Jaundice → Total serum bilirubin → > 2 mg/dl.

Normal Bilirubin Metabolism

1. Synthesis
2. Hepatic Phase
3. Intestinal Phase





1) Synthesis



2) Hepatic Phase

Hepatic uptake →

- Albumin-bound unconjugated bilirubin upon entry into hepatocyte → dissociated into bilirubin and albumin.
- Bilirubin gets bound to cytoplasmic protein glutathione-S-transferase (GST)

Conjugation →

- Occurs in ER
- Involves conversion of Unconjugated bilirubin to Conjugated bilirubin (bilirubin mono- and diglucuronide) by the action of microsomal enzyme, bilirubin- UDP-glucuronosyl transferase

Secretion in bile →

- Conjugated (water-soluble) bilirubin → transported into bile canaliculi and then excreted into the bile

3) Intestinal Phase





Types of Bilirubin

1. Unconjugated bilirubin → Blood, liver
2. Conjugated bilirubin → Liver, Intestine, Blood

	Unconjugated bilirubin	Conjugated bilirubin
Normal levels	**0.3 – 1.6 mg/dl**	**0 to 0.3 mg/dl**
Lipid insolubility	Lipid soluble (lipophilic)	Lipid insoluble
Water insolubility	Water insoluble	Water soluble
Excretion in urine	Tightly bound to albumin →cannot be excreted in urine	Loosely bound to albumin and thus is easily filtered by glomeruli and excreted in urine.

Excretion in stools	Never comes to intestine → cannot be excreted in stools	Comes to intestine → excreted in stools
Brain affinity	Unbound fraction, which is increased in hemolytic reactions can cross blood brain barrier because of lipophilic nature→ cause brain damage of newborn→ Kernicterus	

Classification of Jaundice

2 types of classifications

1. Prehepatic jaundice → Haemolytic J.
2. Hepatic jaundice → Hepatocellular J.
3. Post hepatic (extrahepatic) jaundice → Obstructive J.

1. Unconjugated Hyperbilirubinaemia
2. Conjugated Hyperbilirubinaemia

1) Prehepatic jaundice

Hemolytic anemias
↓
Increased production of bilirubin
↓
Liver is normal → able to conjugate bilirubin at a normal rate
↓
Due to overproduction of bilirubin
↓
Conjugation capacity of liver is overwhelmed
↓
Unconjugate bilirubin is increased
↓
Unconjugated hyperbilirubinemia

Causes

Hemolytic anemias →
- Hereditary spherocytosis
- Thalassemia
- SCA
- Malaria
- Drugs (methyldopa, quinine)

2) Hepatic jaundice

Cause lies within hepatocytes
↓
a) Defect in uptake of bilirubin
b) Defect in conjugation
c) Defect in excretion of conjugated bilirubin into bile canaliculi (Intrahepatic cholestasis)

↓
Both Conjugated and Unconjugated hyperbilirubinemia

Causes

- Hepatitis (infection, alcohol, drugs, toxin or autoimmune)
- Cirrhosis
- Congenital hyperbilirubenemias

3) Post hepatic (extrahepatic) jaundice

Obstruction of excretion into intestine (Extrahepatic cholestasis)
↓
Conjugated hyperbilirubinemia

Causes

- Stone in CBD
- Carcinoma pancreas
- CBD carcinoma

Causes of Jaundice

Pre Hepatic Cause	Hepatic Cause	Post Hepatic Cause
↑ Bilirubin production leading to unconjugated hyperbilirubinemia e.g. hemolytic anemia.	Due to defect in hepatocyte leading to defective conjugation or decreased excretion of conjugated bilirubin.	Due to impaired excretion of conjugated bilirubin as a result of obstruction e.g. stone, cancer of pancreas and bile duct etc.

Unconjugated Hyperbilirubinaemia

1. Increased bilirubin production (Haemolytic, acholuric or prehepatic jaundice)

    i. Intra- and extravascular haemolysis
    ii. Ineffective erythropoiesis

2. Decreased hepatic uptake

    i. Drugs
    ii. Prolonged starvation
    iii. Sepsis

3. Decreased bilirubin conjugation

    i. Hereditary disorders (e.g. Gilbert's syndrome, Crigler-Najjar syndrome)
    ii. Acquired defects (e.g. drugs, hepatitis, cirrhosis)
    iii. Neonatal jaundice

Conjugated Hyperbilirubinaemia (Cholestasis)

1. Intrahepatic cholestasis (Impaired hepatic excretion)

    i. Hereditary disorders or 'pure cholestasis' (e.g. Dubin Johnson syndrome, Rotor's syndrome, fibrocystic disease of pancreas, benign familial recurrent cholestasis, intrahepatic atresia, cholestatic jaundice of pregnancy)
    ii. Acquired disorders or 'hepatocellular cholestasis' (e.g. viral hepatitis, drugs, alcohol-induced injury, sepsis, cirrhosis)

2. Extrahepatic cholestasis (Extrahepatic biliary obstruction)
   Mechanical obstruction (e.g. gallstones, inflammatory strictures, carcinoma head of pancreas, tumours of bile ducts, sclerosing cholangitis, congenital atresia of extrahepatic ducts)

Intrahepatic versus Extrahepatic Cholestasis



Intrahepatic Cholestasis

Liver biopsy

- Biliary canaliculi of hepatocytes are dilated
- Contain characteristic elongated green-brown bile plugs
- Intrahepatic Cholestasis



Extrahepatic Cholestasis

Liver biopsy

Progressive retrograde extension of bile stasis into intrahepatic duct system
Results in dilatation of bile ducts
Rupture of canaliculi with extravasation of bile producing bile lakes
Bile is Toxic
Focal necrosis of hepatocytes surrounding bile lakes



Intrahepatic Cholestasis Extrahepatic Cholestasis

Types of jaundice	Pre hepatic	Hepatic	Post hepatic
Type of bilirubin elevated	unconjugated bilirubin	Both conjugated A unconjugated bilirubin	conjugated bilirubin
Serum bilirubin- Van den Bergh test	Indirect positive	Biphasic	Direct positive
Urine ✓ Conj. Bilirubin ✓ Urobilinogen ✓ Bile salt	 Absent +++ Absent	 ++ + early, obst. -dec. +	 +++ Absent ++
Urine color	Normal-Acholuric	Dark - Choluric	Dark - Choluric
Stool color	Dark brown colour	N /decreased	Clay colored stools
AST A ALT	Normal	Very high	inc.
ALP Levels	Normal	2-3 times increased	10-12 times inc. 21

(17.02) NEONATAL JAUNDICE / CONGENITAL HYPERBILIRUBINEMIA

University Exams

Short Questions

1. Define Congenital hyperbilirubinemia. Write its Classification
1. Define Neonatal Jaundice. Write Pathogenesis, clinical course and hepatic morphology in Dubin-Johson syndrome.
2. Pathogenesis, clinical course and hepatic morphology in Criggler-Najjar Syndrome I and II
3. Pathogenesis and clinical course in Gilbert syndrome

Overview

- Definition
- Classification

1. Gilbert syndrome
2. Criggler-Najjar Syndrome →Type I
3. Criggler-Najjar Syndrome →Type II
4. Duhin Jhonson syndrome
5. Rotor Syndrome

} Inheritance, Hyperbilirubinemia, Defect, Clinical course, Effect of Phenobarbitone Hepatic morphology, Prognosis

Definition →

Jaundice appears in neonates when total serum bilirubin → **> 3 mg/dl**

Classification

Unconjugated Hyperbilirubinaemia

1. Physiologic and prematurity jaundice
2. Haemolytic disease of the newborn and kernicterus
3. Congenital haemolytic disorders
4. Perinatal complications (e.g. haemorrhage, sepsis)
5. Gilbert's syndrome
6. Crigler-Najjar syndrome (type I and II)

Conjugated Hyperbilirubinaemia

1. Dubin-Johnson syndrome
2. Rotor's syndrome
3. Infections (e.g. hepatitis B, hepatitis C or non-A non-B hepatitis, rubella, coxsackievirus, cytomegalovirus, echovirus, herpes simplex, syphilis, toxoplasma, gram-negative sepsis)
4. Metabolic (e.g. galactosaemia, alpha-1-antitrypsin deficiency, cystic fibrosis, Niemann-Pick disease)
5. Idiopathic (neonatal hepatitis, congenital hepatic fibrosis)
6. Biliary atresia (intrahepatic and extrahepatic)
7. Reye's syndrome

Feature	Gilbert's Syndrome	Type 1 Crigler-najjar Syndrome	Type 2 Crigler-najjar Syndrome	Dubin-johnson Syndrome	Rotor Syndrome
Inheritance	Autosomal dominant	Autosomal recessive	Autosomal dominant	Autosomal recessive	Autosomal recessive
Hyperbilirubinaemia	Unconjugated	Unconjugated	Unconjugated	Conjugated	Conjugated
Defect	↓ UDP-glucuronosyl transferase activity	Absence of UDP-glucuronosyl transferase	↓ UDP-glucuronosyl transferase	Impaired biliary excretion of bilirubin glucuronides due to mutation in canalicular multidrug resistance protein 2 (MRP2) (Prolonged BSP excretion test)	Decreased hepatic uptake and storage? Decreased biliary excretion?
Clinical course→ jaundice	Mild (< 5 mg/dl)	Marked (>20 mg/dl) Kernictrus	Mild to moderate (<20 mg/dl)	Mild (<5 mg/dl)	Mild chronic jaundice
Effect of Phenobarbitone	Induces enzyme →↓ Unconjugated bilirubin	No effect	Induces enzyme →↓ Unconjugated bilirubin		
Hepatic morphology	Normal (except slightly increased lipofuscin)	Normal (except mild canalicular stasis)	Normal	Epinephrine metabolites not excreted normally → black pigment	Normal
Prognosis	Excellent	Poor (due to kernicterus)	Good	Excellent	Good

(17.03) VIRAL HEPATITIS

University Exams

Long Questions

1. Classify viral hepatitis. Describe the structure, modes of transmission, course of disease and serological markers for hepatitis B virus.
2. Briefly describe etiologic agents of chronic viral hepatitis. Discuss in details morphology of chronic hepatitis.
3. Describe pathogenesis, lesions and laboratory investigations in viral hepatitis.
4. Discuss Morphology, modes of transmission, Clinical features and Laboratory Diagnosis in Viral Hepatitis A

Short Questions

1. Serological markers of hepatitis B viral hepatitis and their clinical significance.
2. Outcomes of hepatitis B virus infection.
3. Morphology of liver in hepatitis A.
4. Microscopy in Chronic hepatitis.
5. Enumerate microscopic features of acute hepatitis.
6. Describe the sequence and importance of serologic markers for hepatitis B viral hepatitis in acute infection.
7. Non-A non -B hepatitis.

Very Short Questions

1. Name different causative viruses of hepatitis.
2. Laboratory diagnosis of acute viral hepatitis.
3. List serological markers for hepatitis B viral hepatitis.
4. Mention antibodies of hepatitis B infection.
5. Co-infection and Super-infection in hepatitis D viral hepatitis.

Overview

- Liver → Anatomy
- Liver → Dual blood supply
- Liver → Classical lobule

1. HAV
2. HBV
3. HCV
4. HDV
5. HEV

} Causative agent, Morphology, MOT, IP, Clinical features, Laboratory Diagnosis

- Clinicopathologic Spectrum →

1. Acute Hepatitis
2. Chronic Hepatitis

Liver → Anatomy



Liver → Dual blood supply



Liver → Classical lobule

1. A central hepatic vein (branch of hepatic vein)
2. 4 to 5 portal tracts or triads at the periphery → containing branches of bile duct, portal vein and hepatic artery.
3. Cords of hepatocytes (which are one cell thick) and blood containing sinusoids radiate from the central vein to the peripheral portal triads.





Properties	HAV	HBV	HCV	HDV	HEV
Common name	Infectious hepatitis	Serum hepatitis	Non A non B or post-transfusion hepatitis	Delta agent	Non A non B enteric transmitted hepatitis
Genome	ssRNA	dsDNA	ssRNA	ssRNA	ssRNA
Family	Enterovirus-72 (Picornaviridae)	Hepadnaviridae	Flaviviridae	Deltavirus	Hepeviridae
Envelope	No	Yes (HBsAg)	Yes	Yes (HBsAg)	No
Route	Feco-oral	Blood (MC) Sexual, Vertical	Blood (MC) Sexual (+/-) Vertical (+/-)	Blood (MC) Sexual (++) Vertical (+)	Feco-oral
I.P (days)	15–45, (Average 30)	30–180, (Average 60–90)	15–160, (Average 50)	30–180, (Average 60–90)	14–60, (Average 40)
Acute Hepatitis	Yes	Yes	Yes	Yes	Yes
Fulminant disease	Rare (0.1%)	Rare (0.1–1%)	Rare (0.1%)	Frequent (5–20%)	Usually rare (1-2%) Pregnancy-20-40%
Chronicity	None	Occasional (1–10%)	Common (85%)	Common	None
Carrier	None	Yes (0.1–30%)	Yes (1.5–3.2%)	Variable	None
Oncogenic	No	Yes (neonate)	Yes	+/-	No
Cirrhosis	No	Yes	Yes	Yes	No

Prognosis	Excellent	Worse with age	Moderate	Acute-good Chronic-poor	Good
Prophylaxis	Immunoglobulin, Inactivated vaccine	HBIG, Recombinant vaccine	None	HBV vaccine (no vaccine for HBV carriers)	Vaccine (HEV239) (only in China)
Therapy	None	Pegylated interferon Lamivudine	Pegylated interferon plus ribavirin	Interferon ±	None

1) Hepatitis A



Causative agent →

Belongs to family Picornaviridae.

Morphology →

- Linear ssRNA
- Spherical particle with icosahedral symmetry

MOT →

Fecal-oral route



IP →

15–45 days (mean 30 days)

Clinical features

- Onset → abrupt (subacute)
- 2 Phases →
    1. Pre-icteric phase → nausea and vomiting
    2. Icteric phase → jaundice
- Most common cause of acute viral hepatitis in children.
    - No chronic state
    - No carriers state
- Complete recovery → 98%
- Complications → Fulminant hepatitis



Laboratory Diagnosis

1. Anti-HAV antibody detection:
    - Anti-HAV IgM positive → indicates acute infection
    - Anti-HAV IgG antibody in absence of IgM → indicates past infection or recovery
2. Detection of HAV particles: Appears in stool from –2 to +2 weeks of jaundice.



2) Hepatitis B

Causative agent →

Belongs to family Hepadnaviridae

Morphology →

- Only DNA virus among hepatitis viruses.
- 3 forms →

    1. Spherical forms
    2. Tubular or filamentous forms
    3. Complete form or Dane particles



MOT →

1. Parenteral transmission
2. Sexual transmission
3. Perinatal transmission

IP →

30–180 days



Clinical features

Patients may present with

a) Subclinical infection
b) Acute infection
c) Chronic hepatitis



Laboratory Diagnosis

1. Antigen markers
2. Antibody markers

1) Antigen markers:

- **HBsAg:** First marker to appear; elevated in acute, chronic or in carriers
- **HBeAg:** Indicates active viral replication and high infectivity
- **HBcAg:** Not detectable in serum; can be detected in hepatocytes by immunofluorescence test

2) Antibody markers:

- Anti-HBc Ab:
    - IgM → indicates acute hepatitis
    - IgG → appears in chronic hepatitis or carriers or after recovery

- **Anti-HBs Ab:** Marker of recovery or vaccination
- **Anti-HBe Ab:** Indicates low viral replication and low infectivity







HBsAg ↓	• Present in acute disease • Continued presence indicates chronic disease or carrier state
HBeAg **HBVDNA** **DNA polymerase**	• Seen with active viral replication and denotes high infectivity
↓ **IgM anti HBc** ↓	• Antibody detectable shortly before onset of symptoms • Marker of window period • IgM anti HBc is indicator of recent disease whereas IgG anti HBc is seen with chronic infection or prior infection
Anti HBe Ab ↓	• Detected after HBe Ag disappears and denotes low infectivity
IgG anti HBc ↓ **Anti Hbs**	• Appears after disappearance of HBs Ag • Provides protection against Hepatitis B and indicates immunity • Seen in prior infection and in vaccinated persons

	HBs Ag	Anti HBs Ab	HBe Ag	Anti HBe Ab	Anti HBc Ab
Acute HBV (high infectivity)	+	-	+	-	IgM
Acute HBV (low infectivity)	+	-	-	+	IgM
Chronic HBV (high infectivity)	+	-	+	-	IgG
Chronic HBV (low infectivity)	+	-	-	+	IgG
Recovery	-	+	-	+	IgG

3) Hepatitis C

Causative agent →

Belongs to family Flaviviridae

Morphology →

- ssRNA hepatitis viruses
- Spherical and enveloped





MOT →

1. Parenteral transmission
2. Sexual transmission
3. Perinatal transmission

IP →

15–160 days (average 50 days).

Clinical features

1. Asymptomatic infection
2. Acute hepatitis
3. Chronic disease

Extrahepatic manifestations: Due to deposition of circulating immune complexes in extrahepatic sites →

a) Mixed cryoglobulinemia
b) Glomerulonephritis
c) Arthritis and joint pain

Laboratory Diagnosis

1. HCV Antibody Detection Assay (ELISA) →
   - **First generation Ab →** against **C100-3** region proteins and appear 1 to 3 months after infection.
   - **Second generation Ab →** against **C200 and C33c** proteins and appear about one month earlier than first generation.
   - **Third generation Ab → against C22-3 and NS-5** region proteins and are detected even earlier.



2. HCV Core Antigen Assay
3. Molecular Methods

4) Hepatitis D

Causative agent →

- Belongs to family delta virus
- Defective virus for which HBV is helper.
- Replication is defective and can cause infection only when encapsulated by HBs Ag

Morphology →

- ssRNA
- Protein coat is made up of single protein called hepatitis D antigen (HDAg)
- Surrounded by an envelope protein derived from HBsAg from hepatitis B virus → defective virus



MOT →

1. Parenteral transmission
2. Sexual transmission
3. Perinatal transmission

IP →

30–180 (Average60–90)

Clinical features

2 types

1. Co-infection
2. Super-infection



Laboratory Diagnosis

- IgM anti-HBc + IgM anti-HDV→ Indicates co-infection
- IgG anti-HBc + mixture of IgM & IgG anti-HDV → Indicates super-infection

5) Hepatitis E

Causative agent →

Belongs to family Hepeviridae

Morphology →

- ssRNA
- Non-enveloped with icosahedral symmetry



MOT →

Fecal-oral route

IP →

14–60 days (average 40 days).

Clinical features

- Self-limiting acute hepatitis followed by complete recovery
- Fulminant hepatitis → 1–2% of cases
- Fulminant hepatitis in Pregnant women → higher risk (20%)

- No chronic infection
- No carrier state

Laboratory Diagnosis

- HEV RNA (by reverse transcriptase PCR)
- Serum antibody detection by ELISA:

    - IgM anti-HEV → indicates acute infection
    - IgG anti-HEV re→ indicates recovery or past infection.

Clinicopathologic Spectrum

1. Acute hepatitis
2. Chronic hepatitis

1) Acute Hepatitis → Microscopy

Seen in all types (Hepatitis A, B, C, D, E)

1. Hepatocellular injury →

    - Ballooning degeneration → Some hepatocytes appear swollen with granular cytoplasm
    - Councilman body or acidophil body → Some hepatocytes show acidophilic degeneration in which cytoplasm becomes intensely eosinophilic, nucleus becomes small and pyknotic
    - Dropout necrosis →Small clusters of hepatocytes undergo lysis
    - Bridging necrosis → by bands of necrosis linking



2. Inflammatory infiltrate → Infiltration by mononuclear inflammatory cells in portal tracts and lobules.
3. Kupffer cell hyperplasia → Reactive hyperplasia of Kupffer cells

2) Chronic Hepatitis → Microscopy

- Seen in HBV, HCV, HDV
- HAV and HEV do not produce chronic hepatitis.

1. Limiting plate lesion

    - Piecemeal necrosis → Piecemeal necrosis is defined as periportal destruction of hepatocytes at limiting plate (piecemeal = piece by piece).
    - Interface hepatitis

2. Portal tract lesions

- Inflammatory cell infiltration by lymphocytes, plasma cells and macrophages (triaditis)
- Proliferated bile ductules in the expanded portal tracts.

3. Intralobular lesions

- Acidophilic bodies in lobule.
- Kupffer cell hyperplasia.

4. Bridging necrosis

- portal tract-to-central vein
- central vein-to-central vein
- portal tract-to-portal tract



Bridging necrosis from central vein to portal tract

5. Bridging fibrosis

- Dense collagenous septa destroying lobular architecture and forming nodules resulting in cirrhosis





Remember

- Chronic hepatitis due to HBV → HBs Ag in cytoplasm of hepatocytes produce finely granular appearance→ shows ground-glass hepatocytes

Properties	HAV	HBV	HCV	HDV	HEV
Causative agent					
Morphology					
MOT					
IP					
Clinical features					
Laboratory Diagnosis					

(17.04) CIRRHOSIS (DEFINITION, PATHOGENESIS, CLASSIFICATION)

University Exams

Short Questions

1. Morphologic and Etiologic classification of cirrhosis.
2. Define and classify cirrhosis. Describe its Pathogenesis
3. Write causes of micronodular cirrhosis.

Overview

- Definition
- Features
- Pathogenesis
- Classification →

    1. Morphologic Classification
    2. Etiologic Classification
    3. Combined

Definition→

- It represents the **irreversible end-stage** of several diffuse diseases causing hepatocellular injury

Features →

- Involves entire liver
- Normal lobular architecture of hepatic parenchyma is disorganised.
- Formation of nodules separated from one another by irregular bands of fibrosis.

Pathogenesis

Injurious agent
↓
Continued destruction of hepatocytes
↓
Collapse of normal lobular hepatic parenchyma
↓
Fat storing Ito cells transformed into myofibroblasts and fibrocytes
↓
Increased synthesis of type I and III collagen in space of Disse
↓
Fibrosis in lobules → portal-central, portal-portal, or both
↓
Compensatory proliferation of Surviving hepatocytes under influence of growth factors
↓
Restricted within fibrous nodules forming regenerative nodules
↓
Cirrhosis

Classification

1. Morphologic Classification
2. Etiologic Classification
3. Combined

Morphologic	Etiologic
1. Micronodular (nodules less than 3 mm) 2. Macronodular (nodules more than 3 mm) 3. Mixed  	1. Alcoholic cirrhosis (the most common, 60-70%) 2. Post-necrotic cirrhosis (10%) 3. Biliary cirrhosis (5-10%) 4. Pigment cirrhosis in haemochromatosis (5%) 5. Cirrhosis in Wilson's disease 6. Cirrhosis in a-1 –antitrypsin deficiency 7. Cardiac cirrhosis 8. Indian childhood cirrhosis (ICC) 9. Cirrhosis in autoimmune hepatitis 10. Cirrhosis in non-alcoholic steatohepatitis 11. Miscellaneous forms of cirrhosis (metabolic, infectious, GI, infiltrative) diseases 12. Cryptogenic cirrhosis

Causes of Micronodular cirrhosis
• Indian childhood cirrhosis
• Alcoholic cirrhosis
• Hemochromatosis
• Cirrhosis associated with biliary obustruction
• Cirrhosis associated with jejunoileal bypass
• Venous outflow obstruction → veno - occlusive disease, Budd-chiary syndrome.
Causes of Macronodular cirrhosis
• Post necrotic cirrhosis (after viral hepatitis)
• Wilson disease

(17.05) ALCOHOLIC LIVER DISEASE (ALCOHOLIC CIRRHOSIS)

University Exams

Long Questions

1. Define cirrhosis of liver. Describe etiopathogenesis, morphology, histopathology and complications of alcoholic liver disease.
2. A Male aged 52 years developed gradual weakness, anorexia, weight loss. He has ascites, splenomegaly, jaundice, spider angiomas on skin and gynecomastia. Over a time, he developed behavioral abnormalities, stupor and slipped into coma. There is history of chronic alcoholism:

    a) What is the probable diagnosis?
    b) Discuss etiopathogenesis, pathology and complications.

3. Give chronological classification of alcoholic liver disease. Describe gross & microscopic findings of alcoholic cirrhosis.
4. Discuss etiopathogenesis of Fatty change. Describe morphology of fatty liver.

Short Questions

1. Etiopathogenesis of Alcoholic liver disease
2. Mention lesions of alcoholic liver disease.
3. Morphology of Alcoholic steatosis (fatty liver)
4. Morphology of Alcoholic hepatitis (Steatohepatitis)
5. Gross and microscopy of Alcoholic cirrhosis (steatofibrosis)

Very Short Questions

1. Microscopic appearance and conditions associated with Mallory body.

Overview

- Introduction
- Risk factors
- Ethanol Metabolism
- Pathogenesis
- 3 stages → Morphology

    1. Alcoholic steatosis (fatty liver)
    2. Alcoholic hepatitis (Steatohepatitis)
    3. Alcoholic cirrhosis (steatofibrosis)

Introduction →

Spectrum of liver injury associated with acute and chronic alcoholism.

Risk Factors

1. Gender→

    - Females are more susceptible to ALD than males.
    - Estrogen increases gut permeability to endotoxins (gut-derived) and increased production of pro-inflammatory cytokines and chemokines→ injury to liver cells.

2. Genetic factors→
    - Polymorphisms in aldehydedehydrogenase (ALDH)
3. Associated conditions →
    - Iron overload and chronic infections with HCV and HBV → increase severity of ALD
4. Amount and duration of alcohol intake (drinking patterns) →
    - Type of beverage does not affect risk
    - Damage is more in continuous rather than binge drinkers

Ethanol Metabolism



Pathogenesis

Acetaldehyde →

- Toxic
- Cause membrane damage and cell necrosis
- Produces hepatotoxicity by formation of 2 adducts →
    1. Production of protein-aldehyde adducts → extremely toxic
    2. Formation of malon-di-aldehyde-acetaldehyde (MAA) adducts → produce autoantibodies and initiate autoimmune response.



3 stages

1. Alcoholic steatosis (fatty liver)
2. Alcoholic hepatitis (Steatohepatitis)
3. Alcoholic cirrhosis (steatofibrosis)



1) Alcoholic Steatosis (Fatty Liver)

1. Microvesicular droplets → Initially → Droplets of fat in hepatocyte cytoplasm
2. Macrovesicular large droplets→ Later on → fat displacing nucleus to periphery
3. Fat cysts → due to coalescence and rupture of fat-containing hepatocytes.



2) Alcoholic Steatohepatitis

1. Ballooning degeneration→ Hepatocellular necrosis → hepatocytes appear swollen with granular cytoplasm
2. Mallory-Denk bodies or alcoholic hyalin body → eosinophilic, intracytoplasmic inclusions seen in perinuclear location within ballooned hepatocytes
3. Inflammatory response → polymorphs and some scattered mononuclear cells.



4. Fibrosis → Collagen accumulates in the space of Disse of centrilobular region, spreading outward→ encircling individual or small clusters of hepatocytes → chicken wire fence pattern or creeping collagenosis



3) Alcoholic Cirrhosis

Gross →

- Micronodular cirrhosis → nodules less than 3 mm diameter
- Surface of liver in alcoholic cirrhosis is studded with diffuse nodules → hobnail liver → because of resemblance of surface with sole of an old-fashioned shoe having short nails with heavy heads



Microscopy

1. Nodular pattern: Normal lobular architecture is effced and replaced with nodule formation.
2. Fibrous septa
3. Regenerative nodules→ Hepatocytes in islands of surviving parenchyma undergo slow proliferation forming regenerative nodules having disorganised masses of hepatocytes (fatty change)
4. Bile duct proliferation



(17.06) NON-ALCOHOLIC FATTY LIVER DISEASE (NAFLD)

University Exams

Short Questions

1. Define Non-alcoholic fatty liver disease (NAFLD). Discuss its Pathogenesis and Microscopy.
2. Clinical presentation and Diagnosis of NAFLD.

Overview

- Introduction
- Definition
- Pathogenesis
- Microscopy
- Clinical presentation
- Diagnosis

Introduction →

- Also known as →
  - Non-alcoholic steatohepatitis (NASH)
  - Metabolic Liver Disease
- Most common cause of →
  - chronic liver disease in united states
  - cryptogenic cirrhosis (cirrhosis of unknown origin)

Definition →

- NAFLD represents a spectrum of disorders in which presence of hepatic steatosis (fatty liver) in individuals who do not consume alcohol or do so in very small quantities (less than 20 g of ethanol/week)
- NAFLD resembles alcohol induced liver disease but occurs in non-alcoholics

Pathogenesis

Metabolic syndrome

One of	Diabetes mellitus or impaired glucose tolerance or impaired fasting glucose or insulin resistance
and two of:	• Blood pressure: ≥ 140/90 mm Hg • Dyslipidaemia: triglycerides (TG): ≥ 1.695 mmol/L and high-density lipoprotein cholesterol (HDL-C) < 0.9 mmol/L (male), < 1 mmol/L (female) • Central obesity: waist-hip ratio > 0.90 (male); > 0.85 (female), or body mass index > 30 kg/m^2 • Microalbuminuria: urinary albumin excretion rate of ≥ 20 pig/min or albumin-to-creatinine ratio ≥ 30 mg/gm

Microscopy

- Completely overlaps in its histologic features with alcoholic hepatitis

In NASH, compared with alcoholic hepatitis

1. **Mononuclear cells** → more prominent than neutrophis
2. **Mallory-Denk bodies** → less prominent.





Clinical presentation



Diagnosis →

- Liver biopsy

(17.07) WILSON DISEASE

University Exams

Short Questions

1. Pathogenesis and Clinical features of Wilson's disease.
2. Diagnosis of Wilson's disease.

Very Short Questions

1. Kayser-Fleischer rings.
2. Triad of Wilson's disease.

Overview

- Introduction
- Age
- Pathogenesis
- Clinical features
- Triad
- Diagnosis

Introduction →

- Autosomal recessive disorder
- Caused by **mutation of the ATP7B gene** → impaired copper excretion into bile and a failure to incorporate copper into ceruloplasmin

Age→

In children and young adults (5-30 years).

Pathogenesis



Normally

Ingested copper (2 to 5 mg/day)
↓
Absorbed in duodenum and proximal small intestine
↓
Complexed with albumin
↓
Transported to portal circulation
Free copper dissociates and is taken up by hepatocytes
↓
In liver copper binds to an α2-globulin (apoceruloplasmin) to form ceruloplasmin (copper-transporting ATPase is required)
↓
Secreted into blood
↓
Excess copper is transported into bile

Wilson disease

Wilson disease
↓
Mutation in ATP7B gene present on chromosome 13 (Encodes 'copper-transporting ATPase responsible for incorporation into ceruloplasmin in liver)
↓
Decrease in circulating ceruloplasmin
↓
Copper accumulation in liver → toxic liver injury
↓
Spillage of non–ceruloplasmin-bound copper from liver into circulation
↓
Deposited in extrahepatic tissues such as brain, eyes

Clinical features

1. Hepatic →

- Most common organ involved is liver
- Can present with Fatty change, Acute and chronic hepatitis,Cirrhosis
- Mallory-Denk bodies seen

2. CNS →

- Basal ganglia → involved primarily.
- Dystonia, dysarthria, tremor and chorea.
- Emotional liability
- Depression
- Hyperactivitiy.
- Motor deficit and sensory involvement are characteristically absent.

3. Ocular involvement →

- Kayser-Fleischer ring (brown/green deposits in Descemets membrane of cornea
- Sunflower cataract



4. Others →



- Coombs negative hemolytic anemia
- Fanconi syndrome (amino aciduria)
- Cardiomyopathy.

Triad

1. Cirrhosis of the liver
2. Bilateral degeneration of basal ganglia of the brain
3. Greenish-brown pigmented rings in the periphery of the cornea (Kayser-Fleischer rings).

Cirrhosis



Basal Ganglia



Kayser-Fleischer rings

Diagnosis

1. Liver biopsy with quantitative copper assay → Gold standard → concentration of copper in a liver biopsy sample > 200 mg/g dry weight
2. Serum ceruloplasmin level → low
3. Urine copper excretion -> increased
4. KF rings
5. DNA Haplotype analysis.

(17.08) HEMOCHROMATOSIS

University Exams

Short Questions

1. Types and Pathogenesis of Hemochromatosis.
2. Clinical features and Diagnosis of Hemochromatosis.
3. Triad of Hemochromatosis.

Overview

- Introduction
- Age
- Types
- Pathogenesis
- Clinical features
- Triad
- Diagnosis
- Treatment

Introduction →

Iron-storage disorder in which there is excessive accumulation of iron in parenchymal cells with eventual tissue damage

Age →

- In men → 4th to 5th decades of life
- In women → Later → since menstrual bleeding counterbalances accumulation until menopause.

Types

1. Idiopathic or hereditary haemochromatosis
2. Secondary or acquired haemochromatosis

Pathogenesis

1. Idiopathic or hereditary haemochromatosis
2. Secondary or acquired haemochromatosis

1) Idiopathic or hereditary haemochromatosis

- Primary mechanism of disease appears to be the genetic basis→ defect at intestinal mucosal level → excessive iron absorption

Regulation of iron absorption



Increases in intrahepatic iron levels
↓
Hepcidin is synthesized and released from liver
↓
Hepcidin binds to ferroportin
↓
Degrade it
↓
Inhibits iron transfer from enterocyte to plasma
↓
Iron becomes trapped within duodenal cells
↓
Iron is lost as these cells are sloughed

When body is replete with iron
↓
High hepcidin levels
↓
Inhibit iron absorption into blood

When body iron stores are low
↓
Hepcidin synthesis falls
↓
Facilitates iron absorption into blood

Hepcidin synthesis is controlled by

1. Hemojuvelin **(HJV)**
2. Transferrin receptor 2 **(TFR2)**

3. **HFE**

Hereditary haemochromatosis
↓
HFE or HJV or TFR2 or HAMP gene mutations
↓
Reduce hepcidin synthesis
↓
Diminishing circulating hepcidin
↓
Decreased hepcidin-ferroportin interaction
↓
Allows for increased ferroportin activity
↓
Increased iron efflux from enterocytes
↓
Systemic iron overload



Adult form of hemochromatosis is almost always caused by **mutations of HFE**
- Most common HFE mutation → cysteine-to-tyrosine substitution at amino acid 282 **(C282Y)**
- Other common mutation → **H63D** (histidine at position 63 to aspartate)

2) Secondary or acquired haemochromatosis

Due to acquired causes →
- Ineffective erythropoiesis
- Defective haemoglobin synthesis
- Multiple blood transfusions

Enhanced absorption of iron due to alcohol consumption → Bantu siderosis
- Observed in South African Bantu tribals
- Consume large quantities of home-brew prepared in iron vessels.

Clinical Features

1. Liver (95%)→

    - First organ to be involved
    - Iron becomes evident first as golden-yellow hemosiderin granules in cytoplasm of periportal hepatocytes that stain with Prussian blue
    - Hepatomegaly, micronodular cirrhosis (characteristic) and hepatocellular carcinoma (in 30%).



2. Skin (90%) → Skin hyperpigmentation (bronzing of skin) due to deposition of hemosiderin



3. Pancreas → Pancreatic involvement causes diabetes mellitus (bronze diabetes)
4. Joints → Synovitis, arthropathy, Polyarthritis, Pseudogout
5. Cardiac → CHF, Restrictive cardiomyopathy Arrhythmia
6. Pituitary → decrease production of gonadotropins → Hypogonadism

Triad

1. Micronodular cirrhosis
2. Diabetes mellitus (bronze diabetes)
3. Skin hyperpigmentation

Most common causes of death → CHF and hepatocellular carcinoma.

Diagnosis

1. Liver biopsy →

    - Biochemical determination of hepatic tissue iron concentration → gold standard.
    - In normal individuals → iron content of liver tissue <1000 µg per gram dry weight of liver.
    - Hereditary hemochromatosis → iron content of liver tissue > 10,000 µg iron per gram dry weight

2. Genetic testing

Treatment

1. **Phlebotomy** → Removal of excessive iron
2. **Chelating agents like desferrioxamine**

(17.09) PORTAL HYPERTENSION

University Exams

Short Questions

1. Clinical features and Complications of portal hypertension.
2. Define portal hypertension. Write its causes
3. Discuss Sequelae of Portal Hypertension
4. Pathogenesis of Ascites in Portal Hypertension
5. Pathogenesis of Varices in Portal Hypertension

Overview

- Introduction
- Definition
- Classification
- Causes
- Sequelae of Portal Hypertension →

    1. Ascites
    2. Varices (collateral channels or portosystemic shunts)
    3. Splenomegaly
    4. Hepatic encephalopathy

Introduction

Cirrhosis of the liver
↓
Obstruction to the portal blood flow anywhere along its course
↓
Increase in pressure in portal system
↓
Portal Hypertension



Definition →

- Normal portal venous pressure → 10-15 mm saline
- Portal hypertension occurs when portal pressure > 30 mm saline

Classification

Based on site of obstruction to portal venous blood flow→

1. Intrahepatic →

    - Most common cause Cirrhosis

2. Posthepatic →

   - Obstruction to blood flow through hepatic vein into inferior vena cava.
   - Congestive heart failure and constrictive pericarditis cause portal hypertension by transmitting the elevated pressure through hepatic vessels into portal vein.

3. Prehepatic →

   - Blockage of portal flow before portal blood reaches the hepatic sinusoids

Causes

INTRAHEPATIC

1. Cirrhosis
2. Metastatic tumours
3. Budd-Chiari syndrome
4. Hepatic veno-occlusive disease
5. Diff use granulomatous diseases
6. Extensive fatty change

POSTHEPATIC

1. Congestive heart failure
2. Constrictive pericarditis
3. Hepatic veno-occlusive disease
4. Budd-Chiari syndrome

PREHEPATIC

1. Portal vein thrombosis
2. Neoplastic obstruction of portal vein
3. Myelofibrosis
4. Congenital absence of portal vein



Sequelae of Portal Hypertension

1. Ascites
2. Varices (collateral channels or portosystemic shunts)
3. Splenomegaly
4. Hepatic encephalopathy

1) Ascites

1. Accumulation of excessive volume of fluid within peritoneal cavity.
2. Ascites becomes clinically detectable when more than 500 ml of fluid has accumulated in peritoneal cavity.
3. Development of ascites is associated with haemodilution, oedema and decreased urinary output.
4. Ascitic fluid is transudate with specific gravity of 1.010, protein content below 3 gm/dl



Pathogenesis

1. Decreased plasma colloid oncotic pressure → hypoalbuminaemia from impaired hepatic synthesis
2. Impaired renal excretion→ Reduced renal blood flow and excessive release of ADH results in renal retention of Na and water
3. Hyperaldosteronism → In cirrhosis, there is increased aldosterone secretion by adrenal gland due to reduced renal blood flow, and impaired hepatic metabolism and excretion of aldosterone.





2) Varices (Collateral channels or Porto-systemic shunts)

Obstruction in portal circulation
↓
Rise in portal venous pressure
↓
Blood bypass liver and return to heart by development of porto-systemic collateral channels or shunts or varices



Sites

Varices develop at sites where systemic and portal circulations have common capillary beds.

1. Oesophageal varices: massive haematemesis
2. Haemorrhoids: haemorrhoids
3. Caput medusae: Anastomoses between the portal and systemic veins may develop between the hilum of liver and umbilicus along the paraumbilical plexus of veins resulting in abdominal wall collaterals.



3) Splenomegaly

- Enlargement of the spleen in prolonged portal hypertension → congestive splenomegaly

4) Hepatic encephalopathy

- Characterised by disturbed consciousness, neurologic signs and flapping tremors.


Hepatic encephalopathy
Spider angioma (skin)
Esophageal varices
Splenomegaly
Caput medusae
Ascites
Hemorrhoides
Testicular atrophy
Cirrhosis

(17.10) CVC LIVER / NUTMEG LIVER

University Exams

Short Questions

1. Gross and Microscopy of Nutmeg liver
2. Pathogenesis and morphology of CVC liver

Overview

- Introduction
- Gross
- Microscopy



Introduction

- **In left-sided heart failure** → pulmonary congestion (**CVC lungs**)
- **In right-sided heart failure** → systemic venous congestion (**CVC of systemic organs →Liver, spleen**)



Gross→

- Cut surface shows nutmeg appearance → due to red (ongested centre of lobules) and yellow (atty peripheral zone) mottled appearance



Microscopy→

- Red → Centrilobular haemorrhagic necrosis
- Yellow → peripheral zone of the lobule shows fatty change
- So nutmeg appearance



(17.11) LIVER ABCESS (PYOGENIC, AMOEBIC, HYDATID CYST)

University Exams

Short Questions

1. Write clinical features, gross and microscopic features of Hydatid cyst (hydatid disease).
2. Write clinical features, gross and microscopic features of amoebic liver abscess.
3. Write Pathogenesis, clinical features, gross and microscopic features of Pyogenic Liver Abscess

Overview

- Pyogenic Liver Abscess
- Amoebic Liver Abscess
- Hydatid Cyst

} Introduction, Causative agent, MOT
Pathogenesis, Gross, Microscopy
Clinical features

Pyogenic Liver Abscess

Introduction →

- Most liver abscesses are of bacterial (pyogenic) origin
- Incidence is higher in old age and in immunosuppressed patients such as in AIDS, transplant recipients and those on intensive chemotherapy.

Causative agent →

Commonest infecting organisms →

- Gram-negative bacteria chiefly E. coli
- Pseudomonas
- Klebsiella
- Enterobacter
- Actinomyces
- Bacteroides

MOT

1. Following Ascending cholangitis through ascending infection in biliary tract due to obstruction e.g. gallstones, cancer, sclerosing cholangitis and biliary strictures.

2. Portal pyaemia by means of spread of pelvic or gastrointestinal infection resulting in portal pylephlebitis or septic emboli e.g. appendicitis, empyema of gallbladder, diverticulitis, regional enteritis, pancreatitis

3. Septicaemia through spread by hepatic artery.

4. Direct infection e.g. from adjacent perinephric abscess, secondary infection in amoebic liver abscess, metastasis and formation of haematoma following trauma.

5. Iatrogenic eg. liver biopsy, percutaneous biliary drainage and accidental surgical trauma.





Gross

- Common in right lobe of liver
- Occur as single or multiple yellow abscesses, 1 cm or more in diameter, in an enlarged liver.
- Thick fibrous capsule.



Microscopy

- Multiple small neutrophilic abscesses with areas of extensive necrosis
- Adjacent viable area shows pus and blood clots in the portal vein, inflammation, congestion and proliferating fibroblasts.

Clinical features

- Pain in the right upper quadrant
- Fever
- Tender hepatomegaly
- Jaundice

Amoebic Liver Abscess

Introduction →

- Less common than pyogenic liver abscesses

Causative agent →

Entamoeba histolytica

3 morphological forms →

1. Trophozoite
2. Pre-cystic stage
3. Cyst



MOT →

Ingestion of water and food containing these cysts.

Infective form → Quadrinucleate Cyst

Pathogenesis

Ingestion of water and food containing cysts
↓
Cyst reaches caecum
↓
Excystation
↓
Produces metacystic trophozoites
↓
lodge in submucousa of large intestine
↓
Flask-shaped ulcers
↓
Intestinal amoebiasis (Amoebic dysentery)
↓
Some trophozoites enter into portal vein
↓
Carried away to liver
↓
Extraintestinal amoebiasis (Amoebic hepatitis)







Gross

- Solitary
- Located in the right lobe in posterosuperior portion
- Anchovy sauce Abcess → Centre of abscess contains large necrotic area having reddish-brown, thick pus resembling anchovy or chocolate sauce



Microscopy

- Necrotic area consists of degenerated liver cells, leucocytes, red blood cells, strands of connective tissue and debris.
- Amoebae seen at the margin of abscess.
- PAS-staining is employed to confi rm the trophozoites of E. histolytica.

Clinical features

Intestinal amoebiasis→
- Bloody diarrhoea

Extraintestinal amoebiasis →
- Amoebic liver abscess
- posteriosuperior surface of the right lobe of liver
- Pain in the right upper quadrant
- Tender hepatomegaly

Hydatid Cyst (Echinococcosis)

Introduction→

- Disease is common in sheep-raising countries such as Australia, New Zealand and South America.

Causative agent →

- Echinococcus granulosus (Cestode)
- Also called as →
    - Dog tapeworm
    - Hydatid worm

3 morphological forms →

1. Adult
2. Egg
3. Larva

MOT →

- Ingestion of food and water contaminated with eggs

Infective form → Eggs

Pathogenesis

Host → 2

- Definitive host → Dog, wolf, fox and jackal
- Intermediate host →Sheep, pig, cattle, horse, goat
- Accidental intermediate host (dead end) → Man



Gross

- Hydatid cyst grows slowly and may eventually attain a size over 10 cm in diameter in about 5 years.
- Unilocular



Microscopy



1. Pericyst → outer host inflammatory reaction consisting of fibroblastic proliferation, mononuclear cells, eosinophils and giant cells, eventually developing into dense fibrous capsule which may even calcify.
2. Ectocyst → intermediate layer composed of characteristic acellular, chitinous, laminated hyaline material
3. Endocyst → inner germinal layer bearing daughter cysts (brood-capsules) and scolices projecting into the lumen.

- Hydatid sand → grain-like material composed of numerous scolices present in the hydatid fluid.
- Hydatid fluid → contains antigenic proteins → liberation into circulation → Anaphylaxis.

Clinical features



Unilocular hydatid disease / Cystic echinococcosis (CE)

Symptoms: due to →

- Pressure effect of the enlarging cyst
- Obstruction
- Secondary bacterial infection, causing pyogenic abscess in liver
- Anaphylactic reactions

	Pyogenic Liver Abscess	Amoebic Liver Abscess	Hydatid Cyst
Introduction			
Causative agent			
MOT			
Pathogenesis			
Gross			
Microscopy			
Clinical features			

(17.12) MALIGNANT TUMOURS OF LIVER

University Exams

Long Questions

1. Discuss etiopathogenesis, morphology and serum markers of hepatocellular carcinoma.
2. Give classification of Liver cirrhosis. Discuss gross, microscopic findings, clinical features and spread of Hepatocellular Carcinoma.

Short Questions

1. Morphology of hepatocellular carcinoma.
2. Tumour Markers of hepatocellular carcinoma.
3. Describe etiology, gross and microscopy of Fibrolamellar carcinoma liver.
4. Describe etiopathogenesis and morphological features of primary carcinoma of liver.

Overview

Classification

Hepatocellular Carcinoma

- Introduction
- Age
- Gender
- Etiology
- Molecular Basis of HCC
- Gross
- Microscopy
- Clinical features
- Spread
- Serum Markers
- Prognosis

Fibrolamellar Carcinoma

- Introduction
- Age
- Gender
- Etiology
- Gross
- Microscopy
- Serum Markers
- Prognosis

Malignant tumours of Liver

BENIGN	MALIGNANT
a) Hepatocellular tumours Hepatocellular (liver cell) adenoma	Hepatocellular (liver cell) carcinoma Hepatoblastoma (Embryoma)
b) Biliary tumours Bile duct adenoma (Cholangioma)	Cholangiocarcinoma Combined hepatocellular and cholangiocarcinoma Cystadenocarcinoma
c) Mesodermal tumours Haemangioma	Angiosarcoma Embryonal sarcoma

Remember

- Malignant tumors of liver →

    1. Primary
    2. Metastatic

- Metastasis > Primary

Adults →
- Most common primary benign lesion of liver → **Cavernous hemangioma**
- Most common primary malignant lesion of liver → **hepatocellular carcinoma.**

Children →
- Most common primary hepatic tumor of children → **hepatoblastoma**

Hepatocellular Carcinoma

Introduction →

- Most common primary malignant tumour of the liver → hepatocellular carcinoma (HCC)
- Supervenes on cirrhosis in 70-80% of cases

Age →

5th to 6th decades of life

Gender →

M > F (4:1)

Etiology

1. Relation to HBV infection
2. Relation to HCV infection
3. Relation to cirrhosis
4. Relation to alcohol
5. Mycotoxins
6. Chemical carcinogens

Molecular Basis of HCC

- Activation of oncogene → eg. KRAS, TGF- **α**, WNT signal transduction proto-oncogenes **β** -catenin.
- Inactivation of tumor suppressor gene → e.g. TP53

Gross

3 Patterns →

1. **Expanding type→** single, yellowbrown, large mass in right lobe of liver with central necrosis, haemorrhage and bile staining, encapsulated.
2. **Multifocal type →** multifocal, multiple masses,3-5 cm in diameter, scattered throughout liver
3. **Infiltrating (Spreading) type→** diffusely infiltrating tumour mass



Microscopy

Histologic patterns →

1. Trabecular or sinusoidal pattern → Most common.
2. Trabeculae are made up of 2-8 cell wide layers of tumour cells separated by sinusoids
3. **Pseudoglandular or acinar pattern**→Tumour cells are disposed around central cystic space formed by degeneration
4. **Compact pattern** → Tumour cells form large solid masses
5. **Scirrhous pattern**→ abundant fibrous stroma.



Cytologic features →

Tumour cells have→

1. Nucleus → Vesicular nuclei with prominent nucleoli
2. Cytoplasm → granular and eosinophilic
3. Cellular features → pleomorphism, bizarre giant cell formation, spindle-shaped cells, presence of bile within dilated canaliculi, and intracytoplasmic Mallory's hyalin.

Clinical features

- Abdominal pain
- Weight loss
- Ascites
- Hepatomegaly
- Associated paraneoplastic syndromes →
    - Polycythemia (PTH-related protein)
    - Hypoglycemia (insulin-like factor)
    - Hypercalcemia and erythrocytosis (EPO-erythropoietin)

Spread

Local spread:

- HCC first spread within liver itself → develop satellite nodules.
- Local invasion of the diaphragm is common.

Lymphatic spread:

- To portal lymph nodes, perihilar, peripancreatic, and para-aortic nodes.

Blood spread:

- Strong tendency for invasion of vessels.
- Result in extensive intrahepatic metastases.
- Portal vein and its branches are infiltrated by tumor.
- Rarely, tumor may invade → hepatic veins → inferior vena cava → right side of the heart
- Metastasize to the lungs.



Tumour Markers

1. Alpha-fetoprotein (AFP) →
    - About 50% HCC associated with high serum levels of AFP
    - AFP levels are also raised in other neoplastic and non-neoplastic liver diseases and in some extrahepatic disorders.
2. **α** -L-fucosidase →
    - Raised in HCC and also in cirrhosis.
3. Serum des- **α -** carboxy prothrombin →
    - Raised in a majority of HCC

Prognosis

- Unresected hepatocellular carcinoma has a very poor prognosis.
- Survival rate → 40% and 26% at 5 and 10 year repectively

Cause of Death →

1. Cachexia
2. Gastrointestinal or esophageal variceal bleeding
3. Liver failure with hepatic coma
4. Rupture of the tumor with fatal hemorrhage (rare)

Fibrolamellar Carcinoma

Introduction →

- A Distinctive uncommon variant of HCC
- Constitutes about 5% of HCCs

Age →

Young patients without cirrhosis

Gender →

M = F

Etiology →

- Unknown
- Occurs in the absence of cirrhosis.



Gross

- Single large, hard "scirrhous" well-circumscribed tumor with central stellate fibrous scar.



Microscopy

- Tumor cells arranged in nests or cords, and separated by parallel bands of abundant dense collagen bundles
- Consists of large, polygonal cells
- Abundant deeply eosinophilic (oncocytic) cytoplasm
- Prominent nucleoli.

Tumour marker →

Neurotensin

Prognosis →

Better than conventional HCC.

(17.13) CHOLELITHIASIS (GALL STONES)

University Exams

Long Questions

1. Discuss etiopathogenesis of cholelithiasis. Describe morphology of different kinds of gallstones. List the complications.

Short Questions

1. Risk factors for gallstones.
2. Enlist etiological factors for pigment gallstones.
3. Cholelithiasis → types and complications
4. Types of gallstones.
5. Pathogenesis of cholelithiasis

Overview

- Introduction
- Risk factors
- Pathogenesis →

    1. Cholesterol and mixed gallstones
    2. Pigment gallstone

- Types Of Gallstones
- Clinical features and Complications

Introduction →

Gallstones are formed from constituents of the bile (cholesterol, bile pigments and calcium salts) along with other organic components.

Risk Factors

Gallstones are common in 4F's acronym for—
1. Fat
2. Female
3. Fertile (multipara)
4. Forty

1. Geography
2. Genetic factors
3. Age
4. Sex
5. Drugs
6. Obesity
7. Diet
8. Factors in pigment gallstones

1) Geography

- Prevalent in Western world
- American Indians have highest known prevalence.
- Black Africans and populations in Eastern world are relatively free of cholelithiasis.

2) Genetic factors

Mutation in CYP7A1 gene
↓
Deficiency of enzyme → cholesterol 7-hydroxylase
↓
Hypercholesterolaemia
↓
Gallstones

3) Age

- Incidence increases above age of 40

4) Sex

- F > M
- Multiparous women > Nulliparous women

5) Drugs

Women on oestrogen therapy
or on birth control pills
↓
cholestatic effect of oestrogen
↓
production of more lithogenic bile
↓
Higher incidence of gallstones

6) Obesity

Obesity
↓
Increased cholesterol synthesis
↓
Increased cholesterol excretion
↓
Higher incidence of gallstones in obese patients

7) Diet

- Deficiency of dietary fibre → higher prevalence of gallstones.
- Moderate consumption of alcohol → protect against gallstones.

8) Factors in pigment gallstones

Haemolytic anaemias
↓
Increased content of unconjugated bilirubin in bile
↓
Pigment stones

Pathogenesis

1. Cholesterol and mixed gallstones
2. Pigment gallstone

1) Cholesterol and Mixed Gall stones

Cholesterol is insoluble in water and can be solublised by another lipid.

- Cholesterol + phospholipids (lecithin) = bilayered vesicles
- Cholesterol + phospholipids (lecithin) + Bile acids = mixed miscelles

If there is excess of cholesterol compared to other two constituents
↓
Unstable cholesterol-rich vesicles
↓
Aggregate and form cholesterol crystals.
↓
Lithogenic (stone-forming) bile

2) Pigment Gallstones

1. Chronic haemolysis → increased level of unconjugated bilirubin in bile.
2. Alcoholic cirrhosis
3. Chronic biliary tract infection → e.g. by parasitic infestations of biliary tract such as by Clonorchis sinensis and Ascaris lumbricoides.

Types of Gallstones

TYPE	FREQUENCY	COMPOSITION	GALLBLADDER CHANGES	APPEARANCE
1. Pure gallstones	10%	• Cholesterol	Cholesterolosis	Solitary, oval, large, smooth, yellow white; on C/S radiating glistening crystals
		• Bile pigment	No change	Multiple, small, jet-black, mulberry shaped; on C/S soft black
		• Calcium carbonate	No change	Multiple, small, grey-white, faceted; C/S hard
2. Mixed gallstones	80%	Cholesterol, bile pigment and calcium carbonate in varying combination	Chronic cholecystitis	Multiple, multifaceted, variable size, on C/S laminated alternating dark pigment layer and pale-white layer
3. Combined gallstones	10%	Pure gallstone nucleus with mixed gallstone shell, or mixed gallstone nucleus with pure gallstone shell	Chronic cholecystitis	Solitary, large, smooth; on C/S central nucleus of pure gallstone with mixed shell or vice versa



CHOLESTEROL (YELLOW-WHITE) | **PIGMENT** (MULBERRY-SHAPED) | **CALCIUM CARBONATE** (HARD)

MIXED GALLSTONES **(MULTIFACETED)** | COMBINED GALLSTONES **(SMOOTH-SURFACED)**

Clinical features and Complications

- 50% cases → silent gallstones → Asymptomatic and may be diagnosed by chance during investigations for some other condition
- Symptomatic gallstone disease appears only when complications develop.

1. **Cholecystitis →** typical biliary colic precipitated by fatty meal, nausea, vomiting, fever along with leucocytosis and high serum bilirubin.

2. **Choledocholithiasis →** Gallstones may pass down into the extrahepatic biliary passages and small bowel → pain and obstructive jaundice.

3. **Mucocele and empyema →** distension of gallbladder by clear, watery mucinous secretion resulting from impacted stones in neck of gallbladder.



- When it gets infected empyema is formed.

4. **Biliary fistula** → Formation of fistulae between one part of the biliary system and bowel
5. **Gallstone ileus** → Gallstones in intestine may cause intestinal obstruction
6. **Pancreatitis** → Obstructive cholecystasis may result in acute pancreatitis.
7. **Gallbladder cancer** → Rare

(17.14) CHOLECYSTITIS

University Exams

Short Questions

1. Microscopy of acute and chronic cholecystitis.
2. Etiopathogenesis, Gross, Microscopy and Clinical features of acute cholecystitis.
3. Etiopathogenesis, Gross, Microscopy and Clinical features of chronic cholecystitis.

Overview

- Definition
- Types →

1. Acute Cholecystitis
2. Chronic Cholecystitis

} Introduction, Etiopathogenesis, Gross, Microscopy, Clinical features

Definition →

Cholecystitis is inflammation of gallbladder

Types →

1. Acute
2. Chronic
3. Acute superimposed on chronic.

- Acute cholecystitis → surgical emergency.
- Chronic cholecystitis → more common

1) Acute Cholecystitis

Introduction →

Begins with obstruction → followed by infection later.

Etiopathogenesis

Types →
a) Acute calculous cholecystitis (90 %)
b) Acute acalculous cholecystitis (10 %)

a) Acute calculous cholecystitis

Obstruction in neck of gallbladder or in cystic duct by a gallstone (Commonest location is in Hartmann's pouch)
↓
Distension of gallbladder
↓
Chemical irritation
↓
Acute inflammation
↓
Secondary bacterial infection (E. coli and Streptococcus faecalis)



b) Acute acalculous cholecystitis

- Previous nonbiliary surgery
- Multiple injuries
- Burns
- Recent childbirth
- Severe sepsis
- Dehydration
- Torsion of the gallbladder
- Diabetes mellitus

Gross

- Gallbladder → distended and tense.
- Serosal surface → coated with fibrinous exudate with congestion and haemorrhages.
- Mucosa → bright red.
- Lumen → filled with pus mixed with green bile.



Microscopy

- Wall of the gallbladder shows marked inflammatory oedema, congestion and neutrophilic exudate.
- Gangrenous cholecystitis → Abscesses in wall and gangrenous necrosis with rupture into peritoneal cavity

Clinical Features

- Severe pain in upper abdomen
- Features of peritoneal irritation such as guarding and hyperaesthesia.
- Fever
- Leucocytosis with neutrophilia
- Slight jaundice



2) Chronic Cholecystitis

Introduction →

Commonest type of clinical gallbladder disease.

Etiopathogenesis

Supersaturation of bile with cholesterol
↓
Predisposes to
↓
Chronic Inflammation

Gross

- Gallbladder is generally contracted but may be normal or enlarged
- Wall of gallbladder is thickened
- C/S → grey-white due to dense fibrosis
- Mucosal folds intact, thickened, or flattened and atrophied.
- Lumen→ contains multiple mixed stones or a combined stone.



Microscopy

1. Penetration of the mucosa deep into wall of gallbladder up to muscularis layer → Rokitansky-Aschoff 'sinuses.
2. Variable degree of chronic inflammatory reaction, consisting of lymphocytes, plasma cells and macrophage in lamina propria and subserosal layer.
3. Variable degree of fibrosis in subserosal and subepithelial layers.



Clinical Features

- Abdominal distension or epigastric discomfort, especially after a fatty meal.
- Tenderness over the right upper abdomen.
- Constant dull ache in the right hypochondrium and epigastrium
- Nausea and flatulence

	Acute Cholecystitis	Chronic Cholecystitis
Introduction		
Etiopathogenesis		
Gross		
Microscopy		
Clinical features		

(17.15) GALL BLADDER CARCINOMA

University Exams

Long Questions

1. Discuss etiology, gross, microscopic findings, clinical features and spread of Gall bladder carcinoma

Short Questions

1. Gross and microscopy of Gall bladder carcinoma

Overview

- Introduction
- Age
- Gender
- Etiology
- Site
- Gross
- Microscopy
- Clinical Features
- Spread

Introduction →

Remain undetected until time it is widely spread and rendered inoperable.

Age →

Peak incidence in 7th decade of life.

Gender →

F > M (ratio 4:1)

Etiology

1. Cholelithiasis and cholecystitis →
    - Present in about 75% cases of gallbladder cancer.
2. Chemical carcinogens
    - Methyl cholanthrene, various nitrosamines and pesticides.
    - Workers engaged in rubber industry → higher incidence of gallbladder cancer.
3. Miscellaneous
    - Patients who have undergone previous surgery on biliary tract
    - Patients with inflammatory bowel disease (ulcerative colitis and Crohn's disease).

Site

Fundus > Neck



Gross

2 types →

1. Infiltrating type → Irregular area of diffuse thickening and induration of gallbladder wall.
2. Fungating type → Irregular, friable, papillary or cauliflower-like growth into lumen of gallbladder

Infiltrating type Fungating type

Microscopy

1. 90% gallbladder cancers→Adenocarcinomas papillary or infiltrative, well-differentiated or poorly-differentiated
2. 5% gallbladder cancers → squamous cell carcinomas arising from squamous metaplastic epithelium.
3. 5% gallbladder cancers → Adenosquamous carcinoma.



Clinical Features

- Pain
- Jaundice
- Noticeable mass
- Anorexia
- Weight loss.

Spread

- Direct spread → spread to liver, peritoneal surfaces, cystic duct and adjacent bile ducts.
- Lymphatic spread→ spread to perihilar lymphatics and portal-hepatic lymph nodes
- Blood spread → Liver and lungs

(17.16) PANCREATITIS

University Exams

Short Questions

1. Etiology, morphology, Clinical features and Complications of acute pancreatitis.
2. Etiology, morphology, Clinical features and Complications of chronic pancreatitis.

Overview

- Definition
- Types →

1. Acute pancreatitis
2. Chronic Pancreatitis

} Introduction, Etiology, Gross, Microscopy, Clinical features, Complications

Definition →

Pancreatitis is infammation of pancreas with acinic cell injury.

Types →

1. Acute pancreatitis
2. Chronic pancreatitis

1) Acute pancreatitis

Introduction →

Premature activation of trypsin
↓
Activation of other pancreatic enzymes
↓
Autodigestion of pancreatic parenchyma
↓
Liquefactive haemorrhagic necrosis of
pancreas and fat necrosis of the peripancreatic fat
↓
Acute pancreatitis

Etiology

- Alcohol and Gallstones (Most common)
- Trauma
- Steroids
- Mumps
- Autoimmune disease
- Scorpion sting
- Hypercalcemia
- Hypertriglyceridemia > 1000 mg/dL)
- ERCP, Drugs (e.g. Sulpha drug and protease inhibitors).
- Mutations in the cationic trypsinogen (PRSS1) and trypsin inhibitor (SPINK1) genes.

Gross →

- Focal hemorrhage and liquefication accompanied by chalky, white yellow fat necrosis of adjacent adipose tissue.

Microscopy →

- Liquefactive necrosis of the pancreatic parenchyma with acute inflammation
- Enzymatic fat necrosis.



Clinical features

- Acute epigastric abdominal pain radiating to the back.
- Nausea, vomiting
- Periumbilical and flank hemorrhage due to spread of necrosis into the periumbilical soft tissue and retroperitoneum.

Complications

1. Shock due to peripancreatic hemorrhage and fluid sequestration.
2. Pancreatic pseudocyst → formed by fibrous tissue surrounding Liquefactive necrosis
3. Pancreatic abscess → formed due to E. coli.
4. Organ failure (ARDS, shock, renal failure).
5. Hypocalcemia due to precipitation of Ca2+ soaps.

Lab investigations

1. Elevated serum lipase (more specific for pancreatic damage).
2. Elevated amylase level
3. Hypocalcemia due to consumption of calcium during saponification during fat necrosis.

Diagnostic criteria:

Diagnosis is done by 2 of 3 criteria:

- Acute epigastric pain often radiating to the back
- Serum amylase or lipase (Lipase is more specific) to 3x upper limit of normal
- Characteristic imaging findings.

2) Chronic Pancreatitis

Introduction →

Repeated episodes of acute pancreatitis.
↓
Chronic inflammation, atrophy, and fibrosis of pancreas

Etiology

a) Alcohol in adults.
b) Cystic fibrosis in children.

- Hereditary pancreatitis → mutations in pancreatic trypsinogen gene (PRRS1) or SPINK1 gene encoding a trypsin inhibitor

Gross →

Pancreas will be firm, white, and fibrotic.

Microscopy →

Extensive fibrosis with parenchymal atrophy and chronic inflammation.



Clinical features

- Epigastric abdominal pain radiating to back.
- Pancreatic insufficiency → malabsorption with steatorrhea and fat -soluble vitamin deficiencies.
- Secondary diabetes mellitus → due to destruction of islets.
- Increased risk for pancreatic carcinoma
- Dystrophic calcification of pancreatic parenchyma on imaging; contrast studies reveal a 'chain of lakes' pattern due to dilatation of pancreatic ducts.

Complications

1. Diabetes mellitus
2. Pancreatic insufficiency with steatorrhoea and malabsorption
3. Formation of pancreatic pseudocysts



	Acute pancreatitis	Chronic Pancreatitis
Introduction		
Etiology		
Gross		
Microscopy		
Clinical features		
Complications		

(17.17) PANCREATIC TUMOURS

University Exams

Short Questions

1. Gross and Microscopy of Pancreatic Carcinoma
2. Risk factors, Molecular cytogenetics and morphology of Pancreatic Carcinoma

Overview

- Introduction
- Age
- Risk factors
- Molecular cytogenetics
- Site
- Gross
- Microscopy
- Clinical Features
- Tumor markers
- Prognosis

Introduction →

- Infiltrating ductal adenocarcinoma of exocrine pancreas
- most frequent neoplasm of the pancreas

Age →

60 to 80 years

Risk factors →

a) Smoking
b) Age> 50 years
c) Diabetes mellitus
d) Chronic pancreatitis (especially> 20 years)
e) Family history of pancreatic cancer (e.g., BRCA2 mutations account for 10% of pancreatic cancer in Ashkenazi Jews).

Molecular cytogenetics

4 genes are most commonly affected by somatic mutations→

1. KRAS → Overall most common gene mutation
2. CDKN2A/p16 → most commonly inactivated tumor suppressor gene
3. SMAD4
4. Tp53

Site →

Head of pancreas > body > tail



Gross →

Carcinoma of head of pancreas →

- small, homogeneous, poorly-defi ned, grey-white mass
- Extends into the ampulla of Vater, common bile duct and duodenum, producing obstructive biliary symptoms and jaundice

Carcinomas of body and tail→

- Large and irregular masses
- Infiltrate transverse colon, stomach, liver, spleen and regional lymph nodes.

Microscopy →

- Pancreatic adenocarcinoma arises from duct epithelium.
- Tumor desmoplasia
- Perineural invasion

Clinical Features

- Abdominal pain radiating to back.
- Weight loss (due to malabsorption and anorexia).
- Obstructive jaundice with pale stools
- Palpable nontender gallbladder
- Courvoisier sign → Obstructive jaundice with palpable, nontender gallbladder
- Migratory thrombophlebitis (Trousseau syndrome) → swelling, erythema, and tenderness in extremities

Tumor markers

1. CA19-9
2. CEA

Prognosis

- Very poor prognosis
- 1-year survival is< 10%.

(18.01) NORMAL ANATOMY AND HISTOLOGY

Overview

- Normal kidney → Anatomy
- Nephron
- Glomerulus
- Filtration Barrier
- Normal kidney → Histology

Normal kidney → Anatomy

- Bean-shaped paired organs
- Surrounded by a thin fibrous capsule
- **Hilum** of kidney → situated at midpoint on medial aspect → artery, vein,lymphatics and ureter are located



Nephron

- Parenchyma is composed of approx one million microstructures called **nephrons**
- 5 major parts →

    1. Glomerular capsule (glomerulus and Bowman's capsule)
    2. Proximal convoluted tubule (PCT)
    3. Loop of Henle
    4. Distal convoluted tubule (DCT)
    5. Collecting ducts





Glomerulus

- Invagination of blind end of PCT
- Has a **capillary tuft** fed by **afferent arteriole** and drained by **efferent arteriole**



Filtration Barrier

1. Fenestrated endothelial cells lining capillary loops.
2. Glomerular basement membrane (GBM) consists of 3 layers →

    - The central lamina densa
    - Lamina rara interna on endothelial side of capillary
    - Lamina rara externa on visceral epithelial side of capillary.

3. Podocyte (visceral epithelial cells) with Filtration slit pores in between foot processes





Glomerulus – schematic



Normal kidney → Histology



(18.02) NEPHRITIC VERSUS NEPHROTIC SYNDROME

University Exams

Long Questions

1. Describe etiopathogenesis of nephrotic syndrome along with characteristic blood and urine findings.
2. What is nephrotic syndrome? Enlist important causes of nephrotic syndrome in pediatric age group. Add a note on its pathophysiology.

Short Questions

1. Definition, Features and causes of nephritic syndrome.
2. Definition, Features and causes of nephrotic syndrome.
3. Enumerate differences between nephritic and nephrotic syndrome.

Very Short Questions

1. List examples of nephritic syndrome.
2. Four examples of nephrotic syndrome.
3. Define Azotaemia and Uremia.

Overview

Nephritic Syndrome →

- Features →
  1. Microscopic haematuria
  2. Mild proteinuria
  3. Sodium and water retension
  4. Hypertension
  5. Oedema
  6. Oliguria
- Examples
- Azotaemia
- Uremia

Nephrotic Syndrome

- Features →
  1. Heavy proteinuria
  2. Hypo albuminaemia
  3. Oedema
  4. Hyperlipidaemia
  5. Lipiduria
  6. Hypercoagulability
- Examples

Nephritic Syndrome

Features

1) Haematuria →

- Urine smoky appearance
- RBCs detectable by microscopy or by chemical testing for haemoglobin.
- Presence of RBC casts in urine is classical feature of nephritic syndrome

2) Proteinuria →

- Mild (less than 3 gm per 24 hrs)
- Non-selective

3) Sodium and water retention

4) Hypertension →

- Depending upon severity of the glomerular disease
- Results from sodium and water retention

5) Oedema →

- Mild
- Results from sodium and water retention

6) Oliguria →

- Reflects severity of glomerular involvement.

Examples

Primary Glomerulonephritis
1. Post-streptococcal GN
2. Rapidly progressive GN → Most common cause
3. Membranoproliferative GN
4. Focal and diffuse proliferative GN
5. IgA nephropathy

Systemic Diseases
1. SLE
2. Polyarteritis nodosa
3. Wegener's granulomatosis
4. Henoch-Schonlein purpura
5. Cryoglobulinaemia

Nephrotic Syndrome

Features

1) Heavy proteinuria

- Severe → protein loss > 3 gm per 24 hrs
- Selective → loss of low molecular weight proteins.eg. albumin

2) Hypoalbuminaemia →

- Due to urinary loss of albumin
- Plasma albumin level is 1 to 3 gm/dl (normal 3.5 to 5.5 gm/dl)
- A – G ratio reversed

Concentration of other proteins in plasma fall→
- Immunoglobulins → vulnerable to infections → Spontaneous bacterial perotinitis is most common infection caused by pneumococcus (most common) and E. coli (2nd most common)
- Antithrombin III→ vulnerable to thrombosis.
- Transferrin → causing microcytic hypochromic anemia
- Cholecalciferol binding protein → causing hypocalcemia
- Thyroxine binding globulin →causing decreased thyroxine level

- Fibrinogen → increased due to increased hepatic synthesis

3) Oedema →

- Due to fall in osmotic pressure (hypo albuminaemia)
- (not due to sodium and water retension)

4) Hyperlipidaemia →

- Liver faced with stress of massive protein synthesis in response to heavy urinary protein loss → increased synthesis of lipoproteins.

5) Lipiduria →

- Following hyperlipidaemia

6) Hypercoagulability →

- Predisposes to renal vein thrombosis. It is due to →
- Increased urinary loss of antithrombin III
- Hyperfibrinogenaemia

Examples

Primary Glomerulonephritis
1. Minimal change disease (most common in children)
2. Focal segmental glomerulosclerosis
3. (most common in adults)
4. Membranoproliferative GN
5. Membranous GN
6. Focal and diffuse proliferative GN
7. IgA nephropathy

Systemic Diseases
1. Diabetes mellitus
2. Amyloidosis
3. SLE

Remember
- Cause of edema in nephrotic syndrome: Hypoalbuminemia
- Cause of edema in nephritic syndrome: Salt (sodium) and water retention

	Feature	Nephritic Syndrome	Nephrotic Syndrome
1	Proteinuria	Mild (< 3gm per 24 hrs)	Heavy (> 3 gm Per 24 Hrs)
2	Hypoalbuminaemia	Uncommon	Present
3	Oedema	Mild, in loose tissues	Marked, generalise peripheral
4	Mechanism of oedema	Na^+ and water retention	↓ plasma osmotic pressure
5	Haematuria	Present, microscopic	Absent
6	Hypertension	Present	Present in advance disease
7	Hyperlipidaemia	Absent	Present
8	Lipiduria	Absent	Present
9	Oliguria	Present	Present in advance disease
10	Hypercoagulability	Absent	Present

Azotaemia → Biochemical abnormality characterised by elevation of BUN and creatinine levels

Uremia → Azotemia + clinical signs and symptoms

(18.03) GLOMERULONEPHRITIS PATHOGENESIS AND CLASSIFICATION

University Exams

Long Questions

1. Describe briefly immunopathogenesis of glomerulonephritis.

Short Questions

1. Discuss types of Antibody-Mediated Glomerular Injury
2. Location of deposits in glomerulonephritis.
3. Write Classification of glomerulonephritis.

Overview

- Pathogenesis of Glomerulonephritis →

    1. Antibody-Mediated Glomerular Injury →

        a) In situ immune-complex deposition → Fixed Ag
        b) In situ immune-complex deposition → Planted Ag
        c) Circulating immune complex deposition

    2. Cell-mediated Glomerular Injury
    3. Alternative Complement Pathway

- Location of deposits
- Classification of Glomerular Injury →

    1. Primary Glomerulonephritis
    2. Secondary Systemic Glomerular Diseases
    3. Hereditary Nephritis



Pathogenesis of Glomerulonephritis



1) Antibody-Mediated Glomerular Injury

Glomerular disease result from deposits of immune complexes (Ag - Ab complexes)

a) In situ immune-complex deposition → Fixed Ag, Planted Ag.
b) In situ immune-complex deposition → Planted Ag
c) Circulating immune complex deposition

a) In situ immune-complex deposition → Fixed Ag

Anti-GBM antibody induced nephritis or Goodpasture syndrome

Intrinsic fixed antigens
↓
components of GBM
(non-collagenous domain of the alpha 3 chain of collagen type IV.)
↓
Auto-antibodies formed in blood
↓
Ag –Ab complexes formed in kidney itself
↓
Glomerular injury



b) In situ immune-complex deposition → Planted Ag

Ag not normally present in glomerulus but are planted there → eg. bacterial, viral and parasites and drugs
↓
Ab formed in blood
↓
Ag –Ab complexes formed in kidney itself
↓
Glomerular injury



c) Circulating immune complex deposition

Ag in blood → endogenous (e.g. in SLE) or exogenous (e.g. Hepatitis B virus, Treponema pallidum, Plasmodium falciparum, tumour antigens)

Ab formed in blood
↓
Ag –Ab complexes formed in **circulation**
↓
Trapped in glomeruli
↓
Combine with complement
↓
Glomerular injury



2) Cell-mediated Glomerular Injury

(Delayed-type Hypersensitivity)

Activated T cells (CD4 and CD8)
↓
Releases Cytokines
↓
stimulate cytotoxicity
↓
Glomerular injury

No immune complex deposition → pauci immune mechanism

3) Alternative Complement Pathway

Activation of alternate pathway of complement
↓
C3 and properdin deposited in glomeruli without immunoglobulin deposits
↓
Glomerular injury



Location of deposits (seen on EM)

1. Subepithelial deposits
2. Epimembranous deposits
3. Subendothelial deposits
4. Mesangial deposits



Classification of Glomerular Injury

1. Primary Glomerulonephritis
2. Secondary Systemic Glomerular Diseases
3. Hereditary Nephritis

1) Primary Glomerulonephritis

1. Acute GN / Post-streptococcal GN
2. Rapidly progressive GN
3. Minimal change disease
4. Membranous GN
5. Membrano-proliferative GN
6. Focal and diffuse proliferative GN
7. Focal segmental glomerulosclerosis (FSGS)
8. IgA nephropathy
9. Chronic glomerulonephritis

2) Secondary Systemic Glomerular Diseases

1. Lupus nephritis (SLE)
2. Diabetic nephropathy
3. Amyloidosis
4. Polyarteritis nodosa
5. Wegener's granulomatosis
6. Goodpasture's syndrome
7. Henoch-Schonlein purpura
8. Systemic infectious diseases (bacterial e.g. bacterial endocarditis, syphilis, leprosy; viral e.g. HBV, HCV, HIV; parasitic e.g. falciparum malaria, fila riasis)
9. Idiopathic mixed cryoglobulinaemia

3) Hereditary Nephritis

1. Alport's syndrome
2. Fabry's disease
3. Nail-patella syndrome

(18.04) PRIMARY GLOMERULONEPHRITIS (APGN, RPGN, MCD, MGN, MPGN, FSGN, FSGS, IGA NEPHROPATHY, CHRONIC GN)

University Exams

Long Questions

1. A 50-year-old male presented with facial puffiness and proteinuria 5 g/day. His serum albumin was found to be 2.0 g/dl. His urine showed fat globules.

   a) What is your diagnosis?
   b) Discuss in detail the causes of this clinical syndrome.

2. Classify glomerulonephritis. Describe subtypes, gross & microscopic features of rapidly progressive glomerulonephritis.
3. Describe the etiopathogenesis, light microscopy, immunofluorescence, electron microscopic findings and clinical course poststreptococcal glomerulonephritis.
4. Enumerate major Glomerular Syndrome. Describe immunopathogenesis, gross and microscopic picture of acute proliferative glomerulonephritis.
5. Classify glomerulonephritis. Describe etiopathogenesis, gross & microscopic features of IgA Nephropathy
6. A 8-year-old boy was admitted with history of puffiness of face and passing of red colored urine, following upper respiratory tract infection 2 weeks earlier. Urine examination showed moderate proteinuria and hematuria.

   a) What is your most probable diagnosis?
   b) Describe the etiopathogenesis of the condition.
   c) Give the morphology of the organ affected.
   d) List the investigations that you would do to confirm diagnosis.

Short Questions

1. Classification and pathogenesis of rapidly progressive glomerulonephritis.
2. Goodpasture syndrome.
3. Etiology and Morphology (light microscopy) of minimal lesion glomerulonephritis/ minimal change disease.
4. Etiopathogenesis and pathology of membranous glomerulonephritis.
5. Classify glomerular diseases and describe the morphology of membranoproliferative glomerulonephritis.
6. Classify glomerular diseases and describe gross and microscopic findings in acute glomerulonephritis.
7. Microscopy of chronic glomerulonephritis.

Very Short Questions

1. Etiology of crescentic glomerulonephritis.
2. Classification of membranoproliferative glomerulonephritis.
3. Morphology of Focal segmental glomerulosclerosis.

Overview

1. APGN
2. RPGN
3. MCD
4. Membranous GN
5. MPGN
6. FSGN & FSGS
7. IgA Nephropathy
8. Chronic GN

} Introduction, Etiopathogenesis, Gross, Light Microscopy, Electron Microscopy, Immunofluorescence microscopy, Clinical feature

1) Acute Proliferative GN (APGN)

Introduction →

- Also known as **Poststreptococcal Glomerulonephritis (PSGN)**

Etiopathogenesis

Group A β-haemolvtic streptococci
↓
Pharyngitis or skin infection
↓
Comes to kidney through blood
↓
Bacteria gets deposited on visceral podocytes (Planted Ag)
↓
Antibodies against bacteria are formed in - situ
↓
Ag - Ab complexes are formed in - situ with complement deposition
↓
Subepithelial deposits are seen on EM

Gross

- Petechial haemorrhages on surface→ **flea-bitten kidney**



Light microscopy

- Glomeruli → Enlarged and hypercellular





Electron microscopy

- Subepithelial deposits → humps

Immunofluorescence microscopy



- IgG and C3 → granular deposits

Clinical Features

- Nephritic syndrome

2) Rapidly Progressive Glomerulonephritis (RPGN)

Introduction →

- Also known as crescentic GN → Characterised by formation of crescents

Etiopathogenesis

1. Type I → Anti-GBM type→ In situ deposition against fixed Ag
2. Type II → Post-infectious RPGN → In situ deposition against planted Ag
3. Type III → Pauci-immune RPGN → Cellular immunity→ ANCA positive

Type I RPGN: Anti-GBM disease

- **Goodpasture's syndrome →** characteristic example of Type I RPGN



Light microscopy

Crescents
- Pathogmatic
- Formed by proliferation of parietal cells and by migration of monocytes and macrophages into urinary space.
- Neutrophils and lymphocytes may be present.



Electron microscopy

- Type I → Along the GBM
- Type II → Sub epithelial
- Type III → No deposits



TYPE I TYPE II

Immunofluorescence microscopy

- Type I → linear pattern→ containing IgG + C3
- Type II →Granular pattern → consisting of IgG + C3
- Type III → No deposits



Type I Type II

Clinical Features

- Nephritic syndrome

3) Minimal Change Disease (MCD)

Introduction →

Also known as →
- Minimal Change Disease (MCD) → Nephrotic syndrome with no apparent change in glomeruli by light microscopy

- Foot process disease → EM → flattened podocytes
- Nil deposit disease → No deposits

- Age → Children < 16 years
- Most common cause of nephrotic syndrome in children (80% cases)

Etiopathogenesis

- Cell mediated

Gross

- Kidneys → Normal size and shape

Light microscopy

Glomeruli →

- No apparent abnormality
- Slight increase in the mesangial matrix





Electron microscopy

- Diffuse flattening / effacement/ Fusion / Obliteration of foot processes of visceral epithelial cells (podocytes)
- **No deposits**

Immunofluorescence microscopy

- No deposits

Clinical features →

- Nephrotic syndrome with massive and highly selective proteinuria

Prognosis →

- Characteristically responds to steroid therapy.

4) Membranous Glomerulonephritis

Introduction →

- Also known as → Membranous GN → characterised by widespread thickening of glomerular capillary wall

Etiopathogenesis

Antigen is gp 330
(component of podocyte)
↓
Fixed Ag
↓
Ab are formed against it
↓
Form immune complexes in podocytes
↓
In situ deposition against fixed Ag
↓
ARF

Gross

- Kidneys → Enlarged, pale and smooth.



Light microscopy

- Diffuse thickening of GBM
- Duplication of GBM
- No cellular proliferation

Electron microscopy

- Subepithelial deposits → spikes



Immunofluorescence microscopy

- IgG with C3



Clinical Features

- Nephrotic syndrome in an adult.

5) Membrano Proliferative Glomerulonephritis (MPGN)

Introduction →

- Also known as Mesangiocapillary GN

Etiopathogenesis

1. Type I or classic form → 70% cases.
    - Immune deposits in **subendothelial position**
2. Type II or dense deposit disease → 30% cases
    - Example of alternate pathway disease
    - Immune deposits along **GBM**
3. Type III → rare
    - Features of type I MPGN and membranous nephropathy with systemic diseases



Light microscopy

- **Enlarged** with accentuated lobular pattern due to mesangial cellular proliferation and increase in mesangial matrix.
- Splitting, or 'tram track' appearance

Electron microscopy

- Type I → Subendothelial deposits
- Type II → Intramembranous deposition
- Type III → Anywhere



6) FSGN and FSGS

GN →
- Focal → Some, but not all, glomeruli are involved
- Segmental → In affected glomeruli, only a portion of capillary tuft is involved



7) IgA Nephropathy

Introduction →

- Also known as → Berger's Disease
- Most common type of glomerulonephritis world wide
- Characterised by aggregates of IgA deposited in mesangium.
- Occurs after an upper respiratory tract infection or gastrointestinal infection

Etioathogenesis

- Due to exclusive mesangial deposits of IgA an→ entrapment of immune complexes from circulation in mesangium.
- Due to Presence of C3 and properdin in the mesangial deposits → alternate complement pathway
- HLA-B35 association → genetically-determined abnormality

Light microscopy→

Various patterns→

- Focal proliferative GN,
- Focal segmental glomerulosclerosis,
- Membranoproliferative GN

Electron microscopy →

- Granular electron-dense deposits in mesangium

Immunofluorescence microscopy→

- Mesangial deposits of IgA, with or without IgG, and with C3 and properdin.

Clinical Features

- Mainly Nephritic syndrome
- Occasionally nephrotic syndrome

8) Chronic Glomerulonephritis

Introduction →

- Also known as → End-Stage Kidney, Chronic Kidney Disease
- Chronic GN is final stage of a variety of glomerular diseases which result in irreversible impairment of renal function.



Gross

- Kidneys → small and contracted
- Cortical surface → diffusely granular
- On cut section→ cortex is narrow and atrophic, medulla is unremarkable.

Light Microscopy

1. Glomeruli
    - Reduced in number
    - show completely hyalinised tufts
    - Acellular, eosinophilic masses which are PASpositive.
2. Tubules
    - Atrophy of tubules
    - Hyaline-droplets
3. Interstitium → Fibrosis and chronic inflamatory cells
4. Vessels → Arteriolar sclerosis



	Introduction	Etiopathogenesis	Gross	LM	EM	IFM	Clinical feature
APGN							
RPGN							
MCD							
Membranous GN							
MPGN							
FSGN							
FSGS							
IgA Nephropathy							
Chronic GN							

(18.05) DIABETIC NEPHROPATHY

University Exams

Short Questions

1. Renal lesions in diabetes mellitus
2. Kimmelstiel-Wilson lesion.
3. Diffuse and Nodular glomerulosclerosis in diabetes mellitus
4. Armanni-ebstein Lesions

Overview

- Introduction
- Morphology →

    1. Diabetic Glomerulosclerosis →

        a) Diffuse glomerulosclerosis
        b) Nodular glomerulosclerosis

    2. Vascular Lesions
    3. Tubular Lesions (Armanni-ebstein Lesions)

Introduction

- Secondary Systemic Glomerular Diseases
- Renal involvement is a complication of diabetes mellitus
- DM type 1 > DM type 2

Morphology



1) Diabetic Glomerulosclerosis

a) Diffuse glomerulosclerosis

- Most common.
- All parts of glomeruli

1. **Thickening of the GBM →** earliest morphological abnormality
2. **Diffuse increase in mesangial matrix with mild proliferation of mesangial cells**

3. **Capsular drop** → eosinophilic hyaline thickening of parietal layer of Bowman's capsule and bulges into glomerular space.
4. **Fibrin cap** → homogeneous, brightly eosinophilic material appearing on wall of a peripheral capillary



DIFFUSE GLOMERULOSCLEROSIS

b) Nodular glomerulosclerosis

- Also called as Kimmelstiel- Wilson (KW) lesions or intercapillary glomerulo sclerosis
- Pathognomonic (most specific)

Nodule →
- Located within a lobule of glomerulus
- Ovoid or spherical
- Laminated
- Hyaline, acellular mass
- PAS-positive



NODULAR GLOMERULOSCLEROSIS

2) Vascular Lesions →

Hyaline arteriolosclerosis

3) Tubular Lesions (Armanni-ebstein Lesions) →

Epithelial cells of the PCT develop extensive glycogen deposits appearing as vacuoles

(18.06) PYELONEPHRITIS

University Exams

Long Questions

1. Describe etiopathogenesis, gross,microscopy, Clinical features and Complications of chronic pyelonephritis.

Short Questions

1. Etiopathogenesis and types of Pyelonephritis.
2. Complications of acute pyelonephritis.
3. Xanthogranulomatous pyelonephritis.
4. Gross, microscopy and clinical features of chronic pyelonephritis.
5. Gross microscopy and clinical features of acute pyelonephritis.

Overview

- Definition
- Types →

1. Acute Pyelonephritis
2. Chronic Pyelonephritis

} Introduction, Etiopathogenesis, Gross, Microscopy, Clinical features, Complications

- Xanthogranulomatous pyelonephritis (XPN)

Definition

- Inflammatory process that predominantly involves renal interstitial tissue and tubular damage.

Types

1. Acute Pyelonephritis
2. Chronic Pyelonephritis

1) Acute Pyelonephritis

Introduction →

- Acute suppurative inflammation of kidney caused by pyogenic bacteria

Etiopathogenesis

- Bacteria → Escherichia coli (in 90% of cases) > Enterobacter > Klebsiella > Pseudomonas > Proteus
- After renal transplant → Polyomavirus BK (BKV)

2 routes→

a) Ascending infection
b) Haematogenous infection



a) Ascending infection

- Most common route of infection

Organism's inhabitants of colon
↓
Faecal contamination of urethral orifice
↓
Endogenous infection
↓
Ascends upwards
↓
Acute Pyelonephritis

More common in females→

- Shorter urethra in females
- Absence of prostatic secretions
- Urethral trauma during sexual intercourse → H**oneymoon pyelitis**

b) Haematogenous infection

- Blood-borne spread of infection

Gross

- Enlarged and swollen kidney that bulges on section
- C/S → yellow-white abscesses with a haemorrhagic rim

Microscopy

1. Interstitium →
    - Patchy interstitial suppurative inflammation
    - Large number of neutrophils in the interstitial tissue and bursting into tubules
2. Tubules →Tubular necrosis
3. Glomreuli → **Resistant**



Clinical Features

- Chills
- Fever
- Loin pain
- Lumbar tenderness

- Dysuria
- Frequency of micturition
- Urine will show abundance of bacteria, pus cells and pus cell casts in the urinary sediment.



Complications

1. Papillary necrosis
2. Pyonephrosis → Kidney transforms into a multilocular sac filled with pus
3. Perinephric abscess → Abscesses in kidney extend through capsule into perinephric tissue

2) Chronic Pyelonephritis

Introduction →

- Chronic tubulointerstitial disease resulting from repeated attacks of inflammation and scarring

Etiopathogenesis

2 types→

a) Reflux nephropathy (Most common)
b) Obstructive pyelonephritis

a) Reflux nephropathy

Congenital absence or shortening of intravesical portion of ureter
↓
Ureter is not compressed during the act of micturition
↓
Vesicoureteric reflux
↓
Reflux of urine from bladder into one or both ureters during micturition
↓
Urine is forced into renal tubules and interstitium
↓
Damage to kidney and Scar formation
↓
Chronic Pyelonephritis



b) Obstructive pyelonephritis

Obstruction to outflow of urine at any levels
↓
Increase in pressure inrenal pelvis
↓
Urine is forced into renal tubules and interstitium
↓
Damage to kidney and Scar formation
↓
Chronic Pyelonephritis





Gross

- Kidneys → small and contracted
- Surface of kidney → irregularly scarred
- Scars → U-shaped depressions on the cortical surface.
- Blunting and dilatation of calyces

Microscopy

1. Interstitium
    - Chronic interstitial inflammatory reaction composed of lymphocytes, plasma cells and macrophages
    - Interstitial fibrosis
2. Tubules
    - Atrophy and dilatation
    - Dilated tubules → contain eosinophilic colloid casts → thyroidisation of tubules
3. Glomeruli
    - Hyalinisation of glomeruli



Clinical Features

- Patients present with clinical picture of **chronic renal failure**

Xanthogranulomatous pyelonephritis (XPN)

- Unusual variant of chronic pyelonephritis.
- Most common cause → obstruction due to infected renal stones.

Gross →
Yellow, lobulated masses diffusely replace renal architecture.

Microscopy →
Massive destruction of kidney due to granulomatous tissue containing lipid-laden macrophages



(18.07) RENAL CALCULUS AND HYDRONEPHROSIS

University Exams

Short Questions

1. Etiopathogenesis and morphology of various types of renal calculi.
2. Nephrolithiasis → types and pathogenesis.
3. Discuss etiology, gross, stages and microscopy of Hydronephrosis.

Very Short Questions

1. Enumerate types of renal stones.
2. Staghorn calculi.
3. Struvite stones.
4. Write 4 conditions when renal stones occur.
5. Causes of hydronephrosis.
6. Gross morphology of kidney in hydronephrosis.

Overview

- Renal calculus
  - Introduction
  - Age
  - Sex
  - Clinical features
  - Types →
    1. Calcium containing
    2. Mixed (struvite)
    3. Uric acid
    4. Cystine stones

    } Introduction, Etiology, Pathogenesis, Morphology
- Hydronephrosis
  - Introduction
  - Causes
  - Gross
  - Stages
  - Microscopy

Renal calculus / Nephrolithiasis / urolithiasis

Introduction →

Formation of urinary calculi at any level of urinary tract.

Age →

2nd to 3rd decades of life.

Sex →

Male: female ratio → 2:1

Clinical features →

- Colicky pain as they pass down along the ureter
- Haematuria

Types

4 types →

1. Calcium containing
2. Mixed (struvite)
3. Uric acid
4. Cystine stones

1) Calcium Stones

Introduction →

Most common (75% of all urinary calculi)

Etiology →

- 50% → idiopathic hypercalciuria without hypercalcaemia.
- 10% → hypercalcaemia and hypercalciuria due to hyperparathyroidism hyperparathyroidism, or a defect in the bowel (i.e. absorptive hypercalciuria), or in the kidney (i.e. renal hyper calciuria).
- 15% → hyperuricosuria with a normal blood uric acid level and without any abnormality of calcium metabolism.
- 25% → cause is unknown → idiopathic calcium stone disease

Pathogenesis →

Imbalance between degree of supersaturation of ions forming stone and concentration of inhibitors in the urine.

Morphology →

- Small
- Ovoid
- Hard with granular rough surface.
- Dark brown due to old blood pigment deposited in them as a result of repeated trauma

2) Mixed (Struvite) Stones

Introduction →

- 15% of urinary calculi
- Made of magnesium-ammonium-calcium phosphate
- Mixed stones → struvite stones or triple phosphate stones

Etiology →

Urinary infection with urea-splitting organisms like Proteus

Pathogenesis →

Bacterias producing urease (Proteus)
↓
split of urea and form ammonia
↓
Alkaline urinary pH
↓
Precipitate ions
↓
struvite stones

Morphology →



- Yellow-white or grey
- Soft and friable
- Irregular in shape
- Staghorn stone → large, solitary stone that takes shape of renal pelvis

3) Uric Acid Stones

Introduction →

- 6% of urinary calculi
- Uric acid calculi → radiolucent (unlike radio-opaque calcium stones)

Etiology →

Hyperuricaemia and hyperuricosuria →
- Primary gout or secondary gout
- Myeloproliferative disorders (e.g. in leukaemias)
- Chemotherapy
- Administration of uricosuric drugs (e.g. salicylates, probenecid).

Pathogenesis →

- Solubility of uric acid at pH of 7 → 200 mg/dl
- Solubility of uric acid at pH of 5 → 15 mg/dl

Urine becomes acidic
↓
Solubility of uric acid in urine decreases
↓
Precipitation of uric acid crystals
↓
Formation of uric acid stones

Morphology →

- Yellowish-brown
- Smooth
- Hard
- Cut section → laminated structure

4) Cystine Stones

Introduction →

2% of urinary calculi.

Etiology →

- Genetically-determined defect in the transport of cystine
- Cystinuria

Pathogenesis →

Excessive excretion of cystine
↓
Least soluble of naturally-occurring amino acids
↓
Formation of crystals
↓
Cystine calculi

Morphology →

- Yellowish and waxy
- Small
- Rounded
- Smooth

Type	Incidence	Etiology	Pathogenesis
Calcium stones	75%	Hypercalciuria with or without hypercalcaemia	Supersaturation of ions in urine, alkaline pH of urine; low urinary volume, oxaluria and hyperuricosuria
Mixed (struvite) stones	15%	Urinary infection with urea-splitting organisms like Proteus	Alkaline urinary pH produced by ammonia from splitting of urea by bacterially produced urease
Uric acid stones	6%	Hyperuricosuria with or without hyperuricaemia (e.g. gout)	Acidic urine (pH below 6) decreases the solubility of uric acid in urine and favours its precipitation
Cystine stones	2%	Genetically-determined defect in cystine transport	Cystinuria containing least soluble cystine precipitates as cystine crystals

Hydronephrosis

Introduction →

- Dilatation of renal pelvis and calyces due to partial or intermittent obstruction to outflow of urine
- Hydroureter always accompanies hydronephrosis.
- Unilateral or bilateral


Stenosis
Stone
Pelvic and ureteric tumours
Congenital stenosis
Adhesions
Blood clot
Inflammatory stenosis of ureter
Loop
Ureteric stone
External compression
Bladder cancer
Bladder neck contraction
Prostatic carcinoma
Prostatic hyperplasia
Posterior urethral valve
Meatal stenosis
Stricture
Phimosis

Causes


Renal pelvis
- Calculi
- Tumors
Ureter-intrinsic
- Calculi
- Tumors
Ureter-extrinsic
- Pregnancy
- Tumors
Bladder
- Calculi
- Tumors
- Neurogenic
Prostate
- Hyperplasia
- Carcinoma
Urethra
- Stricture

Gross

Kidneys → enlarged
↓
Extrarenal hydronephrosis → dilatation of renal pelvis medially in the form of a sac
↓
Obstruction persists
↓
Progressive dilatation of pelvis and calyces
↓
Pressure atrophy of renal parenchyma
↓
Intrarenal hydronephrosis → thin rim of renal cortex is stretched over dilated calyces (lobulated appearance)



Stages



Microscopy

- Wall of hydronephrotic sac is thickened due to fibrous scarring and chronic inflammatory cell infiltrate
- Glomerulus sclerosis
- Atrophy of tubules
- Interstitial fibrosis



(18.08) RENAL CHANGES IN HYPERTENSION (NEPHROSCLEROSIS)

University Exams

Short Questions

1. Pathogenesis of essential hypertension.
2. Kidney changes in hypertension.
3. Describe gross and microscopy of Benign nephrosclerosis.
4. Describe gross and microscopy of malignant nephrosclerosis.
5. Differences between Benign Nephrosclerosis and Malignant Nephrosclerosis

Overview

- Hypertension Classification
- Clinical classification
- Renal changes in Hypertension →

  1. Benign Nephrosclerosis
  2. Malignant Nephrosclerosis

  } Introduction, Gross, Microscopy

- Differences between Benign Nephrosclerosis and Malignant Nephrosclerosis

Hypertension Classification

Category	Systolic (Mmhg)	Diastolic (Mmhg)
Normal	<120	and < 80
Prehypertension	120-139	or 80-89
Benign Hypertension		
Stage 1	140-159	or 90-99
Stage 2	>160	or >100
Isolated systolic hypertension	>140	and < 90
Malignant hypertension	>200 (sudden onset)	>140 (sudden onset)

Clinical classification

Benign hypertension

- 90-95% patients
- Moderate elevation of BP and rise is slow over years.

Malignant hypertension

- 5% patients
- Marked and sudden increase of BP to 200/140 mmHg or more
- Papilloedema, retinal haemorrhages and hypertensive encephalopathy

Renal changes in Hypertension

1. Benign Nephrosclerosis
2. Malignant Nephrosclerosis

1) Benign Nephrosclerosis

Introduction →

- Renal disease that occurs in **benign phase of hypertension.**

Gross

- Kidneys → reduced in size and weight
- Surface of kidney is finely granular and shows V-shaped areas of scarring → leather grain kidney

Microscopy

1. Vascular changes→
   a) Hyaline arteriolosclerosis → homo generous and eosinophilic thickening of the wall of small blood vessels.
   b) Intimal thickening → due to proliferation of smooth muscle cells in the intima
2. Glomerulus → Periglomerular fibrosis, glomerular sclerosis
3. Tubules → Tubular atrophy
4. Interstitium → Interstitial fibrosis



2) Malignant Nephrosclerosis

Introduction →

- Renal disease that occurs in **malignant or accelerated hypertension.**

Gross

- Kidneys → enlarged, oedematous
- Petechial haemorrhages on the surface→ flea-bitten kidney



Microscopy

1. Vascular changes →

    a) **Hyperplastic intimal sclerosis** → onionskin proliferation → concentric laminae of proliferated smooth muscle cells
    b) **Necrotising arteriolitis** → vessel wall shows fibrinoid necrosis

2. Glomerulus → Fibrinoid necrosis
3. Tubules → Tubular atrophy
4. Interstitium → Interstitial fibrosis



MALIGNANT NEPHROSCLEROSIS

Features	Benign nephrosclerosis	Malignant nephrosclerosis
Condition	Benign hypertension	Malignant hypertension
Gross appearance of kidneys	Leather grain appearance	Flea bitten appearance (petechial hemorrhage)
	Small or reduced kidneys	Variable size of the kidney
Microscopy	1. Vascular changes a) Hyaline arteriolosclerosis b) Intimal thickening 2. Glomerulus → Periglomerular fibrosis, glomerular sclerosis 3. Tubules → Tubular atrophy 4. Interstitium → Interstitial fibrosis	1. Vascular changes a. Hyperplastic intimal sclerosis → onionskin proliferation b) Necrotising arteriolitis 2. Glomerulus → Fibrinoid necrosis 3. Tubules → Tubular atrophy 4. Interstitium → Interstitial fibrosis
Activity of renin angiotensin system	Normal	↑ed
Proteinuria	Mild	Marked
Sign of malignant hypertension	Absent	Present such as retinopathy encephalopathy
Renal failure	Rare	More common

(18.09) POLYCYSTIC KIDNEY DISEASE

University Exams

Short Questions

1. Types of Polycystic Kidney Disease
2. Cystic disease of kidney.
3. Discuss inheritance, mutation and gross findings in Adult polycystic kidney disease.
4. Discuss inheritance, mutation and gross findings in Infantile polycystic kidney disease.
5. ADPKD and ARPKD

Overview

- Introduction
- Types →

1. ADPKD
2. ARPKD

} Inheritance, Mutation, Laterality, Age of presentation, Gross, Other features

Introduction→

- Major portion of the renal parenchyma is converted into cysts of varying size

Types→

1. Adult Polycystic Kidney Disease → ADPKD
2. Infantile Polycystic Kidney Disease → ARPKD

1) Adult Polycystic Kidney Disease

Inheritance →

Autosomal dominant trait → ADPKD

Mutation→

PKD gene (Polycystein protein)

a) Mutation in PKD-1 gene located on chromosome 16 (85% cases)
b) Mutation in PKD- 2 gene located on chromosome 4 (15% cases)

Laterality →

Bilateral and diffuse.

Age of presentation →

Kidneys are abnormal at birth → symptoms appear in adult life (30 and 50 years)

Gross →

- Kidneys bilaterally enlarged, symmetrically
- Lobulated appearance on external surface
- Cut surface → cysts throughout the renal parenchyma
- Cysts do not communicate with pelvis of kidney (distinguish from hydronephrosis)



Other featutes →

- Cysts of the liver (1/3rd patients)
- Intracranial aneurysms
- Other associated congenital anomalies → cysts in the pancreas, spleen, lungs and other organs.

2) Infantile Polycystic Kidney Disease

Inheritance →

Autosomal recessive trait → ARPKD

Mutation →

Mutation in PKHD1 gene located on in chromosome 6 (Fibrocystin protein)

Laterality →

Bilateral and diffuse

Age of presentation →

Present at birth → death from renal failure in early childhood.

Gross →

- Kidneys bilaterally enlarged
- Smooth external surface
- Cut surface → small, fusiform or cylindrical cysts radiating from medulla to outer cortex → sponge-like appearance



Other featutes →

- Hepatic cysts
- Congenital hepatic fibrosis

	Feature	ADPKD	ARPKD
1	Inheritance	Autosomal dominant	Autosomal recessive
2	Cytogenetic defect	Chromosome 16: ADPKD-1 Chromosome 4: ADPKD-2	Chromosome 6
3	Mutations	PKD 1 gene (85%) PKD 2 gene (15%)	PKHD1
4	Incidence	1:400 to 1:1000	1:20,000
5	Age at presentation	Adults (3rd to 5th decades)	Infancy, perinatal
6	Gross	Symmetric bilateral enlargement Macrocysts	Micro- and macrocysts radiating from medulla to outer cortex → sponge-like
7	Microscopy	Cysts derived from all parts of nephron (glomeruli, tubules)	Cysts from dilated collecting ducts
8	Other manifestations	Intracranial aneurysms, cysts of other organs	Hepatic cysts Congenital hepatic fibrosis

(18.10) TUMOURS OF KIDNEY

University Exams

Long Questions

1. Classify tumors of kidney. Describe pathology of any two.
2. Classify tumors of kidney. Describe etiopathogenesis and morphology of various types of renal cell carcinoma in detail.
3. Classify tumors of kidney. Describe etiopathogenesis, gross, microscopy, clinical features and spread of various types of Wilms' Tumour (nephroblastoma).

Short Questions

1. Renal cell carcinoma → gross and microscopic features.
2. Gross and microscopy of Nephroblastoma
3. Gross and microscopic features of Clear cell renal cell carcinoma.

Very Short Questions

1. Enumerate 4 microscopic variants of renal cell carcinoma.
2. Morphology of Papillary renal cell carcinoma.

Overview

- Classification
- Renal cell carcinoma (RCC) →
  - Introduction
  - Age
  - Gender
  - Gross
  - Classification →
    - a) Clear cell carcinoma
    - b) Papillary Carcinoma
    - c) Chromophobe carcinoma
    - d) Bellini duct carcinoma
  - Clinical Features
  - Spread
  - Prognosis
- Wilms' Tumour
  - Introduction
  - Arises from
  - Age
  - Gender
  - Etiology
  - Gross
  - Microscopy
  - Clinical Features
  - Spread
  - Prognosis

Classification

Benign	Malignant
A. Epithelial Tumours of Renal Parenchyma	
Adenoma Oncocytoma	Adenocarcinoma / **Renal cell carcinoma (RCC)**
B. Epithelial Tumours of Renal Pelvis	
Transitional Cell Papilloma	Transitional cell carcinoma Others (squamous cell carcinoma, adenocarcinoma of renal pelvis, undifferentiated carcinoma of renal pelvis)

C. Embryonal Tumours	
Mesoblastic Nephroma Multicystic Nephroma	Wilms' tumour (nephroblastoma)
D. Non-epithelial Tumours	
Angiocnyolipoma Medullary Interstitial Tumour (Fibroma)	Sarcomas (rare)
F. Metastatic Tumours	

Renal cell carcinoma (RCC)

Introduction →

Also known as Hypernephroma
Most common malignant tumor of kidney.

Age →

50 to 70 years of age

Gender →

M > F

Gross



- Arises from poles of kidney (more common in upper pole)
- Solitary and unilateral tumour
- Large, golden yellow and circumscribed

Classification of RCC

a) Clear cell carcinoma
b) Papillary Carcinoma
c) Chromophobe (oncocytic) renal cell carcinoma
d) Collecting duct (Bellini duct) carcinoma

a) Clear cell type RCC

Incidence → Most common pattern (70% to 80% of RCC)

Arises from → PCT

Microscopy →
- Clear cytoplasm of tumour cells → due to removal of glycogen from cytoplasm during processing of tissues.

Cytogenetic abnormalities →
- VHL (Ch 3)



b) Papillary type RCC



Incidence → 2nd most common

Arises from → DCT

Microscopy →
- Tumour cells arranged in papillary pattern over fibrovascular stalks.
- Psammoma bodies may be seen.

Cytogenetic abnormalities →
- Sporadic: Trisomy of 7,16,17; loss of Y chromosome (Y); t (X:1).
- Hereditary: Trisomy 7.

c) Chromophobe type RCC



Incidence → 3rd common

Arises from → cortical collecting ducts

Microscopy →
- Prominent cell membranes
- Pale eosinophilic cytoplasm
- Perinuclear halo

Cytogenetic abnormalities →
- Hypodiploidy or monosomy due loss of multiple chromosomes: 1,2,6,10,13,17,21, & Y

d) Collecting duct (Bellini duct) carcinoma

Incidence → least common

Arises from → medullary collecting duct.

Microscopy → Cells have prominent fibrotic stroma.

Clinical Features

Triad →

1. Gross haematuria (most common)
2. Flank pain
3. Palpable abdominal mass

Spread

1. Haematogenous route → to lungs (most common) brain and bone
2. Locally → to liver and perirenal lymph nodes.

Prognosis

- 5-year survival rate → 70%

Poor prognostic factors →

1. Presence of metastases
2. Renal vein invasion
3. Higher nuclear grade of the tumour

Wilms' Tumour

Introduction →

- Also known as Nephroblastoma
- Most common abdominal malignant tumour of young children

Arises from →

Embryonic tumour derived from primitive renal epithelial and mesenchymal components

Age →

1 to 6 years of age

Gender →

M = F

Etiology

Mutation → Wilms' tumour associated gene, WT1 gene (located on chromosome 11p13)

Gross

- Large → replacing most of the kidney
- Solitary and unilateral
- On cut section → variegated appearance



Microscopy

Triphasic →

- Blastemal → small, round, blue, anaplastic tumour cells (blast)
- Epithelial → abortive tubules and poorly-formed glomerular structures
- Mesenchymal → elements such as smooth and skeletal muscle, cartilage and bone, fat cells and fibrous tissue



Clinical Features

- Palpable abdominal mass in a child detected by mother

Spread

Haematogenous route → to lungs

Prognosis

- With combination therapy of nephrectomy, post-operative irradiation and chemo therapy → prognosis is good
- 5-year survival → 80-90%

(18.11) BLADDER CARCINOMA

University Exams

Short Questions

1. Etiopathogenesis, Gross and Microscopy of transitional cell carcinoma of bladder.
2. Classify bladder tumours. Write a note on Papillary urothelial neoplasms of low malignant potential (PUNLMP)
3. Transitional cell papilloma of bladder
4. Risk factors of urinary bladder (urothelial) tumors.

Overview

- Classification of bladder tumours
- Urothelial (Transitional Cell) Bladder Tumours →
    - Introduction
    - Age
    - Gender
    - Etiopathogenesis
    - Gross
    - Microscopy →
        1. Papilloma
        2. PUNLMP

        a) Low-grade TCC
        b) High-grade TCC

        3. Invasive urothelial carcinoma
    - Clinical features
    - Complications

Classification of bladder tumours

EPITHELIAL TUMOURS

a) Urothelial (transitional cell) tumours (90%)

1. Transitional cell papilloma (exophytic, Inverted)
2. Papillary urothelial neoplasms of low malignant potential (PUNLMP)
3. Carcinoma in situ (CIS, or fl at non-invasive urothelial carcinoma)
4. Transitional cell carcinoma (low grade, high grade)

b) Squamous cell carcinoma (5%)
c) Adenocarcinoma (Rare)
d) Small cell carcinoma (Rare)
e) Mixed carcinoma (5%)

NON-EPITHELIAL TUMOURS (Rare)

- Benign (leiomyoma, granular cell myoblastoma, neurilemmoma, haemangioma)
- Malignant (Sarcomas)

Urothelial (Transitional Cell) Bladder Tumours

Introduction →



- 90% of bladder tumours arise from transitional epithelial (urothelium) lining of bladder

Age →

5th decade of life

Gender →

M > F

Etiopathogenesis

f) Smoking
g) Industrial occupations
h) Schistosomiasi
i) Dietary factors
j) Local lesions
k) Drugs
l) Prior irraditation

a) Smoking →

- 2-to-4-fold increased risk of developing bladder cancer

b) Industrial occupations →

- Workers in industries that produce aniline dyes, rubber, plastic, textiles, and cable have high incidence of bladder cancer.
- Carcinogenic substances responsible for bladder cancer → metabolites of β-naphthylamine and benzene.

c) Schistosomiasis →

- Increased risk of squamous cell carcinoma

d) Dietary factors →

- Artificial sweeteners like saccharin, coffee or caffeine and chronic alcoholism

e) Local lesions →

- ectopia vesicae (extrophied bladder), vesical diverticulum, leukoplakia of bladder mucosa

f) Drugs →

- Immunosuppressive therapy with cyclophosphamide and patients having analgesic-abuse → high risk of bladder cancer.

g) Prior irraditation →

- Patients who have received prior irradiation for some other pelvic cancers have higher risk of developing bladder cancer.

Gross

- 90% → Papillary (non-invasive or invasive)
- 10% → flat indurated (non-invasive or invasive)
- Location in bladder → Lateral walls > posterior wall > region of trigone.



Microscopy

- Histologic criteria for classifying urothelial tumours as per WHO/ISUP

FEATURE	Papilloma	PUNLMP	Low-grade TCC	High-grade TCC
Architecture				
1. Papillae	Delicate	Delicate	Fused, branching	Fused, branching
2. Cellular organisation • Resemblance to normal • Thickness • Loss of polarity • Cohesiveness • Crowding	 Yes Normal No Yes No	 Generally, yes Any thickness No Yes No	 Orderly Any thickness Minimal Yes Minimal	 Mainly disordered Any thickness Frequent Often lost Yes
Cellular Features				
1. Nuclear size	Normal	May be uniformly enlarged	Enlarged with variation in size	Enlarged with variation in size
2. Nuclear shape	Normal	Elongated, uniform	Round-to-oval, slight pleomorphism	Moderate to marked pleomorphism
3. Nuclear chromatin	Fine	Fine	Mild variation	Moderate to marked variation
4. Nucleoli	Absent	Rare	Inconspicuous, small, regular	Conspicuous, multiple
5. Mitosis	Absent	Rare, basal	Occasional, any level	Frequent, any level
6. Umbrella cells	Present	Present	Usually, present	May be absent

1) Papilloma



Exophytic papillomas →
- Small
- Delicate papillae
- Each papilla is composed of fibrovascular stromal core covered by normal-looking transitional cells having normal number of layers (upto 6-7) in thickness.
- Individual cells resemble the normal transitional cells and do not vary in size and shape.
- Mitoses are absent
- Basal polarity is retained

Inverted papillomas → an endophytic growth pattern



2) Papillary urothelial neoplasms of low malignant potential (PUNLMP)

Similar to papillomas with additional features →
- Increase in the number of epithelial layers
- Round to oval nuclei
- Diffuse nuclear enlargement

Papillary urothelial (Transitional cell) carcinoma

Histologic criteria →
- Architecture → type of papillae and relationship of cell layers with basement membrane as regards polarity.
- Cytologic criteria → extent of resemblance with normal cells, crowding, variation in nuclear size, shape, chromatin, nucleoli (their presence and type), and mitotic figures.
- Criteria for invasion → penetration of basement membrane of bladder mucosa and presence or absence of invasion by neoplastic cells in muscle.

2 grades→

a) Low grade - TCC
b) High grade - TCC



a) Low grade - TCC

- Tumours show fused and branching papillary pattern
- But overall, there is an orderly arrangement of layers of cells.
- Tumour cells are cohesive
- Show mild variation in polarity, nuclear size, chromatin and shape and inconspicuous small and regular nucleoli.
- Mitotic figures are infrequent

b) High grade - TCC

- Increased thickness
- Fused and branching papillae
- Disorderly arrangement.
- Tumour cells show nuclear enlargement, moderate to marked variation in nuclear size, shape, hyperchromatism, and multiple prominent nucleoli.
- Mitoses are frequent and atypical





3) Invasive urothelial carcinoma

- Any grade of papillary urothelial carcinoma → invasion into lamina propria or further into muscularis propria (detrusor).

Clinical features

Painless hematuria:
- Only clinical feature.
- Sometimes associated with frequency, urgency and dysuria.

Complications

When the tumor obstructs the ureteral orifice→
- Pyelonephritis
- Hydronephrosis

(19.01) CRYPTORCHIDISM

University Exams

Short Questions

1. Define Cryptorchidism. Write its Etiology, Gross and Microscopy
2. Describe Etiology, Clinical features, Complications and Treatment of Cryptorchidism.

Overview

- Normal testis →
  - Anatomy
  - Histology
  - Development of testis
- Cryptorchidism
  - Introduction
  - Etiology
  - Gross
  - Microscopy
  - Clinical features
  - Treatment

Normal testis → Anatomy

- Contents of scrotal sac → testicle, epididymis, lower end of spermatic cord
- Testies Covered by thick inseparable covering of fibrous tissue → tunica albuginea
- Serous space in front and lateral surface of testis → tunica vaginalis





Normal testis → Histology



NORMAL TESTIS

Seminiferous tubules →

Cells lining seminiferous tubules are of 2 types→

1. Spermatogonia or germ cells → produce spermatozoa.
2. Sertoli cells → supportive cells to germ cells → produce testosterone (mainly) and oestrogen (little)

Germ cells	Sperm production
Sertoli cells	Testosterone and oestrogen production
Leydig cells	Testosterone production

Fibrovascular stroma between seminiferous tubules →

Interstitial cells of Leydig →

- Abundant cytoplasm
- Lipid granules
- Reinke's crystals.
- Main source of testosterone

Development of testis

- Testis develops in retroperitoneum close to posterior abdominal wall from genital tubercles
- Guided to scrotum by **gubernaculum**



- **7th month** → reaches deep inguinal ring
- **8th month** → inguinal canal
- **9th month** → superficial inguinal pouch
- **Full term** → reaches scrotum

Cryptorchidism

Introduction →

- Complete or partial failure of the intra-abdominal testes to descend into the scrotal sac
- Incidence → **0.2%** in adult male population.
- Association → testicular dysfunction and increased risk of testicular cancer
- Unilateral → in **80%** cases
- Bilateral → in **20%** cases



Etiology

1. **Mechanical factors** → e.g. short spermatic cord, narrow inguinal canal, adhesions to peritoneum.
2. **Genetic factors** → e.g. trisomy 13, maldevelopment of scrotum or cremaster muscles.
3. **Hormonal factors** → e.g. deficient androgenic secretions.

In Cryptorchidism →

- 70% of cases → undescended testis lies in inguinal ring
- 25% of cases → undescended testis lies in abdomen
- 5% of cases → undescended testis lies at other sites along its descent from intra-abdominal location to the scrotal sac



Gross

Small in size, firm and fibrotic.

Microscopy

1. Seminiferous tubules:
   - Progressive loss of germ cell elements
   - Tubular basement membrane is thickened.
   - Hyalinized tubules
2. Interstitial stroma:
   - Increase in interstitial fibrovascular stroma
   - Leydig cells are prominent



Clinical Features

- Asymptomatic and discovered only on physical examination.
- But if surgical correction by orchiopexy is not undertaken by about 2 years of age →

    1. Sterility-infertility
    2. Inguinal hernia
    3. Malignancy → most commonly seminoma→ Cryptorchid testis is at 30-50 times increased risk of developing testicular malignancy than a normally descended testis.

Treatment

Orchiopexy → placement in scrotal sac

Age of surgery→

- During first year of life → majority of inguinal cryptorchid testes descend spontaneously into scrotum.
- Those that remain undescended require surgical correction before 2 years of age

(19.02) TESTICULAR TUMOURS

University Exams

Long Questions

1. Write Pathological classification of testicular tumors. Describe etiology, morphology and clinical features of most common testicular tumor.
2. Write Clinical classification of testicular tumors. Describe etiology, morphology and clinical features of Seminoma.
3. Classify testicular tumors. Describe etiology, morphology and clinical features of Germ cell tumours of testis

Short Questions

1. Name germ cell tumors of testis and describe any one of them.
2. Pathological and Clinical classification of testicular tumors.
3. Etiology and Spread of testicular neoplasms.
4. Gross and Microscopic appearance of seminoma.
5. Gross and Microscopic appearance of teratoma of testis
6. Serological markers (Tumour markers) in testicular tumors.

Very Short Questions

1. Draw a well labelled diagram of seminoma.
2. Modes of spread of classical seminoma.
3. Non-seminomatous germ cell tumors of testis.

Overview

- Testicular Tumours Classification →
    1. Pathological classification
    2. Clinical classification
- Etiologic Factors
- Clinical Features
- Spread
- a) Seminoma
  b) Yolk sac tumour
  c) Teratoma
  } Introduction, Gross, Microscopy, Tumour markers

Testicular Tumours Classification

1) Pathological classification

Germ Cell Tumours → (YESPCT)

1. Yolk sac tumour (Endodermal sinus tumour)
2. Embryonal carcinoma
3. Seminoma
4. Polyembryoma
5. Choriocarcinoma
6. Teratomas

Sex Cord-stromal Tumours → (LSG)

1. Leydig cell tumour
2. Sertoli cell tumour (Androblastoma)
3. Granulosa cell tumour

Combined Germ Cell-sex Cord-stromal Tumours

1. Gonadoblastoma

2) Clinical classification



FEATURE	SGCT	NSGCT
Primary tumour	Larger, confined to testis; testicular contour retained	Smaller, testicular contour may be distorted
Metastasis	Regional lymph nodes	Haematogenous spread early
Response to radiation	Radiosensitive	Radioresistant
Serum markers	OCT -4, KIT, PLAP → high hCG, AFP → generally low levels	hCG, AFP, or both→ high levels
Prognosis	Better	Poor

Etiologic Factors

1. Cryptorchidism
2. Dysgenetic gonads
3. Genetic factors
4. Orchitis
5. Trauma
6. Carcinogens

Clinical Features

- Gradual gonadal enlargement
- Dragging sensation in testis
- Metastatic involvement → secondary symptoms such as pain, lymphadenopathy, haemoptysis and urinary obstruction.

Spread

1. Lymphatic spread → retroperitoneal para-aortic lymph nodes, mediastinal lymph nodes and supraclavicular lymph nodes.
2. Haematogenous spread → lungs, liver, brain and bones.

a) Classic Seminoma

Introduction →

- Commonest malignant tumour of testis
- constitutes about 45% of all GCTs
- Corresponds to dysgerminoma in female

Gross

- Tumour replaces entire testis
- Cut section → homogeneous, grey-white lobulated appearance

Microscopy

1. Tumour cells arrangement → lobules.
2. Tumour cells cytology→
    - Uniform in size
    - Clear cytoplasm
    - Well-defined cell borders.
    - Cytoplasm contains variable amount of glycogen that stains positively with PAS reaction.
    - Nuclei are centrally located, large, hyperchromatic and usually contain 1-2 prominent nucleoli.
3. Stroma → characteristic lymphocytic infiltration



Tumour markers

1. KIT
2. OCT4
3. Placental alkaline phosphatase (PLAP)

- Negative for alpha-fetoprotein (AFP) and human chorionic gonadotropin (HCG)

Prognosis
- Highly radiosensitive
- Prognosis of classic seminoma is better than other germ cell tumours.

Spermatocytic seminoma
- Found in old age
- Do not metastasize
- Have excellent prognosis

b) Yolk Sac Tumour

Introduction →

- Also known as Endodermal Sinus Tumour or Orchioblastoma
- Most common testicular tumour of infants and young children (up to age of 4 years)

Gross

- Soft, yellow-white, mucoid with areas of necrosis and haemorrhages

Microscopically

- Tumour cells arranged in **reticular network**
- Tumour cells → **flattened to cuboid** epithelial cells with clear vacuolated cytoplasm.
- **Schiller-Duval bodies** → Tumour cells form papillary projections having a central blood vessel with perivascular layer of anaplastic embryonal germ cells.
- **Hyaline globules** → intracellular and extracellular, PAS-positive



Schiller Duval body

Tumour marker

1. α-fetoprotein (AFP) (100% cases)
2. α1- antitrypsin

c) Teratoma

Introduction →

- Composed of Totipotential cells
- All 3 germ cell layers → endoderm, mesoderm and ectoderm
- More common in infants and children



Gross

- Cut surface → variegated appearance
- Grey-white solid areas, cystic and honey-combed areas, and foci of cartilage and bone, sometimes hairs

Microscopy

Teratoma is composed of disorderly **mixture** of a variety of well differentiated structures such as

1. Cartilage
2. smooth muscle
3. Intestinal and respiratory epithelium
4. Mucus glands
5. Cysts lined by squamous and transitional epithelium



6. Neural tissue
7. Fat
8. Bone

Tumour marker

- hCG or AFP levels or both (**in 50% cases)**

	Seminoma	Yolk sac tumour	Teratoma
Introduction			
Gross			
Microscopy			
Tumour markers			

(19.03) BENIGN PROSTATE HYPERPLASIA & PROSTATE CARCINOMA

University Exams

Long Questions

1. Prostate in benign prostatic hyperplasia.
2. Describe etiology, Gross and microscopy of benign prostatic hyperplasia.
3. Describe etiology, Gross, microscopy, spread and staging of Prostate Carcinoma

Short Questions

1. Pathogenesis of nodular hyperplasia of prostate.
2. Morphology of nodular hyperplasia of prostate.
3. Spread of carcinoma prostate.
4. Staging of prostatic cancer.
5. Prostate-specific antigen.

Very Short Questions

1. Tumour markers in prostatic cancer
2. Glesons score in prostatic cancer

Overview

Normal Prostate →

- Anatomy
- Histology

BPH

- Introduction
- Site
- Etiology
- Gross
- Microscopy

Prostate Carcinoma

- Introduction
- Site
- Gross
- Microscopy
- Spread
- Tumour markers
- Grading → Glesons score

Normal Prostate → Anatomy

- Weighs approx **20 gm.**
- Surrounds commencement **of male urethra**





5 lobes →
- Anterior
- Middle
- Posterior
- 2 lateral lobes

3 zones →
- Central
- Peripheral
-
- Transitional



LOBAR DISTRIBUTION OF PROSTATE GLAND



ZONAL DISTRIBUTION OF PROSTATE GLAND

Remember

- BPH → arises from transitional zone (Middle lobe)
- Carcinomas → arise from peripheral zone (Posterior lobe)



Normal Prostate → Histology

Glandular epithelium consists of 2 layers →
1. A basal layer of low cuboidal cells
2. An inner layer of mucussecreting tall columnar cells.

Fibromuscular Stroma



Benign prostate hyperplasia (BPH)

Introduction →

- More frequent above age of **50 years**
- Incidence → 80% in men above 80 years.

Site →

Arises from **transitional zone (Middle lobe)**

Etiology

Enzyme type 2 5α-reductase located in stromal
cells (not in epithelial cells)
↓
Converts testosterone → DHT
↓
DTH moves to epithelial cells
↓
Binds to nuclear androgen receptor (AR) present in both
stromal and epithelial prostate cells
↓
Stimulates transcription of androgen-dependent genes
↓
Growth factors and their receptors are formed
↓
Prostatic hyperplasia
↓
BPH



Gross

- Enlarged prostate → nodular, smooth and firm
- Cut surface → firm, homogeneous





Microscopy

Glandular hyperplasia→

- Exaggerated intra-acinar papillary infoldings with delicate fibrovascular cores.
- Lining epithelium → two-layered→ inner tall columnar mucus-secreting with poorly-defined borders and outer cuboidal epithelium with basal nuclei.

Fibromuscular stroma hyperplasia→ aggregates of **spindle cells**

Prostate Carcinoma

Introduction →

- Second most common cancer in males (Most common is lung cancer)
- Above age of 50 years
- Incidence → More than 50% of men 80 years old have asymptomatic (latent) carcinoma of the prostate.

Site →

Arises from arise from peripheral zone (Posterior lobe)

Gross

- Enlarged or normal in size
- Cut section → homogeneous and contains irregular yellowish areas.





Microscopy

- Loss of intra-acinar papillary convolutions.
- Acini are closely packed in back-to-back arrangement without intervening stroma
- Glands are lined by a single layer of cuboidal cells.
- Malignant acini have little or no stroma between them.
- Perineural Invasion

Spread

- Direct spread → extend into bladder neck, seminal vesicles, trigone and ureteral openings.
- Lymphatic spread → spread to sacral, iliac and para-aortic lymph nodes
- Haematogenous spread → osteoblastic osseous metastases, especially to pelvis, and lumbar spine
- Other sites of metastases are lungs, kidneys, breast and brain.

Tumour markers

1. Prostatic acid phosphatase (PAP)
2. Prostate-specific antigen (PSA)

Grading → Glesons score

- Tumors contain more than one pattern →

    1. **Primary grade →** dominant pattern
    2. **Secondary grade →** sub-dominant pattern.

- Both primary and secondary architectural patterns are graded from 1 to 5, with 1 being most differentiated and 5 being undifferentiated.
- Gleason score is derived by **addition of these two grades**



SCORES 2-5	SCORES 6-7	SCORES 8-10
Low-grade Prostate Cancer	**Intermediate-grade** Prostate Cancer. Most prostate cancers fall into this group.	**High-grade** Prostate Cancer

(19.04) MALE INFERTILITY

University Exams

Short Questions

1. What are the causes of male infertility?
2. Pre-testicular and testicular causes of male infertility?

Overview

- Introduction
- Causes →

    1. Pre-testicular causes
    2. Testicular causes
    3. Post-testicular causes

Introduction

Causes

1. Pre-testicular
2. Testicular
3. Post-testicular



1) Pre-Testicular Causes

1. Hypopituitarism → from tumour, trauma, infarction, cyst and genetic deficiency of FSH and LH secretion.
2. Oestrogen excess → Endogenous excess such as from hepatic cirrhosis, adrenal tumour, Sertoli and Leydig cell tumour; or exogenous excess such as in treatment of carcinoma of the prostate.
3. Glucocorticoid excess → Endogenous excess in Cushing's syndrome, Exogenous excess in treatment of ulcerative colitis, bronchial asthma, rheumatoid arthritis etc.
4. Other endocrine disorders ➔ Hypothyroidism and diabetes mellitus → associated with hypospermatogenesis.



Pretesticular azoospermia

2) Testicular Causes


Testicular azoospermia

1. Agonadism → total absence of testes.
2. Cryptorchidism → undescended testis
3. Maturation arrest → failure of spermatogenesis beyond one of the immature stages.
4. Hypospermatogenesis → presence of all maturation stages of spermatogenesis but in decreased number.
5. Sertoli cell-only syndrome → congenital or acquired absence of all germ cells so that seminiferous tubules are lined by Sertoli cells only.
6. Klinefelter's syndrome → XXY intersexuality characterised by primary hypogonadism, azoospermia, gynaecomastia, eunuchoid built and subnormal intelligence.
7. Mumps orchitis → occurring as a complication
8. Irradiation damage → resulting in permanent germ cell destruction.

3) Post-Testicular Causes


Post-testicular or obstructive azoospermia

1. Congenital block → absence or atresia of vas deferens.
2. Acquired block → due to gonorrhoea and surgical intervention.
3. Impaired sperm motility → e.g. immotile cilia syndrome

(20.01) CERVICAL PRECANCERS (CIN / SIL)

University Exams

Long Questions

1. Describe in brief etiopathogenesis, classification and morphology of Cervical Dysplasia and Cervical Intraepithelial Neoplasia.

Short Questions

1. Define Cervical Precancers. write its Etiology
2. Classification of Cervical Precancers
3. Cervical intraepithelial neoplasia (CIN).
4. Classification of cervical intraepithelial neoplasia.
5. Cervical intraepithelial neoplasia grade I, II, III (CIN I, CIN II, CIN III).
6. Well labelled diagram of cervical dysplasia and carcinoma in situ.
7. Write precursor of Ca cervix.
8. Bethesda system for Classification of precursor of Ca cervix.
9. Squamous Intraepithelial Lesion → L- SIL, H- SIL

Overview

Cervix →
- Anatomy
- Histology

Cervical Precancers →
- Definition
- Age
- Etiology
- Classification of Cervical Precancers →

    1. Carcinoma In Situ (CIS) System → Mild, Moderate, Severe dysplasia
    2. Cervical Intraepithelial Neoplasia (CIN)Classification → CIN – 1, CIN – 2, CIN - 3
    3. Bethesda system (presently used) → L- SIL, H- SIL

Cervix →

Anatomy

- **Internal os** → communicating with endometrial cavity above
- **External os** → opening into vagina below.



Histology

- Ectocervix → stratified squamous epithelium
- Endocervix → columnar mucus-secreting epithelium.



Cervical Precancers

Definition →

Spectrum of intraepithelial changes → begins from minimal atypia and progresses through more marked stages of intraepithelial abnormalities to invasive squamous cell carcinoma.

Age →

< 40 years

Etiology

1. Human Pappiloma virus (HPV)
2. Environmental Factors

1) Human Pappiloma virus (HPV)

- Most common risk factor associated with CIN and cancer cervix
- Sexually transmitted DNA virus.
- Risk features associated with increased exposure to HPV →

  a) Multiple sexual partners
  b) First intercourse at young age
  c) High parity
  d) A male partner having multiple previous or current sexual partners
  e) Inefficient immune response

Classification of HPV

Low risk HPV

- **Subtypes 6 and 11**
- cause genital warts (condyloma acuminatum)

High risk HPV

- **Subtypes 16,18,31,33,45,52,56,58,59,68**
- cause squamous cell carcinoma of vagina, vulva, penis, anus

Oncogenesis by HPV



Environmental Factors

1. Presence of co-carcinogens: Cigarette smoke (contains polycyclic hydrocarbons → acts as a co-carcinogen)
2. Coexisting microbial infections.
3. Dietary deficiencies.
4. Hormonal changes.
5. Use of oral contraceptives.

Classification of Cervical Precancers

1. Carcinoma In Situ (CIS) System → Mild dysplasia, Moderate dysplasia, Severe dysplasia
2. Cervical Intraepithelial Neoplasia Classification → CIN – 1, CIN – 2, CIN - 3
3. Bethesda system (presently used) → Squamous Intraepithelial Lesion → L- SIL, H- SIL



1) Carcinoma In Situ (CIS) System

- Based on dysplasia
- Dysplasia → alteration in cell morphology and disorderly arrangement of the cells of stratified squamous epithelium
- Premalignant lesion

Dysplastic cells show following features→

1. Loss of polarity
2. Increased nuclear cytoplasmic size ratio (normal 1:4 to 1:6)
3. Increased number of mitosis with normal (typical) pattern
4. Hyperchromatasia
5. Pleomorphism

- Basement membrane is intact and there is no stromal infiltration.
- Depending upon thickness of squamous epithelium involved by atypical cells→

    1. Mild dysplasia → < 1/3rd involvement of thickness of epithelium
    2. Moderate dysplasia → 1/3rd to 2/3rd involvement
    3. Severe dysplasia → > 2/3rd involvement

- Carcinoma in situ → When dysplastic changes involve entire thickness of epithelium but basement membrane is intact
- Invasive carcinoma → When cancer cells breach basement membrane

2) Cervical Intraepithelial Neoplasia (CIN) Classification

1. CIN-1 (Mild dysplasia) → < 1/3rd involvement of thickness of epithelium
2. CIN-2 (Moderate dysplasia) → 1/3rd to 2/3rd involvement
3. CIN-3 (Severe dysplasia) → > 2/3rd involvement

Rate of progression of C1N			
	CIN I	**CIN II**	**CIN III**
Regression to Normal	60%	40%	30%
Persistence	30%	35%	50%
Progression to	10%	20%	
Progression to cancer	< 1%	5%	20%

3) Bethesda system

L-SIL →

- Corresponds to CIN-1 (includes mild dysplasia)
- Related to HPV 6 and 11 infection
- About 10% cases of L-SIL may progress to H-SIL.

H-SIL →

- Corresponds to CIN-2 and 3 (Includes moderate dysplasia, severe dysplasia, and carcinoma in situ)
- Related to HPV 16 and 18 infection
- Approximately, 10% cases of H-SIL may progress to invasive cervical cancer over a period of 2 years.

WHO	CIN	Description	Bethesda classification
Mild dysplasia	CIN I	Dysplastic cells seen in lower 1/3 of epithelial lining of cervix	LSIL
Moderate dysplasia	CIN II	Dysplastic cells seen in 2/3 of epithelial lining of cervix	HSIL
Severe dysplasia	CIN III	Dysplastic cells seen in more than 2/3 of epithelial lining of cervix	HSIL
	Carcinoma in situ	Dyplastic cells seen full thickness but basement membrane is intact	HSIL





(20.02) CERVIX CANCER

University Exams

Long Questions

1. Describe the etiology, pathogenesis, clinical features and spread of carcinoma cervix.
2. Describe the Predisposing factors, Precursor, Classification, Gross and Microscopy of carcinoma cervix.
3. A 48-year-old lady presented with bleeding per vagina. Per speculum examination showed an ulcer proliferative growth involving the external os which bleeds on touch.

    a) What Is the most probable diagnosis and why?
    b) What is the etiopathogenesis of this condition?
    c) Add a note on screening method for this condition.

Short Questions

1. Carcinoma cervix → etiology
2. Role of virus in carcinoma of cervix.
3. Enumerate etiologic factors of carcinoma of cervix. Describe gross and microscopic features of carcinoma of cervix.
4. Write precursor lesion, gross and microscopy of Ca cervix.
5. FIGO Staging of Ca cervix.

Very Short Questions

1. Risk factors for carcinoma cervix.
2. Screening of carcinoma in cervix.
3. Morphological features seen in carcinoma of cervix.
4. Pap smear

Overview

- Introduction
- Age
- Etiology / Predisposing factors
- Precursor
- Classification
- Gross
- Microscopy
- Clinical features
- FIGO Staging
- Screening → Pap smear
- Spread

Introduction →

Most common gynaecological malignancy in the world

Age →

Bimodal peak →
- 1st peak → 35-39 years
- 2nd peak → 55-60 years

Etiology

5. Human Pappiloma virus (HPV)
6. Environmental Factors →

    a) Presence of co-carcinogens: Cigarette smoke (contains polycyclic hydrocarbons → acts as a co-carcinogen)
    b) Coexisting microbial infections
    c) Dietary deficiencies
    d) Hormonal changes
    e) Use of ral contraceptives

Precursor

- Immediate Precursor → H - SIL



Classification

1. Squamous cell carcinoma (Epidermoid)
2. Adenocarcinoma



Gross

3 types of patterns→

1. Fungating
2. Ulcerating
3. Infiltrating

Microscopy

Squamous cell carcinoma (Epidermoid)

- More common (70%)
- Arises from → squamocolumnar junction.
- Tumour cells →
    - Arranged in sheets
    - Large with hyperchromatic nuclei and high N/C ratio
    - Non keratinising

Adenocarcinoma

- Less common (20-25%)
- Arises from → endocervix
- Tumor cells →
    - Arranged in glandular pattern
    - Form glands
    - Produce mucin

Clinical features

- Vaginal bleeding after sexual intercourse

FIGO Staging

- Cancer Committee of the International Federation of Gynaecology and Obstetrics (**FIGO classification**)

Stage I: The carcinoma Is strictly confined to the cervix uteri (extension to the corpus should be disregarded)
IA : Invasive carcinoma that can be diagnosed only by microscopy, with maximum depth of invasion ≤5 mm
IA1 Measured stromal invasion <3 mm in depth
IA2 Measured stromal invasion ≥3 mm and ≤5 mm in Depth
IB : Invasive carcinoma with measured deepest invasion ≤5 mm (greater than stage IA), lesion limited to the cervix uteri
IB1 Invasive carcinoma ≥5 mm depth of stromal invasion and <2 cm in greatest dimension
IB2 Invasive carcinoma ≥2 cm and <4 cm in greatest Dimension
IB3 Invasive carcinoma ≥4 cm in greatest dimension
Stage II: The carcinoma Invades beyond the uterus, but has not extended onto the lower third of the vagina or to the pelvic wall
IIA : Involvement limited to the upper two-thirds of the vagina without parametria! involvement
IIA1 Invasive carcinoma <4 cm in greatest dimension
IIA2 Invasive carcinoma ≥4 cm in greatest dimension
IIB : With parametria! involvement but not up to the pelvic wall
Stage III: The carcinoma Involves the lower third of the vagina and/or extends to the pelvic wall and/or causes hydronephrosis or non-functioning kidney and/or Involves pelvic and/or paraaortic lymph node dissection.
IMA : Carcinoma involves the lower third of the vagina, with no extension to the pelvic wall
1MB : Extension to the pelvic wall and/or hydronephrosis or non-functioning kidney (unless known to be due to another cause)
IMC Involvement of pelvic and/or paraaortic lymph nodes, irrespective of tumor size and extent (with r and p notations)
IIIC1 Pelvic lymph node metastasis only
IIIC2 Paraaorticlymph node metastasis
Stage IV: The carcinoma has extended beyond the true pelvis or has Involved (biopsy proven) the mucosa of the bladder or rectum. A bullous edema, as such, does not permit a case to be allotted to stage IV
IVA Spread of the growth to adjacent organs
IVB Spread to distant organs



Screening → Pap Smear

- Exfoliative Cytology
- Obtaining sample of Exfoliative cells of cervix by brushing or scraping



Procedure

Undress from waist down
↓
Lie down on your back on an exam table with knees bent (lithotomy position)
↓
Insert speculum into vagina
↓
Take samples of cervical cells using a soft brush and spatula

Pap smear:
cells are scraped from the cervix and examined under a microscope to check for disease or other problems





Cervix viewed through speculum with patient in lithotomy position

Spread

Direct extension → Tumor spreads directly to adjacent organs → uterus, vagina

Lymphatic spread →

- H → Hypogastric
- O → Obturator
- P → Presacral and Parametrial
- E → External iliac

Hematogenous spread → M/ C site → lungs

(20.03) ENDOMETRIOSIS

University Exams

Short Questions

1. Define Endometriosis. Write gross, microscopic and complications of Endometriosis.
2. Pathogenesis and Molecular changes of Endometriosis.
3. Define Endometriosis. Write its Sites and Pathogenesis
4. Pathogenesis and Clinical features of Chocolate cyst (Endometriomas)
5. Morphology of endometriosis of ovary.

Overview

- Definition
- Age
- Sites
- Pathogenesis
- Molecular changes
- Gross
- Microscopy
- Chocolate cyst (Endometriosis of ovary)

Definition →

Presence of endometrial tissue outside uterus

Age →

Reproductive life (3rd and 4th decades)

Sites →

- Most common site → **Ovary**

Other sites →

- Uterine ligaments
- Rectovaginal septum
- Cul-de sac
- Pelvic peritoneum
- Gastrointestinal tract
- Mucosa of cervix
- Vagina
- Fallopian tube
- Laparotomy scars





Pathogenesis

1. Retrograde menstruation theory
2. Benign metastases theory
3. Metaplastic theory

1) Retrograde menstruation →

Most widely accepted theory

Retrograde menstruation through fallopian tubes
↓
Diffuse seeding of endometrial tissue
↓
Endometrial tissue deposited outside uterus
↓
Endometriosis



2) Benign metastases theory →

Endometrial tissue spread from uterus to distant sites (e.g. bone, lung, and brain) via blood vessels and lymphatic channels.

3) Metaplastic theory →

Endometrium arises directly from metaplasia of celomic epithelium (mesothelium of pelvis or abdomen).

Molecular changes

- Endometriotic tissue differs from normal endometrium by exhibiting→
    1. Marked activation of inflammation → prostaglandin production
    2. Increased stromal aromatase activity → estrogen productiong



Gross

Nodules with a red-blue to yellow-brown appearance

Microscopy

- Both endometrial glands and stroma with or without presence of hemosiderin-laden macrophages at ectopic site



Chocolate cyst (Endometriomas)

Endometriotic lesions bleed periodically
↓
Convert ovaries into large cystic masses
↓
Filled with brown fluid (resulting from previous hemorrhages)
↓
Chocolate cyst (Endometriomas)

Age → Reproductive years

Clinical features → Triad →
1. Dysmenorrhea
2. Dyspareurnia
3. Infertility

Risk → Endometriosis may be a precursor to carcinoma (endometrioid and clear cell carcinoma).

(20.04) ADENOMYOSIS

University Exams

Short Questions

1. Adenomyosis uterus.
2. Define Adenomyosis. Write its Gross and Microscopy

Overview

- Definition
- Clinical features
- Gross
- Microscopy

Definition →

- Presence of endometrial tissue into and between the smooth muscle fascicles of underlying myometrium.
- Coexist with endometriosis



Clinical Features →

- No cyclic bleeding
- Metrorrhagia (irregular and heavy menses)
- Colicky dysmenorrhea
- Dyspareunia
- Pelvic pain (during premenstrual period)



Gross →

- Uterus → mild to moderate enlargement.
- Cut section → coarse trabeculations with ill-defined area of hemorrhage

Microscopy →

- Irregular nests of benign endometrial glands and stroma deep within the myometrium.
- Minimum distance between basal endometrium and the endometrial tissue of adenomyosis → at least 2–3 mm



(20.05) ENDOMETRIAL HYPERPLASIA

University Exams

Short Questions

1. Classification of Endometrial hyperplasia.
2. Morphology of various types of endometrial hyperplasia.
3. Pathogenesis and Clinical significance of endometrial hyperplasia.
4. Risk of malignant transformation in endometrial hyperplasia.
5. Causes of endometrial hyperplasia.
6. Morphology of cystoglandular hyperplasia.
7. Morphology of Complex atypical hyperplasia

Overview

- Introduction
- Pathogenesis
- Clinical significance
- Classification →

    1. Simple hyperplasia without atypia
    2. Complex hyperplasia without atypia
    3. Complex hyperplasia with atypia

- Risk of malignant transformation

Introduction

Increased proliferation of endometrial glands relative to stroma
↓
Increased gland-to-stroma ratio when compared with normal proliferative endometrium
↓
increased thickness of endometrium
↓
Abnormal uterine bleeding



Pathogenesis

Prolonged oestrogenic stimulation unopposed with any progestational activity →

Endogenous sources→
- Polycystic ovarian disease, e.g. Stein-Leventhal syndrome.
- Functioning granulosa -theca cell tumours
- Adrenocortical hyper function

Exogenous sources→ Prolonged administration of oestrogen.

Clinical significance

- Prolonged, profuse
- Irregular uterine bleeding in a menopausal or postmenopausal woman.
- Presence of cellular atypia which is closely linked to endometrial carcinoma.
-

Classification

1. Simple hyperplasia without atypia (Cystic glandular hyperplasia).
2. Complex hyperplasia without atypia (Complex nonatypical hyperplasia).
3. Complex hyperplasia with atypia (Complex atypical hyperplasia).



1) Simple Hyperplasia Without Atypia (Cystic Glandular Hyperplasia)

- Presence of varying sized glands → large and cystically dilated
- Lined by atrophic epithelium.
- Mitoses are scanty
- No atypia
- Stroma between glands → sparsely cellular and oedematous



2) Complex Hyperplasia Without Atypia (Complex Non-atypical Hyperplasia)

- Glands are increased in number
- Variation in size and are irregular in shape.
- Lined by multiple layers of tall columnar epithelial cells with large nuclei
- Basal polarity maintained
- No significant atypia.
- Glandular epithelium at places is thrown into papillary infolds into adjacent stroma i.e. there is crowding and complexity of glands without cellular atypia.
- Stroma → dense, cellular and compact



3) Complex Hyperplasia with Atypia (Complex Atypical Hyperplasia)

Complex hyperplasia with atypia is distinguished from complex non-atypical hyperplasia by →

- Presence of atypical cells
- Loss of polarity
- Large size
- Irregular and hyperchromatic nuclei
- Prominent nucleoli
- Altered nucleocytoplasmic ratio

Risk of malignant transformation

- Cystic hyperplasia → Minimal risk (<1%)
- Complex hyperplasia without atypia → 3%
- Complex hyperplasia with atypia → 20-25%

(20.06) ENDOMETRIAL CARCINOMA

University Exams

Long Questions

1. Describe the etiology, pathogenesis, clinical features and spread of endometrial carcinoma.
2. Describe the Predisposing factors, Classification, Gross and Microscopy of endometrial carcinoma.

Short Questions

1. Risk factors for endometrial carcinoma
2. Gross and microscopy of endometrial carcinoma.

Overview

- Introduction
- Age
- Classification
- Etiology / Predisposing factors
- Pathogenesis
- Gross
- Microscopy
- Clinical features
- Spread

Introduction →

- Most common pelvic invasive cancer of female genital tract.
- Also called uterine cancer

Age →

Postmenopausal women (6th to 7th decades)

Classification

1. Type I
2. Type II

Etiology

Type I

- Chronic unopposed oestrogen excess
- Obesity
- Diabetes mellitus
- Hypertension
- Nulliparous state
- Heredity

Type II

- Thin physique
- Atrophy of endometrium

Pathogenesis

Type I

- Mutations or inactivation of PTEN
- Mutations in PIK3CA, PIK3R1, ARID1A, KRAS, and TP53.

Type II

- Mutations in TP53, PIK3CA, FBXW7, PPP2R1A.
- Germline BRCA1/2 mutations

Gross

2 patterns →

1. Localised polypoid tumour
2. Diffuse tumour



Microscopy

Depending upon the pattern of glands and individual cell changes →
a) Well-differentiated
b) Moderately-differentiated
c) Poorly-differentiated

a) Well differentiated →

Increase in number of glands→ 'back-to-back crowding' due to obliterated intervening stroma.

b) Moderately differentiated →

Shows all above features along with presence of some solid sheets (< 50 %)of malignant cells.

c) Poorly differentiated →

Presence of solid sheets (> 50 %) of malignant cells which show marked cytologic atypia and frequent mitoses.

- Glandular pattern is hard to find.



Well Differentiated | Moderately Differentiated | Poorly Differentiated

Clinical features

Abnormal bleeding in postmenopausal woman or excessive flow in premenopausal years.

Spread

- Direct invasion → invade underlying myometrium and extend into periuterine structures.
- Lymphatic spread → To the regional lymph nodes.
- Hematogenous spread → To lungs, liver, bones, and other organs.

Characteristics	Type I	Type II
Age	55-65 years	65-75 years
Predisposing conditions	Unopposed estrogen stimulation, obesity, hypertension, infertility, diabetes	Atrophy of endometrium Thin physique
Precursor lesion	Atypical endometrial hyperplasia	Serous endometrial intraepithelial carcinoma
Molecular features	Mutations or inactivation of *PTEN*, mutations in *PIK3CA, PIK3R1, ARID 1 A, KRAS, and TP53.* Microsatellite instability	Mutations in *TP53, PIK3CA, FBXW7, PP2R1A.* Germline *BRCA1/2* mutations
Microscopy	Endometrioid adenocarcinoma ranges from well to poorly differentiated	• Serous • Clear cell • Mixed Mullerian tumor
Spread	Via lymphatics	Intraperitoneal and via lymphatic
Behaviour	Indolent	Aggressive

(20.07) TUMOURS OF MYOMETRIUM → LEIOMYOMA

University Exams

Short Questions

1. Leiomyoma→ sites, Etiology, types, gross and microscopy.
2. Leiomyoma-uterus (morphology).

Very Short Questions

1. Types of leiomyoma.
2. Microscopy of leiomyoma.

Overview

- Introduction
- Age
- Etiology
- Types
- Gross
- Microscopy
- Clinical features

Introduction →

- Most common uterine tumours of smooth muscle origin
- Also known as fibroids
- Benign

Age →

Rare before age 20 and most regress after menopause.

Etiology

Stimulus to their proliferation → Oestrogen

Evidences →

- Increase in their size in pregnancy and high dose oestrogen-therapy
- Regression following menopause and castration

Types

1. Intramural or interstitial → within myometrium
2. Subserosal → Beneath serosa
3. Submucosal → Underneathe endometrium



Gross

- Circumscribed
- Firm
- Nodular
- Grey-white masses of variable size.
- On cut section → whorled pattern



Microscopy



- Composed of 2 tissue elements → whorled bundles of smooth muscle cells admixed with variable amount of connective tissue.
- Smooth muscle cells are uniform in size and shape with abundant cytoplasm and central oval nuclei
- No mitosis seen

Secondary changes in leiomyomas include:

- Hyaline degeneration
- Cystic degeneration
- Infarction
- Calcification
- Infection and suppuration
- Necrosis
- Fatty change
- Sarcomatous change.

Clinical features

- Usually, asymptomatic.
- Common symptoms →
    - Abnormal bleeding
    - Urinary frequency (compression of the bladder)
    - Infertility
    - Malignant transformation (leiomyosarcoma) → extremely rare

(20.08) OVARIAN TUMOURS

University Exams

Long Questions

1. Classify ovarian tumors. Describe morphological features of surface epithelial tumors.
2. Classify tumors of ovary. Describe risk factors, gross and microscopy of Serous tumors of ovary.
3. Classify tumors of ovary. Describe risk factors, gross and microscopy of mucinous tumors of ovary
4. Classify ovarian tumors. Describe gross and microscopy of germ cell tumors.
5. Classify ovarian tumors. Describe gross and microscopic features of teratoma of ovary.
6. Classify germ cell tumors of ovary and write down gross and microscopic findings of dysgerminoma.

Short Questions

1. Classification of ovarian tumors.
2. Classify ovarian tumors and discuss in brief about mature cystic teratoma.
3. Serous tumors of ovary.
4. Define and classify teratoma. Mention difference teratomas occurring in ovary.
5. Morphology of Serous cystadenoma of ovary.
6. Morphology of mucinous cystadenoma of ovary.
7. Morphology of Dysgerminoma
8. Gross and microscopy of Yolk sac tumour of ovary

Very Short Questions

1. Name four ovarian tumors.
2. Classification of ovarian germ cell tumors.
3. Borderline tumors of ovary.
4. Enlist types of ovarian teratomas.
5. Dermoid cyst → ovary.
6. Struma ovarii.

Overview

- Classification
- Tumours of Surface Epithelium →

  a) Serous Tumors
  b) Mucinous tumours

  } Introduction, Types, Age, Bilateraliy, Risk factors, Gross, Microscopy, Clinical features, Tumour markers

- Germ Cell Tumors →

  a) Dysgerminoma
  b) Yolk sac tumour
  c) Teratoma

  } Introduction, Most common, Age group, Tumour markers, Gross, Microscopy, Prognosis

Classification

Tumours of Surface Epithelium

1. Serous tumour
2. Mucinous tumours
3. Endometrioid tumours

4. Clear cell tumours
5. Brenner tumours

Germ Cell Tumours (YED PCT)

1. Yolk sac tumour (Syn. endodermal sinus tumour, orchioblastoma, infantile type embryonal carcinoma)
2. Embryonal carcinoma
3. Dysgerminoma
4. Polyembryoma
5. Choriocarcinoma
6. Teratomas



Sex Cord-stromal Tumours

- Granulosa-theca cell tumours
- Sertoli-Leydig cell tumours (Androblastoma, arrhenoblastoma)
- Gynandroblastoma

Metastatic Tumours

- Krukenberg tumour

Tumours of Surface Epithelium

a) Serous Tumors
b) Mucinous Tumors

a) Serous Tumors

Introduction →

Most common malignant ovarian tumor (50 % of all cancers of ovary)

Types →

1. Benign
2. Borderline
3. Malignant

Age →

- Benign and borderline tumors → Most common between 20 and 45 years.
- Serous carcinomas → later in life.

Bilateraliy →

- Benign → 20%
- Borderline → 30%
- Malignant → 66%

Risk factors →

1. Family history → High risk
2. Parity: Nulliparous and women with low parity → high risk.
3. Gonadal dysgenesis in children → High risk
4. Genetic factors: Heritable mutations in BRCA1 and BRCA2 → High risk
5. Oral contraceptives and sterilization: reduces risk.

Gross

Benign serous cystadenoma →

- Appearance → Unilocular cysts filled with clear watery serous fluid.
- C/S→ smooth internal surface and may show small papillae



Borderline tumors →

- Appearance → Cystic
- C/S→ cystic cavities filled by increased number of dense and closely packed, cauliflower-like papillary projections

Malignant serous adenocarcinoma →

- Appearance → mixture of solid and cystic areas
- C/S→ solid or papillary, Areas of necrosis and hemorrhage, Adhesion to adjacent structures



Microscopy

Benign →

- Cysts are lined by cuboidal to columnar epithelial cells
- Papillae with a fibrovascular core
- Psammoma bodies.



Borderline →
- Micropapillary carcinoma
- Budding or cellular tufting: tiny, irregular tightly packed clusters of tumor cells which gets detached and float into lumen of cyst.
- Mild nuclear atypia.
- Increased mitotic activity.
- Psammoma bodies.
- Absence of stromal invasion.
- Precursor to low-grade serous carcinoma





Malignant→
- More complex growth patterns
- Solid masses of tumor cells
- Marked nuclear atypia
- Pleomorphism
- Hyperchromatic nuclei
- Atypical mitotic figures
- Psammoma bodies
- Widespread invasion

b) Mucinous Tumors

Introduction →

Less common than serous tumors (30% of all ovarian neoplasms)

Types →

1. Benign
2. Borderline
3. Malignant

Age →

- Middle adult life
- Rare before puberty and after menopause.

Bilateraliy →

Only 5% of mucinous cystadenomas and mucinous cystadenocarcinomas are bilateral.

Risk factors →

1. Smoking
2. Mutations in KRAS (oncogene)

Gross

Benign →
Multilocular, thin-walled cysts with smooth external surface.
Cyst content: Sticky semi-solid mucinous material.



Borderline → Similar to mucinous cystadenoma.

Malignant →
- Similar to mucinous cystadenoma with additional solid regions showing necrosis and hemorrhage.

Microscopy

Benign mucinous cystadenoma →
- Composed of multiple cysts and glands.
- Lined by simple nonstratified tall, non-ciliated, columnar mucinous cells with apical mucin and basally situated nuclei.



Microscopically the mucinous cystadenoma lined by tall columnar cells

Borderline mucinous cystadenoma →
- Stratification of the epithelium → multilayering
- Budding or cellular tufting
- Papillary intraglandular growth
- Mild nuclear atypia.
- Increased mitotic activity.
- Absence of stromal invasion



Mucinous carcinomas→
- Epithelial cell stratification and atypia
- Loss of gland architecture.
- Necrosis.
- Confluent glandular growth
- Expansile invasion.
- Infiltration of the serosa



Clinical features

Benign and Borderline tumors →
- Lower abdominal pain and distention/enlargement
- Vaginal bleeding
- Urinary frequency, dysuria (Due to compression)

Malignant tumors → detected when these spread outside ovary.
- Progressive weakness, weight loss, and cachexia.
- Massive ascites

Tumour Markers

- CA-125 → present in serum of 80% of patients with serous
- Osteopontin → expressed at higher levels in ovarian cancer patients.

	Serous Tumors	Mucinous tumours
Introduction		
Types		
Age		
Bilateraliy		
Risk factors		
Gross		
Microscopy		
Clinical features		
Tumour markers		

Germ Cell Tumors

a) Dysgerminoma
b) Yolk sac tumour
c) Teratoma

a) Dysgerminoma

Introduction →

- Ovarian counterpart of seminoma of testes
- All dysgerminomas are malignant
- Extremely radiosensitive

Most common →

- Malignant germ cell tumour of ovary.
- Ovarian malignancy detected during pregnancy.

Age group →

2nd and 3rd decades.

Tumour markers →

a) Placental alkaline phosphate
b) LDH

- It does not secrete AFP and HCG

Gross →

- Usually unilateral (80–90%)
- External surfaceround to oval, encapsulated tumors., convoluted/bosselated.
- C/S → lobulated, soft and fleshy with foci of haemorrhages and necrosis.

Microscopy →



- Tumour cells arranged in lobules
- Separated by scanty fibrous stroma.

Tumour cells →
- Uniform in appearance
- Large
- vesicular nuclei
- clear cytoplasm rich in glycogen.

Fibrous stroma → lymphocytic infiltrate (good prognostic sign)

Prognosis

- All dysgerminomas are malignant.
- Dysgerminoma is treated surgically
- Highly radiosensitive and also responsive to chemotherapy.
- 5-year survival for patients with stage I is almost 100%.

b) Endodermal Sinus (Yolk Sac) Tumour

Introduction →

- Malignant, aggressive, rapidly growing germ cell tumor.

Most common →

- Second most common germ cell tumour
- Most deadly malignant ovarian germ cell tumor.

Age group →

Children or young women.

Tumour markers →

- α-fetoprotein
- α1-antitrypsin

Gross →



- Usually unilateral
- Solid with areas of cystic degeneration

Microscopy →



- Tumour cells form a variety of patterns → loose reticular network (Most common), papillary, tubular and solid arrangement.
- Tumour cells→
    - Flattened to cuboid epithelial cells
    - Clear vacuolated cytoplasm.

Characteristic features→
- Schiller-Duval bodies → glomerulus-like structure composed of a central blood vessel enveloped by germ cells within a space lined by germ cells
- PAS positive Hyaline droplets → intracellular and extracellular

Prognosis:

- Aggressive and rapidly growing tumor.

c) Teratomas

Introduction →

- Arise from Totipotent cells→ Have the capacity to differentiate into all 3 germ cell layers → ectoderm, mesoderm and endoderm

Most common →

- Most common **benign tumor of ovary** in reproductive age group.
- Most common **benign neoplasm diagnosed during pregnancy**.
- Most common **Germ cell tumor.**
- It is the commonest tumor to **undergo torsion.**

Age group →

40-50 years

Tumour markers →

AFP in immature teratoma

3 types:
1. Mature (benign)
2. Immature (malignant)
3. Monodermal or highly specialized teratomas.

1) Mature (Benign) Teratoma

- Also known as →**dermoid cyst**
- Have predominant **ectodermal elements**



Gross→
- A unilocular cyst
- C/S → cyst is filled with paste-like sebaceous secretions and desquamated keratin admixed with masses of hair.
- A solid prominence seen where tissue elements such as tooth, bone, cartilage and various other odd tissues are present → Rokitansky's protuberance



Microscopy→
- Cyst wall consists of skin (stratified squamous epithelium) with skin appendages (sebaceous glands, hair shafts, and other skin adnexal structures).
- In most cases, structures from all 3 germ layers can be identified, which includes:
- Ectoderm (e.g. skin, neural tissue, glia)
- Mesoderm (e.g. smooth muscle, cartilage, bone, fat)
- Endoderm (e.g. respiratory tract epithelium, gut, thyroid)

2) Immature (Malignant) Teratoma

- Contain **immature or embryonal structures** in contrast to the mature or adult structures of the benign teratomas.

Gross→
- Unilateral solid mass
- C/S → variegated appearance revealing areas of haemorrhages, necrosis, tiny cysts and heterogeneous admixture of various tissue elements.



Microscopy→
- Parts of tumour may show mature tissues, while most of it is composed of immature tissues having an embryonic appearance.
- Prognostic factor → Relative amount of immature neural tissue.

3) Monodermal (Specialised) Teratoma

- Composed entirely of one tissue type.
- Most common among these → struma ovarii
- struma ovarii → teratoma composed exclusively of **thyroid tissue** → recognisable grossly as well as microscopically.
- Rarely struma ovarii hyperfunctioning → **hyperthyroidism.**



Gross→
Cystic lesion composed entirely of mature thyroid tissue



Microscopy →
Appearance of a follicular adenoma of the thyroid.

Prognosis
- Mature Teratoma → good prognosis
- Immature Teratoma → extremely poor prognosis

Germ Cell Tumors

	Dysgerminoma	Yolk sac tumour	Teratoma
Introduction			
Most common			
Age group			
Tumour markers			
Gross			
Microscopy			
Prognosis			

(20.09) GESTATIONAL TROPHOBLASTIC DISEASE

University Exams

Long Questions

1. Classify Gestational trophoblastic Disease. Describe gross and microscopy of choriocarcinoma.
2. Classify Gestational trophoblastic Disease. Describe gross and microscopy of Complete Mole, Partial Mole and Invasive Mole
3. Discuss pathogenesis and morphology of trophoblastic diseases.
4. Describe the etiology, morphology, clinical features and spread of choriocarcinoma.

Short Questions

1. Enlist types of vesicular moles.
2. etiopathogenesis of complete and partial Hydatidiform mole
3. Write the karyotype and morphological features of complete mole.
4. Write the karyotype and morphological features of partial mole.
5. Morphology of hydatidiform mole.
6. Laboratory diagnosis of hydatidiform mole.
7. Differences between hydatidiform mole and choriocarcinoma.
8. Microscopy of choriocarcinoma.

Overview

- Introduction
- Classification

1. Complete Mole
2. Partial Mole
3. Invasive Mole
4. Choriocarcinoma

} Pathogenesis, Gross, Microscopy, Clinical features, Diagnosis, Spread

Introduction

Consists of tumors and tumor-like lesions characterized by proliferation of placental tissue (villous or trophoblastic)

Classification

Molar Pregnancy	Neoplasm
Complete Partial Invasive	Choriocarcinoma

1) Complete Hydatidiform Mole

Pathogenesis →

- Results from fertilization of an egg that has lost its chromosomes → either by one or two sperms.
- So characteristic feature → complete absence of maternal chromosomes (empty ovum) and the genetic material is completely paternally derived (from sperm).
- Whole placenta → neoplastic
- Fetal parts → Not present
- Risk for choriocarcinoma → Increased

Most commonly (90%)

- Fertilization of an empty egg/ovum by a one sperm.
- Chromosomes of sperm (23, X not 23, Y) undergoes duplication → Androgenesis.
- Karyotype → 46, XX diploid pattern, all derived from sperm. Never 46YY.

Less commonly (10%)

- Fertilization of an empty ovum by two sperm (dispermy).
- Depending on genetic material of two sperms (23, X or 23, Y)
- Karyotype → 46, XX (two sperms with 23, X) or 46, XY (one sperm with 23X and other with 23Y)



COMPLETE MOLE

Gross →

- Uterus is enlarged and characteristically filled with grape-like vesicles.
- Vesicles contain clear watery fluid.



Microscopy →

- Large, round, oedematous and acellular villi due to hydropic degeneration forming central cisterns.

- Decreased vascularity of villous stroma.
- Trophoblastic proliferation in the form of masses and sheets of both cytotrophoblast and syncytiotrophoblast circumferentially around villi.



Clinical features →

- Abdominal swelling (due to rapid increase in uterine size) mistaken for normal pregnancy
- Rapid increase is disproportionate to stage of pregnancy
- Vaginal bleeding
- Passage of grape-like masses per vaginum
- Nausea, vomiting



Diagnosis →

Serial determination of **β**-hCG → High levels.

2) Partial Hydatidiform Mole

Pathogenesis →

- Fertilization of a single ovum with two sperms (23, X or 23, Y).
- Karyotype → triploid {(e.g. 69, XXY- ova (23X) + sperm (23X + 23Y) or 69, XYY- ova (23X) + sperm (23Y + 23Y)} or even occasionally tetraploid (92, XXXY).
- Fetal parts → present
- Risk for choriocarcinoma → Not increased



PARTIAL MOLE

Gross →

- Uterus is smaller than expected and contains some cystic villi.
- A foetus with multiple malformations present.



Microscopy →



- Some villi show oedematous change while others are normal or even fibrotic.
- Trophoblastic proliferation is focal.

Clinical features →

- Fetus usually dies after 10 weeks gestation
- Mole (pregnancy) is aborted shortly thereafter
- Uterine size small or appropriate for gestational age

Diagnosis →

Serial determination of **β**-hCG → High HCG but not as complete.

3) Invasive Mole (Chorioadenoma Destruens)

Pathogenesis →

Complete or partial hydatidiform mole
↓
Penetrates or perforates uterine wall
↓
Enter into veins of myometrium,
↓
spread to distant sites → mostly lungs
↓
Invasive Mole (Chorioadenoma Destruens)

- Occurs in about 15% of complete moles and rarely in partial mole.
- Can cause hemorrhage and uterine perforation.

Gross →

Invasion of molar tissue into uterine wall

Microscopy →

Lesion is benign and identical to classic mole but has potential for haemorrhage.

Clinical features →

- Irregular uterine enlargement
- Vaginal bleeding

Diagnosis →

Serial determination of **β**-hCG → elevated more in both blood and urine as compared with levels in normal pregnancy.

Spread →

- Local spread → into parametrial tissue and blood vessels.
- Blood spread → Lungs and brain

4) Choriocarcinoma

Pathogenesis →

Highly malignant and widely metastasising tumour of trophoblast

- 50% of cases occur → following hydatidiform mole
- 25% → following spontaneous abortion
- 20% → after an otherwise normal pregnancy
- 5% → develop in an ectopic pregnancy.



Gross →

Tumour appears as haemorrhagic, soft and fleshy mass.



Microscopy →

- Absence of identifiable villi.
- Masses and columns of highly anaplastic and bizarre cytotrophoblast and syncytiotrophoblast cells which are intermixed.
- Invariable presence of haemorrhages and necrosis.
- Invasion of underlying myometrium and other structures, blood vessels and lymphatics.

Clinical features →

- Vaginal bleeding following a normal or abnormal pregnancy.
- Metastases in the brain or lungs.

Diagnosis →

Persistently high levels of **β**-hCG in the plasma and urine.

Spread →

Haematogenous metastases → in lungs, vagina, brain, liver and kidneys.



Feature	Complete Mole	Partial Mole	Invasive Mole	Choriocarcinoma
Pathogenesis				
Gross				
Microscopy				
Clinical features				
Diagnosis				
Spread				

(21.01) STROMAL BREAST TUMORS (FIBROSDENOMA AND PHYLLODES)

University Exams

Long questions

1. Write important gross and microscopic features of fibroadenoma and Phyllodes tumor of breast.
2. Mention classification of breast tumors. Write gross and microscopic features of stromal tumours of breast.

Short questions

1. Types of Fibroadenoma of breast.
2. Describe gross and microscopic features of fibroadenoma of breast.
3. Microscopy and types of Phyllodes tumor.
4. Classify breast tumors. Write a note on fibroadenoma of breast.

Overview

- Normal Breast histology
- 1. Fibroadenoma
  2. Phyllodes tumor

  } Introduction, Age, Clinical features, Gross, Microscopy
- Differences between Fibroadenoma and Phyllodes Tumor

Normal Breast histology





1) Fibroadenoma

Introduction →

Most common benign tumor of Breast

Age →

15 to 30 years of age

Clinical features →

solitary, discrete, freely mobile nodule within breast.



Gross

- **Small** (2-4 cm diameter)
- **Solitary**
- **Well encapsulated**
- Spherical mass
- C/S → firm, grey-white, show **slit-like spaces** formed by compressed ducts.



Microscopy

2 types of patterns →

1. Intracanalicular pattern → Stroma compresses ducts → reduced to slit-like clefts lined by ductal epithelium
2. Pericanalicular pattern → Encircling masses of fibrous stroma around patent or dilated ducts.



2) Phyllodes Tumor (Cystosarcoma Phyllodes)

Introduction →

Phyllodes = leaf-like

Age →

30 to 70 years (peak 60 years)

Clinical features →

- Aggressive clinical behavior.
- Local recurrences are much more frequent than metastases.

Grossly

- 10-15 cm in diameter
- Round to oval, bosselated
- Less fully encapsulated than a fibroadenoma.
- C/S → grey-white with cystic cavities, areas of hemorrhages, necrosis and degenerative changes

Microscopy

- Extremely hypercellular stroma
- Stromal overgrowth accompanied by benign ductal structures → typical Leaf like architecture
- Phyllodes tumor ressembles fibroadenoma except for marked stromal overgrowth.



WHO Classification	Benign PT	Borderline PT	Malignant PT
Stromal cellularity	Modest	Modest	Marked
Cellular pleomorphism	Little	Moderate	Marked
Mitoses	Few, if any	Intermediate	Numerous (>10/HPF)
Margins	Well circumscribed (Pushing)	Intermediate	Invasive

Stromal pattern	Uniform stromal distribution	Heterogenous stromal distribution	Marked stromal overgrowth
Heterologous stromal distribution	Rare	Rare	Not uncommon
Overall average distribution (%)	60	20	20

Fibroadenoma	Phyllodes Tumor
• 20-30 yrs old • 2-3cm • Firm, rubbery, painless, movable, well-circumscribed mass • Epithelial elements and connective tissue stroma • Rapid growth but not premalignant	• Any age • >5cm • Firm, mobile, less circumscribed, nontender mass • Epithelial elements and connective tissue stroma (more cellular, more pleomorphic and mitotically more active) • Tendency to grow rapidly and aggressively

(21.02) BREAST CARCINOMA

University Exams

Long questions

1. A 52-year-old female presented with lump in right breast which was noticed 6 months back. On examination of lump, it was firm to hard and fixed to underlying structures and skin with 5 palpable lymph nodes in the right axilla. After FNAC she underwent mastectomy:

   a) What is probable diagnosis?
   b) Write risk factors of the condition.
   c) Write morphology of lesion in breast.

2. Write molecular classification of breast cancer. Discuss etiopathogenesis and morphology of carcinoma breast.
3. Discuss etiology, gross,microscopic appearance and clinical features of carcinoma of breast.
4. How will you classify Breast Carcinoma? Discuss Grading, Staging and Prognostic Factors in Breast Cancer.
5. Discuss morphology of Ductal Carcinoma in – situ (DCIS) and lobular Carcinoma in – situ (LCIS)
6. Discuss morphology of Invasive ductal carcinoma and Invasive lobular carcinoma

Short questions

1. Carcinoma of breast.
2. Carcinoma in situ of breast.
3. Morphology of Lobular carcinoma of breast
4. Morphology of infiltrating duct carcinoma.
5. Etiology of carcinoma of breast.
6. Classification of tumors of breast.
7. Enumerate significant risk factors in etiology of breast carcinoma.
8. Paget's disease of the nipple (breast).
9. Prognostic factors of carcinoma breast.
10. Molecular classification of breast cancer
11. TMN Staging of breast cancer

Very Short questions

1. Enumerate four risk factors for breast cancer.
2. Classification of breast carcinoma.
3. Histopathological types (microscopic variants) of breast carcinoma.
4. Paget's disease of breast (nipple}- morphology.
5. Prognostic factors of carcinoma of breast.

Overview

- Introduction
- Age
- Clinical features
- Triple technique
- Risk factors
- Classification
- Individual tumor microscopy → DCIS, LCIS, IDC, ILC

- Paget's Disease of Nipple → Introduction, Pathogenesis, Gross, Microscopy
- Molecular classification
- TMN Staging
- Prognostic factors

Introduction→

- Most common Cancer of females in world
- Cancer of male breast is rare
- Male: Female breast cancer ratio → 1:150



Age →

Perimenopausal age group

Clinical features

Solitary, painless, palpable lump detected by self-examination

Triple technique

PALPATION



MAMMOGRAPHY



FNAC

Risk Factors

1. Geographic and racial factors→ 4-6 times higher in developed countries and low in developing countries of Asia and Africa
2. Family history → First-degree relatives (mother, sister, daughter) of women with breast cancer have 2 to 6-fold higher risk of development of breast cancer.
3. Menstrual and obstetric history → Total length of menstrual life is directly related to increased risk. Increased risk of breast cancer in→

   d) Early menarche
   e) Nulliparity
   f) Delayed menopause

4. Estrogen exposure → Menopausal hormone therapy increases risk of breast cancer

   - Reducing endogenous estrogens by oophorectomy decreases risk
   - Drugs that block estrogenic effects (e.g., tamoxifen) or block the formation of estrogen (e.g., aromatase inhibitors) decrease risk

5. Fibrocystic change → Atypical epithelial hyperplasia → 5-fold higher risk of developing breast cancer

6. Dietary factors →

    - Consumption of large amounts of animal fats, high calorie foods
    - Cigarette smoking
    - Alcohol consumption

7. Exercise → Protective effect
8. Breast density → Women with very dense breasts on mammography have higher risk of breast cancer compared to women with the lowest density.
9. Environmental toxins → organochlorine pesticides have estrogenic effects on humans.
10. Radiation exposure → Radiation to the chest (cancer therapy, atomic bomb exposure, nuclear accidents)
11. Genetic Factors

    BRCA 1 gene →
    - Located on chromosome 17
    - Breast and ovarian cancer in inherited cases

    BRCA 2 gene →
    - Located on chromosome 13
    - Breast and ovarian cancer in inherited cases

    Mutation in p53 →
    - Located on chromosome 17

Classification







Individual Tumor microscopy

Ductal Carcinoma in – situ (DCIS)

4 patterns →

a) Solid pattern → Plugging of ductal lumina with tumor cells.
b) Comedo pattern → centrally placed necrotic debris surrounded by neoplastic cells
c) Papillary pattern → Intraductal papillary projections of tumor cells which lack a fibrovascular stalk
d) Cribriform pattern → Punched out fenestrations in intraductal tumor.



Lobular Carcinoma in – Situ (LCIS)



- Filling up of terminal ducts by uniform cells which are loosely cohesive and have small, rounded nuclei with indistinct cytoplasmic margins
- Higher incidence of developing a **contralateral breast cancer**

Invasive Duct Carcinoma (IDC)

- Anaplastic tumor cells forming solid nests, cords, poorly-formed glandular structures
- Infiltration of tumor cells into diffuse fibrous stroma and fat.
- Invasion into perineural spaces
- Lymphatic and vascular invasion.



Invasive Lobular Carcinoma (ILC)

- Single file (Indian file) linear arrangement of stromal infiltration by the tumor cells
- Tumor cells → round and uniform, regular with very little pleomorphism and infrequent mitoses.



INDIAN FILE PATTERN

Paget's Disease of Nipple

Introduction →

- Eczematoid lesion of nipple associated with an invasive or non-invasive ductal carcinoma of underlying breast.
- Nipple bears a crusted, scaly and eczematoid lesion with a palpable subareolar mass
- Malignant lesion

Pathogenesis



Tumor cells from underlying ductal carcinoma
↓
Migrated up into lactiferous ducts
↓
Invaded epidermis of nipple
↓
Producing skin lesions

Gross
Skin of nipple and areola is **crusted, fissured and ulcerated** with oozing of serosanguineous fluid from erosions

Microscopy



- Presence of Paget's cells singly or in small clusters in epidermis
- Paget's cells → larger than epidermal cells, spherical, hyperchromatic nuclei with cytoplasmic halo

Molecular classification

a) ER / PR Receptors
b) HER2/neu (cerbB2) Receptors





	ER/PR	Her-2
Luminal A	+	-
Luminal B	+	+
Basal	-	-
Her-2 Positive	-	+

TNM Staging



Tumor Size **T**	Tumor size < 2 cm T1	Tumor size 2-5 cm T2	Tumor size > 5 cm T3	Tumor extends to skin or chest wall T4
Lymph Nodes **N**	**NO** No lymph node metastasis	**N1** Metastasis to ipsilateral, movable, axillary LNs	**N2** Metastasis to ipsilateral fixed axillary, or IM LNs	**N3** Metastasis to infraclavicular/ supraclavicular LN, or to axillary and IM LNs
Metastasis **M**	**M0** No distant Metastasis	**M1** Distant metastasis		

Stage	TNM	Category
I	T_1, N_O, M_0	Early cancer (operable, curable)
II	T_1, N_1, M_0 T_2, NO_1, M_0	Early cancer (operable, curable)
III	Any T, N_{2-3}, M_0 T_3, Any N, M_0	Locally advanced breast cancer (Neoadjuvant treatment to make operable)
IV	Any T, Any N, M_1	Metastatic (Not curable, Inoperable, Palliative care)

Prognostic Factors

Based on →

- Routine histopathology criteria
- Hormone receptor status
- Biological indicators

FACTOR	FAVOURABLE PROGNOSIS	POOR PROGNOSIS
I. ROUTINE HISTOPATHOLOGY CRITERIA		
i. Histologic type	Medullary ca., Tubular ca., Mucinous (colloid) ca., Lobular ca.	Inflammatory ca.
ii. Tumor size	Size < 1 cm	Size > 1 cm
iii. Histologic grading (Score range of 3-9)	Low grade (grade I) Tumor = score 3-5, moderate grade (grade II) Tumor = score 6-7	High grade (grade III) Tumor= score 8-9
iv. Axillary nodal status	Node negative: recurrence after 10 years 10-30%; sentinel node negative	Node positive: recurrence after 10 years 70%; sentinel node positive
v. Lymphatic and / or vascular invasion	Negative for both: good	Positive for one or both: poor
II. **HORMONE RECEPTOR STATUS**		
i. Oestrogen-progesterone receptors (ER-PR)	ER-PR positive cases respond better to adjuvant therapy	ER-PR negative cases respond poorly to adjuvant therapy
ii. HER-2/neu (C-erb B-2)	Under expression	Overexpression (predictive of response to erceptin)
III. BIOLOGICAL INDICATORS		
vi. Mitotic index (by Ki67, MIB-1)	Low mitotic count	High mitotic count
vii. Angiogenesis (VEGF, CD31, CD34, micro vessel density counts)	Angiogenic activity low	High angiogenic activity
viii. Oncogene dysregulation		
a) BRCA 1, BRCA2	BRCA negative	BRCA positive
b) p53	p53 positive respond better to chemotherapy and radiotherapy	p53 negative respond poorly to chemotherapy and radiotherapy

(22.01) AUTOIMMUNE DISORDERS OF THYROID (HASHIMOTO THYROIDITIS, GRAVES DISEASE)

University Exams

Long Questions

1. Enumerate non-neoplastic lesions of thyroid. Discuss pathogenesis, Gross, Microscopy and clinical features of Hashimoto thyroiditis.

Short Questions

1. Etiopathogenesis and morphology of Hashimoto's thyroiditis
2. Pathogenesis and microscopy of Grave's disease

Overview

- Normal thyroid → Histology
- Autoimmune disorders of thyroid → Introduction

1. Graves' Disease
2. Hashimoto Thyroiditis

} Introduction, Age, Gender, Pathogenesis, Gross, Microscopy, Clinical features

Normal thyroid → Histology

- Follicular cells secrete →**T4 and T3**
- Parafollicular cells secrete → **calcitonin**





Autoimmune disorders of thyroid → Introduction

2 extremes of autoimmune thyroid disorders →

1. Graves' disease → hyperthyroidism
2. Hashimoto thyroiditis → hypothyroidism

1) Graves' Disease

Introduction →

- An autoimmune disorder presenting clinically with feature of hyperthyroidism
- Most common cause of endogenous Hyperthyroidism

Age →

20 to 40 years of age

Gender →

F > M

Pathogenesis

Ab against TSH receptors (Thyroid stimulating immunoglobulins (TSI) and long-acting thyroid stimulator (LATS) are produced)
↓
Ab react with TSH receptor present on follicular cells of thyroid
↓
Thyroid cells are stimulated
↓
T3, T4 (free and bound) is raised
TSH is suppressed
↓
Hyperthyroidism
↓
Graves' disease



Gross

- Thyroid gland is symmetrically enlarged
- Diffuse hypertrophy and hyperplasia of thyroid follicular epithelial cells.
- C/S → soft and meaty (resembling normal muscle)

Microscopy



- Epithelial hyperplasia and hypertrophy →
  - Increased height of follicular lining cells
  - Formation of papillary infoldings of piled up epithelium into lumina of follicles
- Colloid → markedly diminished and is lightly staining, watery and finely vacuolated.
- Stroma → Lymphoid inflammation

Clinical Features

Triad of Graves' disease →
1. Hyperthyroidism → diffuse hyperplasia of the thyroid
2. Infiltrative ophthalmopathy→ Exophthalmos
3. Pretibial myxedma → Localized infiltrative dermopathy

2) Hashimoto Thyroiditis (Chronic Lymphocytic Thyroiditis)

Introduction →

- An autoimmune disorder that results in destruction of thyroid gland and gradual and progressive thyroid failure
- Most common cause of hypothyroidism in areas of world where iodine levels are sufficient

Age →

45 to 65 years of age

Gender →

F > M

Pathogenesis

Breakdown in self-tolerance to thyroid autoantigens.

Genes regulating T-cell responses including cytotoxic T lymphocyte associated antigen-4 (CTLA4) and protein tyrosine phosphatase-22 (PTPN22)
↓
Polymorphisms in these genes
↓
Breakdown in self-tolerance
↓
Induction of thyroid autoimmunity
↓

a. CD8+ cytotoxic T cell-mediated cell death
b. Cytokine-mediated cell death
c. Antithyroid antibodies



Gross



- Diffuse and symmetric enlargement of thyroid gland
- Gland → firm and nodular.
- Capsule → Intact
- C/S → pale, gray-tan

Microscopy



- Extensive infiltration of parenchyma by a mononuclear inflammatory infiltrate containing small lymphocytes, plasma cells→ germinal centers are formed
- Thyroid follicles are atrophic
- Thyroid follicles are lined by epithelial cells with abundant eosinophilic, granular cytoplasm → Hurthle cells

Clinical features

- Painless enlargement of thyroid
- Hypothyroidism gradually develops
    - Early stages → transient thyrotoxicosis → Hashitoxicosis →due to destruction of thyroid follicles with secondary release of thyroid hormones
    - Late stages → Hypothyroidism

Feature	Graves' Disease	Hashimoto Thyroiditis
Introduction		
Age		
Gender		
Pathogenesis		
Gross		
Microscopy		
Clinical features		

(22.02) GOITER

University Exams

Short Questions

1. Pathogenesis of Toxic goiter.
2. Iodine deficiency goiter.
3. Stages in evolution of multinodular goiter.
4. Morphology of Multinodular goiter
5. Goitrogens.
6. Gross and Microscopy of Diffuse goiter and Multinodular goiter

Overview

- Introduction
- Pathogenesis →

    1. Endemic Goiter
    2. Sporadic Goiter

- Types →

    1. Diffuse goiter
    2. Multinodular goiter

    } Introduction, Gross, Microscopy, Clinical features, Functional status

Introduction

- Most common manifestation of thyroid disease

Thyroid hormone deficiency
↓
Compensatory hyperplasia and hypertrophy of follicular epithelium
↓
Enlargement of thyroid without hyperthyroidism
↓
End-result → Euthyroid state
↓
Goiter



Pathogenesis

1. Endemic Goiter
2. Sporadic Goiter

1) Endemic Goiter

- Prevalence of goiter in a geographic area > 10% of population
- Causes →

a) Deficiency of iodine →

- Low iodine in soil, water, and food:
- Common in mountainous areas (e.g. Himalayas)
- Dietary supplementation of iodine has reduced frequency and severity of endemic goiter

b) Goitrogens →

- Substances ingestion of which interferes with thyroid hormone synthesis.
- Eg. Brassicaceae (Cruciferae) family → cabbage, cauliflower, Brussels sprouts, and turnips and Cassava root2) Sporadic Goiter



Autosomal recessive conditions
↓
Hereditary enzymatic defects
↓
Interfere with thyroid hormone synthesis
↓
Hypothyroidism

- e.g. dyshormonogenetic goiter

Types

1. Diffuse goiter (simple nontoxic goiter or colloid goiter)
2. Multinodular goiter (Adenomatous goiter)

1) Diffuse goiter

Introduction →

- D diffuse enlargement of thyroid gland without any nodularity.
- Consists of large thyroid follicles distended with colloid → Colloid goiter



Gross →

- Enlargement of thyroid gland→ symmetric and diffuse.
- C/S → gelatinous and translucent brown

Microscopy→

2 stages →

1. Hyperplastic stage → Tall columnar follicular epithelium showing papillary infoldings
2. Involution stage → Large follicles distended by colloid and lined by flattened follicular epithelium



Clinical features

- Clinically euthyroid
- Mass effects from enlarged thyroid gland

Functional status

- Most patients → euthyroid
- Serum T3 and T4 levels → normal
- Serum TSH → elevated or at upper range of normal

2) Multinodular goiter

Introduction →

In long-standing simple goiters
↓
Recurrent episodes of hyperplasia and involution
↓
Produce a more irregular enlargement of thyroid
↓
Multinodular goiters

- Mistaken for neoplasm

Gross →

- Thyroid → asymmetric and extreme enlargement
- Nodularity with poor encapsulation
- Fibrous scarring
- Haemorrhages
- Focal calcification
- Cystic degeneration



Microscopy →

- Partial or incomplete encapsulation of nodules.
- Follicles lined by flat to high epithelium.
- Areas of hemorrhages, hemosiderin-laden macrophages and cholesterol crystals.
- Fibrous scarring with foci of calcification.



Clinical features

- Usually asymptomatic
- Mass in neck.
- Large multinodular goiter → compression of surrounding structures and may cause:

- Compression of trachea → Airway obstruction
- Compressing the esophagus → Dysphagia
- Compression of recurrent laryngeal nerve→ Hoarseness

Functional status

- Most patients → euthyroid → T4, T3, and TSH → normal
- Some patients → subclinical hyperthyroidism → reduced TSH levels.
- Some patients → hyperthyroidism (toxic multinodular goiter) → Plummer syndrome.

Feature	Diffuse goiter	Nodular goiter
1. Nomenclature	Simple goiter, hyperplastic goiter, nontoxic goiter	Multinodular, adenomatous goiter
2. Etiology	Graves'disease, thyroiditis, puberty	Endemic thyroiditis, cancer
3. Pathogenesis	Hyperplasia-involution	Repeated cycles of hyperplasia with growth and involution with fibrosis
4. Composition	Cellular-rich	Colloid-rich
5. Gross	Moderate, symmetric, diffuse enlargement, colloid-filled follicles, gelatinous	Nodular asymmetric, haemorrhages, scarring, cystic change, calcification
6. Microscopy	Hyperplastic phase: papillary infoldings, Involution stage: large colloid filled follicles with flat epithelium	Incomplete encapsulation, nodularity, variable-sized follicles, fibrous scarring, haemorrhages, calcification, cyst formation
7. Functional status	Hyperthyroidism, euthyroid	Hypothyroidism, euthyroid

(22.03) THYROID NEOPLASM

University Exams

Long questions

1. Classify Thyroid tumours. Discuss gross, Microscopy, Spread and prognosis of Papillary carcinoma of thyroid

Short Questions

1. Classify thyroid tumors. Describe in detail any one of them.
2. Mention major subtypes of carcinomas of thyroid. Write a note on genetic alterations in follicular cell derived malignancies of thyroid.
3. Histopathology of Papillary carcinoma of thyroid
4. Gross and microscopy of Medullary carcinoma of thyroid.
5. Describe gross and microscopic features of Follicular carcinoma of thyroid.

Very Short Questions

1. Hurthle cell.
2. Types of thyroid tumors.
3. Morphology of papillary carcinoma of thyroid.
4. Calcitonin in Medullary carcinoma of thyroid.

Overview

Classification →

1. Papillary carcinoma
2. Follicular carcinoma
3. Medullary carcinoma
4. Anaplastic carcinoma
5.

} Introduction, Cell of origin, Age, F/M ratio, Risk factors, Genetic alterations, Gross, Microscopy, Spread, Prognosis

Classification

1) Papillary Carcinoma

Introduction →

Most common type (75%).

Cell of origin →

Follicular

Age →

20-40 years

F/M ratio →

3 : 1

Risk factors →

radiation exposure in head and neck area.

Genetic alterations →

RET gene over-expression, NTRK gene rearrangement

Gross



- Well-circumscribed
- Encapsulated or
- Ill-defined (infiltrate the adjacent parenchyma).
- Gray white, firm to hard tumor
- Areas of fibrosis and calcification.

- Cut surface → papillary foci

Microscopy



1. Papillary pattern → Papillae composed of fibrovascular stalk and covered by single layer of tumor cells
2. Tumor cells →

    a) Optically clear nuclei = orphan Annie eye = ground glass appearance
    b) Nuclear grooves
    c) Intranuclear pseudo inclusions (cytoplasmic invaginations into nuclei)

3. Psammoma bodies → small, concentric, calcified spherules





Spread →

Lymphatic route

Prognosis →

- Excellent prognosis
- Least malignant thyroid neoplasm
- 10-year survival → 80 – 95 %

2) Follicular Thyroid Carcinoma

Introduction →

2nd common type of thyroid cancer (10-20% of all thyroid carcinomas)

Cell of origin →

Follicular

Age →

40 to 50 yrs.

F/M ratio →

2.5 : 1

Risk factors →

Iodine deficiency (long standing multinodular goiter).

Genetic alterations →

RAS mutation, PAX-PPAR g1 fusion

Gross

- Large tumors penetrate the capsule and infiltrate into the adjacent structures in the neck.
- Follicular thyroid carcinoma differentiated from follicular adenoma:
    - Capsular invasion
    - Vascular invasion



Microscopy

1. Follicular pattern → follicles of various sizes
2. Tumor cells → hyperchromatic nuclei and cytoplasm resembles that of normal follicular cells.
3. Vascular invasion and direct extension
    - Diagnosis is confirmed by tissue biopsy (histopathology) and cytology is inconclusive (follicular adenoma versus carcinoma).
    - Hurthle cell or oncocytic carcinoma is a variant of follicular carcinoma.



Hurthle Cell Tumors

- Variant of follicular carcinoma.
- More than 50% Hurthle cell.
- Usually adult females.
- Solid, tan, well-vascularised, and encapsulated.
- Usually follicular, may be trabecular/solid or papillary.
- Histology of Hurthle cells → granular, acidophilic cytoplasm (filled with mitochondria).
- Aggressive carcinoma.



Spread →

Hematogenous to bone, Lung, Liver, etc

Prognosis →

- Good prognosis
- 10-year survival → 50 – 70 %

3) Medullary Thyroid Carcinoma

Introduction →

Constitutes 5% of thyroid neoplasms.

Cell of origin →

Para- Follicular or C-cells → secrete **calcitonin**

Age →

40 to 50 yrs.

F/M ratio →

1 : 1

Risk factors →

Familial type associated with MEN syndrome 2A or 2B

Genetic alterations →

RET point mutation

Gross

- Firm, pale gray, and infiltrative.
- Areas of necrosis and hemorrhage
- Extend through capsule of thyroid





Microscopy

- Tumor cells → Nests of Tumor cells separated by fibrovascular septa.
- Tumor cells are uniform
- Amyloid stroma → Tumor cells are separated by **amyloid** stroma derived from altered calcitonin
- Areas of irregular calcification

Spread →

Lymphatic and blood both

Prognosis →

- Good prognosis
- 10-year survival → 60 – 70 %

4) Anaplastic Carcinoma

Introduction →

- Constitutes < 5% of thyroid neoplasms.
- One of the most malignant Tumor in human
- Tumor is widely aggressive and rapidly growing

Cell of origin →

Follicular cell

Age →

Old age (7th-8th decades)

F/M ratio →

1.5 : 1

Genetic alterations →

p53 loss, (β-catenin mutation

Gross

Diffusely infiltrative tumor.

Microscopy

- Tumor is poorly-differentiated
- Tumor is composed of 3 types of cells occurring in varying proportions:
    a) Squamoid: keratinize.
    b) Spindle cell: Storiform, form bone/cartilage.
    c) Giant cells: scattered among spindle cell pattern.





Spread →

1. Local spread into thyroid capsule and adjacent neck structures.
2. Hematogenous spread to lungs.

Prognosis →

- Worst prognosis
- 10-year survival → 5 – 10 %
- Essentially 100% mortality (mean survival 2 months) from local extension into neck structures.

Feature	Papillary carcinoma	Follicular carcinoma	Medullary carcinoma	Anaplastic carcinoma
Introduction	75-80%	10-20%	5%	5%
Cell of origin	Follicular	Follicular	Parafollicular	Follicular
Age	All ages	Middle to old age	Middle to old age; familial too	Old age
F / M ratio	3:1	2.5:1	1:1	1.5:1
Relation to radiation	Maximum	Present	None	Present
Genetic alterations	RET gene over-expression, NTRK gene rearrangement	RAS mutation, PAX-PPAR g1 fusion	RET point mutation	p53 loss, (β-catenin mutation
Gross	Small, multifocal	Moderate size, nodular	Moderate size	Invasive growth
Microscopy	Nuclear features, papillary pattern	Vascular and capsular invasion	Solid nests, amyloid stroma	Undifferentiated, spindle-shaped, giant cells
Regional metastases	Common	Rare	Common	Common
Distant metastases	Rare	Common	Rare	Common
Prognosis → 10-year survival	80-95%	50-70%	60-70%	5-10% (median survival about 2 months)

(22.04) DIABETES MELLITUS

University Exams

Long Questions

1. Discuss pathogenesis of type 2 diabetes mellitus. Mention its major long-term complications.
2. Discuss laboratory diagnosis of Diabetes mellitus.
3. Describe Risk factors, etiology, pathogenesis and classification of diabetes mellitus. Discuss laboratory diagnosis of diabetes mellitus.

Short Questions

1. Differences between Type I DM and Type II DM
2. Etiopathogenesis of type I diabetes mellitus.
3. Pathogenesis of type II diabetes mellitus.
4. Describe method and interpretation of oral glucose tolerance test.
5. Ketonuria.
6. Glycosuria.
7. Glucose tolerance curve.
8. Write short Questions on Glycosylated Hemoglobin.

Very Short Questions

1. List the complications of diabetes mellitus.
2. Renal glycosuria.
3. HbA1C

Overview

- Normal Pancreas → Histology
- Diabetes Mellitus →
    - Introduction
    - Classification
    - Risk factors
    - Insulin regulation → Synthesis, release, action
    - Pathogenesis → Type I DM and Type II DM
    - Clinical features
    - Differences between Type I DM and Type II DM
    - Complications
    - Diagnosis

Normal Pancreas → Histology

1. Exocrine part → secretes enzymes
2. Endocrine part → secretes hormones



CELL	HORMONE SECRETED
Alpha (α) cell	Glucagon
Beta (β) cell	Insulin
Delta (δ) cell	Somatostatin
Pancreatic polypeptide (PP) cells	Pancreatic polypeptide

Diabetes Mellitus

Introduction →

- Diabetes mellitus (DM) is a group of metabolic disorders sharing the common feature of hyperglycemia
- DM is diagnosed when blood glucose levels becomes too high



Body controls how much glucose is in blood relative to how much is in the cell by 2 hormones →

1. Insulin → (Glucose → Glycogen)
2. Glucagon → (Glycogen → Glucose)

DM
↓
Insulin absent or non-functional
↓
Body has trouble in moving glucose from Blood to Cells
↓
Increase glucose in blood (Hyperglycemia) → secondary damage in multiple organ systems
↓
Decrease glucose in cells → Cells starve for energy despite having glucose right on doorstep



Classification

Type 1 DM →
- 10% cases of DM.
- Juvenile-onset diabetes (JOD) → due to its occurrence in younger age
- Insulin-dependent DM (IDDM) → because these patients have absolute requirement for insulin replacement as treatment.

Type 2 DM →
- 90% cases of DM.
- Maturity-onset diabetes
- Non-insulin dependent diabetes mellitus (NIDDM)

Type 1 diabetes (β-cell destruction, usually leading to absolute insulin deficiency)
Immune-mediated Idiopathic
Type 2 diabetes (combination of insulin resistance and β-cell dysfunction)
Genetic defects of β-cell function
Maturity-onset diabetes of the young (MODY), caused by mutations in: Hepatocyte nuclear factor 4α (HNF4A), M0DY1 Glucokinase (GCK), M0DY2 Hepatocyte nuclear factor 1α (HNF1A), M0DY3 Pancreatic and duodenal homeobox 1 (PDX1), M0DY4 Hepatocyte nuclear factor 1β (HNF1B), M0DY5 Neurogenic differentiation factor 1 (NEUROD1), M0DY6 Neonatal diabetes (activating mutations in KCNJ11 and ABCC8, encoding Kir6.2 and SUR1, respectively) Maternally inherited diabetes and deafness (MIDD) due to mitochondrial DNA mutations (m.3243A→G) Defects in proinsulin conversion Insulin gene mutations
Genetic defects in insulin action
Type A insulin resistance Lipoatrophic diabetes
Exocrine pancreatic defects
Chronic pancreatitis Pancreatectomy/trauma Neoplasia Cystic fibrosis Hemochromatosis Fibrocalculous pancreatopathy
Endocrinopathies
Acromegaly Cashing syndrome Hyperthyroidism Pheochromocytoma Glucagonoma
Infections
Type A insulin resistance Lipoatrophic diabetes

Drugs
Glucocorticoids Thyroid hormone Interferon -α Protease inhibitors b-adrenergic agonists Thiazides Nicotinic acid Phenytoin (Dilantin) Vacor
Genetic syndromes associated with diabetes
Down syndrome Klinefelter syndrome Turner syndrome Prader-Willi syndrome
Gestational diabetes mellitus

Risk Factors

1. Family history of type 2 DM
2. Obesity
3. Habitual physical inactivity
4. Race and ethnicity (Blacks, Asians, Pacific Islanders)
5. Previous identification of impaired fasting glucose or impaired glucose tolerance
6. History of gestational DM or delivery of baby heavier than 4 kg
7. Hypertension
8. Dyslipidaemia (HDL level < 35 mg/dl or triglycerides > 250 mg/dl)
9. Polycystic ovary disease and acanthosis nigricans
10. History of vascular disease

Insulin regulation

1. Synthesis
2. Release
3. Action







Pathogenesis

Type 1 DM

- Body cannot make enough INSULIN

In DM -I
↓
There is genetic abnormality and environmental trigger
↓
Loss of self-tolerance
↓
Self-reactive clonal T cells are not killed or deleted inthymus
↓
Attack β-cells of pancreas (β-cells act as autoantigens)
↓
Decrease insulin
↓
Increase glucose in blood (Hyperglycemia)





Type 2 DM

- Type 2 diabetes is a complex disease that involves an interplay of genetic and environmental factors and a proinflammatory state.

Genetic component
+
Constitutional factors such as obesity, hypertension, and level of physical activity play contributory role
↓
Insulin resistance on target organ
↓
Hyperglycaemia





Clinical Features

Classical triad →

1. Polyuria
2. Polydipsia
3. Polyphagia

- Weight Loss
- Fatigue
- frequency of infection
- Rapid Onset
- insulin Dependent
- Familial Tendency
- Peak incidence from 10 to 15 years

Polyuria
↑ Urination

Polydipisia
↑ Thirst

Polyphagia
↑ Hunger

1. Polyphagia → catabolism of proteins and fats → induce a negative energy balance → increasing appetite → polyphagia
2. Polyuria → Hyperglycemia exceeds renal threshold for reabsorption→ glycosuria → induces an osmotic diuresis → polyuria
3. Polydipsia → obligatory renal water loss → deplete intracellular water → trigger osmoreceptors of thirst centers of brain → polydipsia
4. Weight loss→ Despite increased appetite → catabolic effects prevail → weight loss



Differences between TYPE 1 DM and TYPE 2 DM

	FEATURE	TYPE 1 DM	TYPE 2 DM
1	**Frequency**	10-20%	80-90%
2	**Age at onset**	Early (below 35 years)	Late (after 40 years)
3	**Type of onset**	Abrupt and severe	Gradual and Insidious
4	**Weight**	Normal	Obese/non-obese
5	**HLA**	Linked to HLA DR3, HLA DR4, HLA DQ	No HLA association

6	**Family history**	<20%	About 60%
7	**Genetic locus**	Unknown	Chromosome 6
8	**Diabetes in identical twins**	50% concordance	80% concordance
9	**Pathogenesis**	Autoimmune destruction of (1-cells)	Insulin resistance, Impaired Insulin secretion
10	**islet cell antibodies**	Yes	No
11	**Blood Insulin level**	Decreased Insulin	Normal or Increased Insulin
12	**islet cell changes**	Insulitls, p-cell depletion	No insulitls, later fibrosis of islets
13	**Amyloidosis**	Infrequent	Common in chronic cases
14	**Clinical management**	Insulin and diet	Diet, exercise, oral drugs, Insulin
15	Acute complications	Ketoacidosis	Hyperosmolar coma

Complications

- Acute Metabolic Complications
    - Diabetic ketoacidosis (DKA)
    - Hyperosmolar hyperglycaemic nonketotic coma
- Late Systemic Complications
    - Atherosclerosis → resulting in increased risk of myocardial infarction, stroke, and lower extremity ischemia.
    - Diabetic nephropathy
    - Diabetic neuropathy
    - Diabetic retinopathy
    - Infections



Diabetic ketoacidosis

Insulin deficiency
↓
Stimulates lipoprotein lipase
↓
Breakdown of adipose stores → increase in free fatty acids (FFA)
↓
FFA reach liver → esterified to fatty acyl coenzyme A.
↓
Oxidation of fatty acyl coenzyme A → ketone bodies (acetoacetic acid and β-hydroxybutyric acid)
↓
Ketonemia
Ketonuria
↓
Urinary excretion of ketones is compromised by dehydration
↓
Systemic metabolic ketoacidosis



Hyperosmolar hyperglycemic coma

- Occurs in Type 2 diabetics
- Due to severe dehydration resulting from sustained osmotic diuresis
- Hyperglycemia → 600 to 1200 mg/dL.



Diagnosis

1. Urine Test
2. Blood Test
3. Glycosylated hemoglobin (HbA1C)

1) Urine Test

a) Glucosuria
b) Ketonuria





BENEDICT'S TEST



SEMIQUANTITATIVE INTERPRETATION

Disadvantage of relying on urinary glucose test

- Besides diabetes mellitus, glucosuria also occur in →

    1. Renal glycosuria
    2. Alimentary glucosuria

1) Renal glucosuria

- Normal renal threshold for glucose is 180 mg/dl ie in normal persons glucose will only appear in urine if blood glucose is above 180 mg/dl

Renal glucosuria
↓
Renal threshold for
glucose is reduced
↓
Although the blood glucose level is below 180 mg/dl
(i.e. below normal renal threshold for glucose)
↓
Glucose still appears
regularly and consistently in
the urine due to lowered renal threshold



2) Alimentary (lag storage) glucosuria

In some individuals after a meal
↓
A rapid and transitory rise in blood glucose level above normal renal threshold may occur
↓
During this period → glucosuria is present
↓
Unusually high blood glucose level returns to normal 2 hours after meal.

2) Blood test

Revised criteria for diagnosis of diabetes
(as per American Diabetes Association)

RANDOM VALUE	
200mg/dl (11. 1 mmol/L) or more in a symptomatic patient	Diabetes mellitus
FASTING (FOR > 8 HOURS) VALUE	
Below 100 mg/dl (< 5.6 mmol/L) **100-125 mg/dl (5.6-6.9 mmol/L)** **126 mg/dl (7.0 mmol/L) or more**	Normal fasting value Impaired fasting glucose (IFG)** Diabetes mellitus
TWO-HOUR AFTER 75 GM ORAL GLUCOSE LOAD	
< 140 mg/dl (< 7.8 mmol/L) 140-199 mg/dl (7.8-11.1 mmol/L) **200 mg/dl (11. 1mmol/L) or more**	Normal post-prandial GTT Impaired post-prandial glucose tolerance (IGT)** Diabetes mellitus

3) Glycosylated hemoglobin (HbA1C)

- Minor hemoglobin component present in normal persons.
- Long-term assessment of degree of glycemic control is better monitored by measurement of glycosylated hemoglobin (HbA1C)
- This is because non-enzymatic glycosylation of hemoglobin takes place over 90-120 days (lifespan of RBCs)
- Gives an estimate of diabetic control and compliance for preceding 3-4 months.



- Normal value < 6.5 %
- A glycated hemoglobin (HbA1C) level ≥ 6.5% → diabetics
- It is recommended that HbA1C be maintained below 7% in diabetic patients.

(22.05) ZOLLINGER – ELLISON SYNDROME (GASTRINOMA)

University Exams

Short Questions

1. Pathogenesis of Zollinger-Ellison syndrome.
2. Triad of Zollinger-Ellison syndrome.
3. Sites of Peptic ulcer in Zollinger-Ellison syndrome.

Overview

- Pathogenesis
- Triad
- Sites of Peptic ulcer

Pathogenesis

Gastrinoma (benign or malignant)
↓
Occur in duodenum or Pancreas (Non-B pancreatic islet cell Tumor)
↓
Secrete gastrin → hypergastrinemia
↓
Stimulates parietal cells of stomach
↓
Massive secretion of acid and pepsin
↓
Fulminant peptic ulcer disease
↓
Zollinger-Ellison syndrome



Triad

1. Presence of non-β pancreatic islet cell Tumor.
2. Gastric acid hypersecretion
3. Fulminant peptic ulcer disease





Sites of Peptic ulcer

Multiple and recurrent peptic ulcers

- Usual sites → stomach, first and second part of the duodenum
- Unusual sites → esophagus and jejunum.



ZOLLINGER-ELLISON SYNDROME

(22.06) CUSHING SYNDROME

University Exams

Short Questions

1. List four causes of Cushing syndrome
2. Etiopathogenesis of Cushing's syndrome.
3. Define Cushing's syndrome. Write its Clinical features

Overview

- Adrenal Gland →Anatomy
- Adrenal Gland → Physiology
- Biosynthesis of adrenal steroids
- Glucocorticoids (Cortisol) functions
- Regulation of cortisol secretion
- Hyperadrenalism → Patterns
- Cushing syndrome →
    - Definition
    - Etiopathogenesis
    - Clinical features

Adrenal Gland Anatomy

- 2 adrenal glands
- Each weighing about 4 gms
- Lie at superior poles of two kidneys
- Each gland composed of two distinct parts →
    1. Adrenal cortex
    2. Adrenal medulla



Adrenal Gland Physiology

Adrenal Cortex	Adrenal Medulla
Composed of 3 Layers	Composed of neural crest derived chromaffin cells.
1) Zona Glomerulosa (outermost) → produces mineralocorticoids (e.g., aldosterone).	Main physiologic source of catecholamines (epinephrine and norepinephrine)
2) Zona Fasciculata (middle) → produces glucocorticoids (e.g., Cortisol).	
3) Zona Reticularis (Innermost) → produces androgens or sex steroids (e.g., testosterone).	

Biosynthesis of adrenal steroids

Adrenal cortex produces 3 major classes of steroids →
1. Glucocorticoids
2. Mineralocorticoids
3. Adrenal androgens



Glucocorticoids (Cortisol) functions

- 95% of glucocorticoid activity occurs from secretion of cortisol known as hydrocortisone

Carbohydrate metabolism	• Stimulates gluconeogenesis • ↑ glycogen storage in liver cells • ↓ glucose utilization in cells • ↑ blood glucose levels
Protein metabolism	• ↓ protein stores in all cells except liver cells 1 catabolism of protein in cells • ↑ protein in liver • ↑amino acids in plasma

Fat metabolism	• ↑ mobilization of fatty acids • ↑ oxidation of fatty acid in cells
Anti-inflammatory effects	• Stabilizes lysosomal enzymes • ↓ migration of WBCs • ↓ phagocytosis • Suppresses the immune system

Regulation of cortisol secretion

Hypothalamic-pituitary-adrenal axis

CRH → secreted from hypothalamus → regulated by an inbuilt circadian rhythm and by additional stressors
↓
Adrenocorticotropic hormone (ACTH) → secreted from anterior pituitary under influence of corticotropin-releasing hormone (CRH)
↓
Cortisol → secreted from Adrenal under influence of ACTH
↓
Negative feedback control → Secretion of CRH and ACTH is inhibited by cortisol



Hypoadrenalism

3 patterns →

1. Glucocorticoid excess → Cushing's syndrome
2. Mineralocorticoid excess → Hyperaldosteronism
3. Androgen excess → Adrenogenital syndrome

Cushing's syndrome (Glucocorticoid or cortisol excess):

Definition →

Caused by increased cortisol due to following causes:

Etiopathogenesis

a) Exogenous corticosteroids:

- Most common cause.
- Results in decreased ACTH and bilateral adrenal atrophy.

b) Adrenal Cushing's syndrome:

- Caused by Primary adrenal adenoma (most common), hyperplasia, or carcinoma.
- Results in decreased ACTH, atrophy of uninvolved adrenal gland.

c) Pituitary Cushing disease:

- Caused by ACTH-secreting pituitary adenoma
- leads to bilateral adrenal hyperplasia.

d) Para neoplastic ACTH secretion:

- Most common cause is Carcinoid > small cell carcinoma of lung.
- Leads to bilateral adrenal hyperplasia.

Clinical Features



Clinical features

- Muscle weakness with thin extremities→ because Cortisol breaks down muscle for gluconeo genesis.
- Moon facies, buffalo bump, and truncal obesity → due to high insulin (due to high glucose) increases storage of fat.
- Abdominal striae → due to impaired synthesis of collagen with thinning of skin.
- Hypertension, Osteoporosis, and Immune suppression.



Moon facies



Abdominal striae

Hirsutism → Excess ACTH stimulates zona glomerulosa, zona fasciculata, zona reticularis producing Cortisol, androgens and probably hirsutism

igure showing marked hirsutism in a case of Cushing's syndrome

Plethora →

- Excess Cortisol causes degradation and atrophy of epidermis and underlying connective tissues
- Thinning of the skin and facial plethora

(22.07) PHEOCHROMOCYTOMA

University Exams

Short Questions

1. Discuss gross and microscopy of Pheochromocytoma.
2. Discuss Clinical features and Laboratory Diagnosis of pheochromocytoma.
3. Rule of 10's in pheochromocytoma.

Overview

- Introduction
- Rule of 10's
- Etiology
- Gross
- Microscopy
- Clinical features
- Spread
- Laboratory Diagnosis

Introduction →

- Most common tumor of adrenal medulla in Adults.
    - Derived from chromaffin cells (neural crest) of adrenal medulla → Pheochromocytoma
    - Derived from extra adrenal chromaffin cells → paraganglioma.
- Most often this tumor is benign (10% are malignant)
- Tumor cells secrete → epinephrine, norepinephrine, and dopamine→ surgically correctable Hypertension

Rule of 10's

- 10% malignant
- 10% bilateral
- 10% extra-adrenal (organs of Zuckerkandl and the carotid body)
- 10% calcify
- 10% kids

Etiology

- About 25% of individuals with pheochromocytomas and paragangliomas harbor a germline mutation in one of six known genes →

1. RET
2. NF1
3. VHL
4. SDHB (succinate dehydrogenase complex subunit genes)
5. SDHC (succinate dehydrogenase complex subunit genes)
6. SDHD (succinate dehydrogenase complex subunit genes)

Gross:



- Dark/ dusky-colored tumor
- Cut surface → reddish brown with a prominent area of fibrosis
- Areas of hemorrhage and cystic degeneration.

Microscopy



Tumor cells (Chief cells) →

- Arranged in distinctive cell balls → zellbalballen pattern (means "cell balls" in German).
- Polygonal to oval cells containing finely granular cytoplasm with salt-and-pepper nuclear chromatin

- IHC → positive for chromogranin, synaptophysin

Clinical features

- Hypertension in 90% of patients
- Paroxysmal episodes → Most Symptoms occur in "spells", relapse and remit → sudden release of catecholamines.
- Episodic hyperadrenergic symptoms (5 P's):

1. Pressure (increased BP)
2. Pain (headache)
3. Perspiration
4. Palpitations (tachycardia)
5. Pallor

Spread

- Definitive diagnosis of malignancy in pheochromocytomas is made only when they develop metastases.

1. Lymphatic spread → regional lymph nodes
2. Hematogenous spread → Distant sites → liver, lung, and bone.

Laboratory Diagnosis

- Demonstration of increased urinary excretion of free catecholamines and their metabolites → vanillylmandelic acid and metanephrines.

(22.08) MULTIPLE ENDOCRINE NEOPLASIA (MEN) SYNDROMES

University Exams

Short Questions

1. Multiple endocrine neoplasia (MEN) Syndromes.
2. MEN I syndrome.
3. Enlist four important features of MEN type I (multiple endocrine neoplasia).
4. Components of MEN type II

Overview

- Introduction
- Types →

1. MEN-I (Wermer syndrome)
2. MEN-IIA (sipples syndrome)
3. MEN-IIB

} Mutation, Components

Introduction →

Multiple adenomas and hyperplasia of different endocrine organs

Types

1. MEN-I (Wermer syndrome)
2. MEN-IIA (sipples syndrome)
3. MEN-IIB



1) MEN-I (Wermer syndrome)

Mutation → MEN 1 (or menin) gene

Components →

1. Pituitary → Prolactinoma
2. Parathyroid → parathyroid hyperplasia/ adenoma
3. Pancreas → Islet cell hyperplasia/adenoma (Pancreatic polypeptide Tumor > Gastrinoma > insulinoma > glucagonoma > VIPoma > Somatostatinoma

2) MEN-IIA (Sipples syndrome)

Mutation → RET gene

Components →

1. Thyroid → Medullary carcinoma thyroid
2. Parathyroid → Parathyroid hyperplasia/ adenoma
3. Adrenal medulla → Pheochromocytoma

3) MEN-IIB

Mutation → RET gene

Components →

1. Thyroid → Medullary thyroid carcinoma
2. Adrenal medulla → Pheochromocytoma
3. Mucosa → Mucosal neuroma

(23.01) OSTEOMYELITIS (PYOGENIC, TUBERCULAR)

University Exams

Long Questions

1. What is osteomyelitis? Describe Etiology, Pathogenesis and microscopic features of pyogenic osteomyelitis. Describe clinical course and complications of this disease.
2. Describe briefly the etiology, morphology and complications of Tuberculous osteomyelitis.

Short Questions

1. Pathogenesis of Tuberculous osteomyelitis.
2. Pathogenesis of Pyogenic osteomyelitis.
3. Morphology of pyogenic osteomyelitis.
4. Sequestrum.
5. Involucrum.
6. Complications of pyogenic osteomyelitis.
7. Pott's spine.
8. Brodie abscess
9. Sclerosing osteomyelitis of Garre

Overview

- Definition
- Classification →

1. Pyogenic Osteomyelitis
2. Tuberculous Osteomyelitis

} Introduction, Etiology, Routes, Sites, Pathogenesis, Microscopy, Clinical features, Complications

Definition →

Inflammation of the bone and marrow

Classification →

2 types

1. Pyogenic Osteomyelitis
2. Tuberculous Osteomyelitis

1) Pyogenic Osteomyelitis

Introduction →

- Also known as → Suppurative osteomyelitis
- Caused by bacterial infection

Etiology

- Most common → Staphylococcus aureus
- Other organisms → Escherichia coli, Pseudomonas, Klebsiella, Neisseria gonorrhoeae, Haemophilus influenzae and Salmonella species.

    - Neonatal period → Haemophilus influenzae and group B streptococci.
    - Genitourinary tract infections or intravenous drug abusers → Escherichia coli
    - Patients with sickle cell disease → Salmonella infection.
    - Surgery or open fracture → Mixed bacterial infections

Routes

1. Hematogenous spread:

    - Focus of infection anywhere in the body (e.g. skin pustule or infected teeth and gums, intestinal mucosa).
    - Minor injuries to mucosa (vigorous chewing of hard foods, brushing of teeth), or minor infections of skin, release these organisms into blood
    - Drug addicts (infected needles)

2. Direct implantation → by penetrating wounds, open fractures, or surgical procedures
3. Spread from adjacent site → e.g. infections of the feet may spread into the bone in diabetics.

Sites

Varies with age.

- Adult → Epiphysis and subchondral regions.
- Children and with hematogenous spread → Metaphysis of long bones
- Neonate → Metaphysis, epiphysis, or both.

Pathogenesis

Mild injury or trauma → bacteremia by organisms
↓
Infection begins in metaphyseal end of marrow cavity
↓
Pus formation → Tension in marrow cavity increased
↓
Spread of infection along marrow cavity into endosteum and cortex
↓
Reach subperiosteal space forming subperiosteal abscesses
↓
Suppuration and impaired blood supply to cortical bone
↓
Erosion, thinning and infarction necrosis of cortex → Sequestrum
↓
Cytokines released stimulates osteoclastic bone resorption and deposition of new reactive bone in the periphery→ Involucrum
↓
Pus penetrates periosteum and leads to a soft-tissue abscess may penetrate skin form a draining sinus tracts



- In **vertebral pyogenic osteomyelitis**, infection begins from **disc (discitis)** and spreads to involve the vertebral bodies



Microscopy

- In initial stages → only neutrophils seen.
- As diseases become chronic→ neutrophils admixed with chronic inflammatory cells (lymphocytes and plasma cells) seen.



Morphologic Variants of Osteomyelitis



Brodie abscess→

- Form of subacute pyogenic osteomyelitis.
- Appears as a small, solitary, intraosseous abscess localized to the metaphysis
- Surrounded by reactive bone.
- Due to inadequate treatment of infection by less virulent organisms

Sclerosing osteomyelitis of Garre →

- Extensive new bone formation
- Obscures underlying structure of bone
- Typically develops
- in the jaw.

Clinical features

- Painful and tender limb
- Fever
- Malaise
- Leucocytosis generally accompany the bony lesion.
- Radiologic examination confirms the bony destruction

Complications

1. Septicaemia
2. Pathologic fractures.
3. Development of squamous cell carcinoma
4. Secondary amyloidosis in long-standing cases.
5. Vertebral osteomyelitis may cause vertebral collapse

2) Tuberculous Osteomyelitis

Introduction →

- More destructive and resistant to control than pyogenic osteomyelitis
- Usually solitary but in patients with acquired immunodeficiency syndrome, it is multifocal.

Etiology

- Tubercle bacilli → M. tuberculosis

Routes

1. Blood borne → from a focus of active pulmonary or extrapulmonary disease.
2. Direct extension → From
    - Lung into a rib.
    - Tracheobronchial nodes into adjacent vertebrae.

Sites

1. Spine (thoracic and lumbar vertebrae)

- Known as Pott disease
- Infection breaks through intervertebral discs to involve multiple vertebrae and extends down into soft tissues forming abscesses→ cold abscess or psoas abscess

2. Knees and hips





Pathogenesis

Tuberculosis of spine (occurs in vertebral body)
↓
Pott's disease
↓
compression fractures
↓
Destruction of intervertebral discs
↓
Paraplegia
↓
Extension of caseous material along with pus from lumbar vertebrae to sheaths of psoas muscle
↓
Psoas abscess or lumbar cold abscess
↓
Burst through skin → form sinus

Microscopy

- Dead bone surrounded by tuberculous granuloma.
- Central caseation necrosis surrounded by tuberculous granulation tissue (epitheloid cells, giant cells, lymphocytes) and fragments of necrotic bone

Clinical features

- Low-grade fever with evening rise of temperature.
- Pain on motion
- Localized tenderness.
- Weight loss.

Complications

1. Destruction of vertebrae → Causes severe scoliosis or kyphosis and neurologic deficits due to spinal cord and nerve compression.
2. Psoas abscess → Infection from spine may rupture into soft tissue anteriorly and pus and necrotic debris may drain along the spinal ligaments and form a cold abscess, i.e. an abscess lacking acute inflammation.
3. Tuberculous arthritis
4. Sinus tract formation
5. Amyloidosis

	Pyogenic Osteomyelitis	Tuberculous Osteomyelitis
Introduction		
Etiology		
Routes		
Sites		
Pathogenesis		
Microscopy		
Clinical features		
Complications		

(23.02) BONE TUMOURS

University Exams

Long Questions

1. Classify bone tumors and describe etiology, types, pathogenesis, gross, microscopy and clinical features of osteosarcoma
2. Classify tumors of the bone. Describe the etiopathogenesis, gross and microscopic features of osteoclastoma (giant cell tumor of bone).
3. Classify bone tumors and describe etiology, types, pathogenesis, gross and microscopy of Ewing Sarcoma
4. A 12-year-old boy complained of pain and swelling of knee joint. On X-ray a tibial, metaphyseal lytic lesion invading the cortex and showing periosteal elevation was seen.

    a) What is your most probable diagnosis?
    b) Describe the etiology of the condition.
    c) Describe the gross and microscopy of the condition.
    d) Describe the modes of spread of the lesion.

Short Questions

1. Classification of bone tumors.
2. Gross and microscopy of Chondrosarcoma.
3. Radiological and microscopic features of osteosarcoma.
4. Radiological and microscopic features of Ewing's tumor
5. Write radiologic, gross and microscopic features of osteoclastoma.
6. Patterns of periosteal reaction
7. Surface osteosarcoma

Very Short Questions

1. Osteoid osteoma.
2. Classification of bone tumors.
3. Enumerate chondrogenic tumors/ cartilage producing tumors.
4. Mention the histological types of osteogenic sarcoma.
5. Codman's triangle and Sunray appearances on radiology
6. Soap bubble appearances on radiology
7. Onion skin appearances on radiology
8. Homer-Wright rosette.

Overview

Classification →

- Based on Histologic type
- Based on location of tumours

1. Osteoid Osteoma
2. Osteochondroma
3. Osteosarcoma
4. Chondrosarcoma
5. Giant-cell Tumor
6. Ewing Sarcoma

} Introduction, Age, Gender, Location, Sites, Gross, Microscopy, Radiological appearance, Clinical features, Spread

Classification → Based on Histologic type

Histologic type	Benign	Malignant
Osteogenic/Bone forming	Osteoma Osteoid osteoma Osteoblastoma	Osteosarcoma (primary/secondary)
Chondrogenic/Cartilaginous	Osteochondroma Chondroma Chondroblastoma	Chondrosarcoma Dedifferentiated chondrosarcoma
Hematopoietic		Myeloma Malignant lymphoma
Unknown origin	Giant-cell tumor (primitive osteoblast precursors) Unicameral cyst Aneurysmal bone cyst	Ewing sarcoma
Neuroectodermal		Chordoma

Classification → Based on location of tumours

Epiphyseal tumors	Metaphyseal lesions	Diaphyseal lesions
• **Chondroblastoma** • **Giant cell tumor (osteoclastoma)** • Clear cell chondrosarcoma	• Osteogenic sarcoma • Unicameral (simple) bone cyst • Aneurysmal bone cyst • Fibrous cortical defect • Chondrosarcoma • Osteochondroma • Enchondroma • Osteoblastoma	• Ewing sarcoma • Lymphomas • Fibrous dysplasia • Adamantinoma • Histiocytosis • Osteoid osteoma • Chondromyxoid fibroma • Fibrosarcoma • Fibrous cortical defect • Non ossifying fibroma



Remember

- Commonest bone malignancies → **Secondaries**
- Commonest primary malignant tumor → **Multiple myeloma**
- Commonest primary malignant tumor of long bones → **Osteosarcoma**
- Commonest benign tumor of bone → **Osteochondroma**
- (Osteochondroma is not true neoplasm since its growth stops with cessation of growth at epiphyseal plate)
- Commonest true benign tumor of bone → **Osteoid osteoma**
- Commonest benign tumor of hand → **Enchondroma**

1) Osteoid Osteoma

Introduction →

- Most common true benign tumor of the bone
- Slow-growing lesion

Age →

10 to 30 years of age

Gender →

M > F

Location →

diaphysis of a long bone

Sites →

Tibia

Radiological appearance →

small radiolucent central focus or nidus surrounded by dense sclerotic bone.

Gross →

Well circumscribed
Round-to-oval masses of hemorrhagic gritty tan tissue.



Microscopy →

- Composed of randomly interconnecting **trabeculae of woven bone** that are prominently rimmed by a single layer of osteoblasts
- **Stroma** surrounding neoplastic bone consists of loose connective tissue that contains many dilated and congested capillaries

Clinical features →

- Classically present with nocturnal pain → due to prostaglandin E2 production
- Relieved by aspirin

2) Osteochondromas



Introduction →

- Also known as → Osteocartilaginous Exostoses
- Commonest benign tumor of bone
- Not true neoplasm since its growth stops with cessation of growth at epiphyseal plate
- Cartilage-forming tumour
- Muatation in **EXT1 or EXT2 gene**
- May undergo transformation into chondrosarcoma

Age →

Late childhood or adolescence

Gender →

M > F

Location →

metaphyseal region (near growth plate) of long bones near knee.

Sites →

Lower femur and upper tibia (i.e. around knee) and upper humerus

Radiological appearance →

Cap is composed of benign hyaline cartilage varying in thickness



Gross →

- Broad or narrow base
- Continuous with cortical bone.
- Protrude exophytically as mushroom-shaped
- Cartilage-capped lesions enclosing well-formed cortical bone and marrow



Microscopy →

- Composed of outer cap composed of mature cartilage
- Inner mature lamellar bone and bone marrow

Clinical features →

- Remain asymptomatic and discovered as an incidental radiographic finding
- May produce obvious deformity

3) Osteosarcoma (OS)

Introduction →

- Most common primary malignant tumour of long bone (2nd only to multiple myeloma)
- Characterised by formation of osteoid or bone directly by sarcoma cells.
- Presence of osteoid is mandatory for diagnosis

Pathogenesis

Based on pathogenesis →

1. Primary OS
2. Secondary OS

1) Primary OS

- Underlying bone is unremarkable.

Mutations in tumor suppressor gene→

- RB gene mutations → associated with 1000-fold increased risk of OS and its mutation is found in about two-thirds of patients.
- Patients with Li-Fraumeni syndrome (germline TP53 mutations) → greater incidence of OS.
- Abnormalities in INK4a which codes p16 and p14
- MDM2 and CDK4: They are cell cycle regulators that inhibit functions of p53 and RB, respectively. They are overexpressed in many low-grade OS.

2) Secondary OS

Develop in association with pre-existing bone disorders like →
- Paget disease of bone.
- Radiation exposure:

    - Following therapeutic radiation for tumor, such as lymphoma.
    - Radium watch dial painters, who wetted their brushes by licking them
    - Chemotherapy: Children treated with alkylating agents

- Pre-existing benign bone lesions: e.g. fibrous dysplasia, osteomyelitis, and bone marrow infarcts. Trauma may call attention to an existing OS rather than causing the tumor



Central / Medullary/ Conventional/ Classical OS

Introduction →
More common and classic type

Age →
Bimodal age distribution.

- Primary OS → between 10- and 20-years age.
- Secondary OS → in elderly after 40 years of age.

Gender →
M > F

Location →
Metaphyseal region of the long bones of the extremities.

Sites →
in descending order of frequency →

- Lower end of femur and upper end of tibia (i.e. around knee joint about 60%)
- Upper end of humerus (10%)
- Pelvis and upper end of femur (i.e. around hip joint about 15%)
- In jaw bones, vertebrae and skull.

Radiological appearance →

- When the malignant tumor extends into soft tissue, parallel lines of mineral deposition periosteal region gives an appearance of rays of sun → "sunburst" appearance
- Tumor infiltrates cortex → lifts periosteum→ produces a space between cortex and lifted ends of periosteum → Radiologically appears as triangular shadow → contains reactive new bone which is arranged perpendicular to bone surface → Codman triangle (characteristic but not diagnostic of this tumor).

AP (anterioposterior) view Lateral view

Tumour arises centrally in metaphysis
↓
Extends longitudinally for variable distance into the medullary cavity
↓
expands laterally on either side breaking through the cortex
↓
Lifting periosteum
↓
If periosteum is breached → tumour grows relentlessly into surrounding soft tissues.



Patterns of periosteal reaction





Gross →

- **Grey-white, bulky mass** at metaphyseal end of a long bone of extremity.
- C/S → grey-white, hard, gritty with areas of **haemorrhages and necrotic bone.**

Microscopy →

1. Sarcoma cells

   - Tumour cells are undifferentiated mesenchymal stromal cells
   - show marked pleomorphism and polymorphism i.e. variation in size as well as shape.
   - Show hyperchromatism
   - Atypical mitoses.
   - IIHC → sarcoma cells of osteosarcoma express vimentin, osteocalcin, osteonectin and type I collagen



2. Osteogenesis

   - Sarcoma cells form osteoid matrix and bone directly (lacelike pattern of neoplastic bone)
   - Found interspersed in the areas of tumour cells.

Clinical features→

- Painful, progressively enlarging masses around knee or other involved site.
- Involved area → swollen and tender
- Function of adjacent joint → reduced.
- Serum alkaline phosphatase→ increased.

Spread →

1. Local →

   - Invasion of adjacent cortex and destroys nearby cortex.
   - Elevation or perforation of periosteum by tumor tissue.
   - Spread along the medullary (marrow) cavity.
   - Extention into epiphysis → infiltrate epiphyseal plate and may even involve the joint space.
   - Extension into the soft tissues and may involve skin

2. Blood → to lungs (most common), bones, pleura, brain, and heart.



Surface OS

- About 5% of osteosarcomas occur on surface of bone
- **Slow-growing tumours** compared to medullary osteosarcomas.
- Surface osteosarcoma includes 2 variants:

  1. Parosteal OS
  2. Periosteal OS

1) Parosteal or juxtacortical OS

Introduction →

- Uncommon form of slow-growing OS
- Arises on the surface of bone and grows along external surface of the bone, both linearly and longitudinally, without involving medullary cavity.

Age → older age group (3rd to 4th decade)

Gender → No sex predilection

Location → metaphysis of long bones

Site → most frequently lower end of femur and upper end of humerus.

Clinical features → slow growing, Excellent prognosis

2) Periosteal OS
Introduction →
- Arises between cortex and overlying periosteum
- High-grade OS, and characteristically predominant cartilage component.

Age → young adults (average age 25 years)

Gender → No sex predilection

Location → diaphysis

Site → Tibia or femur

Clinical features → Slow growing

	Feature	Central (Medullary)	Surface (Parosteal And Periosteal)
1	Age	10-20 years	Older patients
2	Sex	More common in male	No sex predilection
3	Anatomic site	Metaphysis	Metaphysis/diaphysis
4	Location	Femur (lower end), tibia (upper end), humerus (upper end), around hip	Femur (lower-end), humerus (upper end)
5	Pathogenesis	Primary: genetic factors (mutations in Rb gene, p53, MDM2) Secondary: Paget's disease, fibrous dysplasia	Parosteal: Arises outer to cortex Periosteal: Arises between cortex and Periosteum
6	Behaviour	Highly malignant	Slow growing
7	G/A	Bulky, necrotic, forms Codman's triangle	Smaller, well-formed bone present
8	M/E	1. Sarcomas cells: Polymorphic and pleomorphic 2. Osteoid formation	1. Parosteal: Fibrous stromal cells with subtle atypia 2. Periosteal: High grade 3. Both form bony trabeculae
9	Histologic types	Telangiectatic, small cell, fibrohistiocystic, well differentiated, anaplastic	Parosteal (Juxta cortical), periosteal
10	Spread and prognosis	Haematogenous spread, prognosis poor	Recurrences common, may metastasise, prognosis generally good, better for parosteal than periosteal

4) Chondrosarcoma

Introduction →

- Malignant tumour of chondroblasts
- Characterized by production of neoplastic cartilage.

Age →

4th to 6th decades

Gender →

M > F

Location →

arise in central medullary cavity

Sites →

- Pelvic bones
- Shoulder
- Ribs
- Long bones

Radiological appearance →

- Shows a tumor in medullary cavity with poorly defined borders.
- Shaft is thickened and tumor perforates cortex.
- Calcified matrix appears as foci of flocculent densities with ring-like ossifications.
- Popcorn calcification, flocculent densities with ring-like ossifications.



Gross →

- Large bulky tumors
- Consists of nodules of light blue with translucent glistening areas (represents cartilage).
- C/S →
    - Mucoid/myxoid feel
    - Spotty calcifications
    - Central area of necrosis may produce cystic spaces.





Microscopy →

- Consists of malignant cartilage cells in various stages of maturity.
- Graded from 1 to 3, depending on degree of cellularity, cytological atypia and mitotic activity.

Grade 1:
- Mild hypercellularity.
- Chondrocytes show plump vesicular nuclei with small nucleoli.
- Binucleate cells are few in number.
- Mitotic figures are rare.
- Matrix: Portions of matrix mineralize, and cartilage undergo endochondral ossification.

Grade 2:
- Moderate cellularity and more cellular than grade 1.
- Chondrocytes show more pronounced nuclear changes.

Grade 3:
- Marked hypercellularity.
- Chondrocytes show extreme pleomorphism
- Bizarre tumor giant cells
- Mitoses frequent.

Clinical features →

- Painful, progressively enlarging masses.
- Biologic behavior of tumor depends on the histological grade.

Spread →

- Local → Locally aggressive tumor→ infiltrate marrow space and may grow with broad pushing fronts into surrounding soft tissue.
-
- Blood → Lungs and skeleton.

5) Giant-cell Tumor

Introduction →

- Also known as osteoclastoma
- Cell of origin: Primitive stromal cells.
- characterized by multinucleated osteoclast-type giant cells randomly and uniformly distributed in a background of mononuclear cells.
- Locally aggressive
- Potentially malignant neoplasm.

Age →

20 to 40 years

Gender →

F > M

Location →

Involve epiphyses but may extend into the metaphysis of a long bone



Site →

- Knee area: Distal femur and proximal tibia.
- Lower end of radius, humerus and fibula.



Soap-bubble appearance

Radiological appearance →

- Produces a lytic lesion and grows slowly, expands bone and destroys overlying cortex.
- Tumor produces a bulging soft tissue mass surrounded by a thin, shell of reactive bone due to periosteal reaction → on radiology has multiloculated or soap-bubble appearance



Gross →

- Large tumor, clearly circumscribed and frequently undergo cystic change.
- C/S →
    - Soft and red-brown without bone or calcification
    - Numerous hemorrhagic areas give an appearance of a sponge full of blood → **honey-combed appearance**



Microscopy →

2 types of cells →

1. Neoplastic mononuclear Stromal cells→
    - Uniform oval and plump
    - Large nuclei with prominent nucleoli
    - Scanty cytoplasm
    - Tumor mostly consists of these neoplastic cells.
    - Mitotic activity is common
    - Diagnosis of malignancy depends upon the morphology of the mononuclear cells rather than that of the multinucleated giant cells.
2. Multinucleated (osteoclastic) giant cells →

- Background shows numerous large osteoclast-type giant cells throughout richly vascularized stroma.
- Have 100 or more nuclei

Clinical features →

- Pain
- Arthritis-like symptoms.
- Microfractures and pathologic fractures (due to thinning of cortex)

Spread →

1. Local → Aggressive tumors may penetrate cortex and periosteum, and involve joint capsule and synovial membrane.
2. Blood → Common site is lung

6) Ewing Sarcoma

Introduction →

- Also known as Primitive Neuroectodermal Tumor
- Primary malignant small round-cell tumors of bone and soft tissue.
- Constitute 6–10% of primary malignant bone tumors.
- Highly aggressive

Age →

- 10 to 15 years of age
- In children → it is second most common group of bone sarcomas (first is osteosarcoma)

Gender →

M > F

Pathogenesis →

Location of genes →
- EWS on chromosome 22q12
- FL1 gene on 11q24
- ERG gene on 21q22.

Ewing sarcoma: Translocations →
- t (11; 22) (q24; q12)
- t (21;22) (q22; q12).

Translocation of EWS gene on chromosome 22 (22q12) with
a gene encoding an ETS family transcription factor (FLI-1 or ERG gene)
↓
Fusion of EWS1 gene to FLI-1 gene (more common)
↓
Fusion protein → EWS/FLI-1
↓
Produces chimeric/aberrant transcription factors
↓
Activation of C-myc promoter
↓
Abnormal cell proliferation and survival



- A less common translocation produces EWS/ERG fusion gene → produces a variant of EWS with worse prognosis

Location →

Arise in diaphysis of medullary cavity of long bones.

Site →

- Long bones→ Humerus, tibia and femur.
- Flat bones→ Pelvis.

Radiological appearance →

- Produces destructive lytic lesion in which border between normal bone and the tumor is indistinct.
- Tumor may extend into surrounding soft tissues.
- Characteristic circumferential, discontinuous layers of periosteal new bone formation produces → onion-skin pattern

Gross →

- C/S → Soft, grayish white
- Show areas of hemorrhage and necrosis.

Microscopy →

Tumor cells →

- Uniform, small and round, Twice the size of a lymphocyte.
- Cytoplasm → Scant and clear because it is rich in glycogen
- Round nuclei having 'salt and pepper' chromatin

Arrangement →

- Closely packed sheets of cells.



- Homer–Wright rosettes → tumor cells arranged in a circle about a central fibrillary space.

Interstitial stroma →

- Fibrous strands separate sheets of tumor cells into irregular nests
- Areas of necrosis

IHC →
- FLI-1
- CD99

Clinical Features →

1. Painful enlarging masses
2. Affected site → tender, warm, and swollen.
3. Systemic features → Fever, weakness, and bone pain.
4. Hematological findings→ Anemia, leukocytosis, and raised erythrocyte sedimentation rate.

Spread →

1. Local → spread from medullary cavity into cortex, periosteum and surrounding soft tissue.
2. Blood → Lungs, brain and other bones (skull).

	Osteoid Osteoma	Osteochondroma	Osteosarcoma	Chondrosarcoma	Giant-cell Tumor	Ewing Sarcoma
Introduction						
Age						
Gender						
Location						
Sites						
Gross						
Microscopy						
Radiological appearance						
Clinical features						
Spread						

(23.03) GOUT

University Exams

Short Questions

1. Gouty arthritis.
2. Pathogenesis of gouty arthritis.
3. Etiology and Pathogenesis of gout.
4. Enlist four causes of gout.
5. Microscopic features of gout.
6. Gouty tophi.

Overview

- Definition
- Classification
- Etiology
- Pathogenesis
- Gouty Arthritis →

    1. Acute Arthritis
    2. Chronic Arthritis

- Morphology
- Clinical Course

Definition

- Gout is a heterogeneous group of diseases characterized by hyperuricemia

Gout
↓
Plasma urate level > 6.8 mg/dL
↓
Hyperuricemia
↓
Urate crystal deposition in joints and kidneys

Classification

Primary gout (90%) → Hyperuricemia occurs without any other disease.
• Increased production of uric acid – Increased dietary intake of purine – Due to inborn errors of purine metabolism • Decreased excretion of uric acid with normal production
Secondary gout (10%) → Hyperuricemia occurs in association with another illness
• Increased production of uric acid with increased urinary excretion – Increased catabolism of nucleic acid due to increased cell turnover (e.g. leukemias, cancers, and during tumor lysis induced by chemotherapy) • Decreased excretion of uric acid with normal production – Chronic renal disease

Etiology

- Uric acid end product of purine metabolism, which is eliminated only in the urine.
- Hyperuricemia is necessary, but not sufficient for development of gout.
- Several factors convert asymptomatic hyperuricemia into primary gout→

  - Age: usually presents after 30 years of age.
  - Duration of hyperuricemia: not less than 20–30 years.
  - Heavy alcohol intake.
  - Obesity.
  - Drugs (e.g thiazides) which reduce excretion of urate.
  - Lead toxicity.

Pathogenesis

Hyperuricemia
↓
Monosodium urate (MSU) crystals formed
↓
Monosodium urate is less soluble in synovial fluid compared to plasma and less soluble at low temperature
↓
Become supersaturated in peripheral joints (ankles and toes→ low temperature)
↓
Precipitation of urate crystals in synovium and joint cartilage
↓
Gouty Arthritis



Gouty Arthritis

1. Acute Arthritis
2. Chronic Arthritis

1) Acute Arthritis

Trauma
↓
Precipitated crystals released into synovial fluid
↓
MSU crystals phagocytosed by macrophages
↓
Release many cytokines (e.g. IL-1**β**, IL-18).
↓
IL-1**β** stimulates expression of adhesion molecules → Accumulation of neutrophils at site of acute inflammation
↓
Neutrophils release leukotriene B4 and lysosomal enzymes
↓
Aggravate inflammation
↓
Acute arthritis



2) Chronic Arthritis

Repeated attacks of acute arthritis
↓
Tophi formation in synovial membranes and periarticular tissue
↓
Chronic arthritis

Morphology

Acute arthritis →

- Acute inflammation: It is characterized by edema, congestion and dense infiltration of synovium by neutrophils.
- Few lymphocytes, plasma cells, and macrophages seen.
- Monosodium urate (MSU) crystals →
    - Long, slender, and needle shaped.
    - Found in cytoplasm of neutrophils and are arranged in small clusters in the synovium.
    - As acute attack subside → resolubilized.



Chronic tophaceous arthritis →

- Urates encrust articular surfaces.
- Synovium shows fibrosis, thickening, and infiltration by inflammatory cells.
- Forms a pannus and destroy cartilage, and juxta-articular bone erosions.
- Tophi →
    - Lesions consisting of large aggregates of urate crystals, surrounded by inflammatory reaction
    - Pathognomonic of gout.
    - Sites: articular cartilage of joints and in periarticular ligaments, tendons, and soft tissues, Achilles tendons, and earlobes.
    - Microscopy: consist of aggregations of urate crystals surrounded by macrophages, lymphocytes, and foreign body giant cells.



Clinical Course

4 Stages →

1. Asymptomatic hyperuricemia: Appears at puberty in males and after menopause in females.
2. Acute gouty arthritis: sudden excruciating joint pain with tenderness.
3. Intercritical gout: asymptomatic interval between first acute attack and subsequent attacks.

4. Chronic tophaceous gout: After several years of acute gouty arthritis → chronic tophaceous gout develop → severe crippling disease.

(24.01) MENINGITIS

University Exams

Long questions

1. Describe CSF findings in case of meningitis.
2. Etiology of Lymphocytic meningitis. Write its CSF findings.
3. CSF finding in pyogenic and tubercular meningitis.
4. Enumerate causative organisms of acute pyogenic meningitis. Describe C.S.F. picture in pyogenic meningitis.
5. List of indications for CSF examination. Give CSF findings in acute Pyogenic meningitis and tuberculous meningitis.
6. Discuss routine, microscopic examination of CSF. and its utility in differential diagnosis of various types of meningitis.
7. Classify meningitis. Describe morphological features and differential diagnosis of Viral meningitis.

Short questions

1. C.S.F. in Viral meningitis.
2. List the CSF findings in Tuberculous meningitis.
3. Etiology and CSF findings in acute pyogenic meningitis.
4. Etiology and CSF findings in Chronic meningitis.
5. Tabulate differences in CSF examination in pyogenic, tuberculous and viral meningitis.

Overview

- Meninges of CNS
- Meningitis
    - Definition
    - Routes of infection
    - Types →

a) Acute Pyogenic Meningitis b) Acute Viral Meningitis c) Chronic Meningitis	} Etiopathogenesis Clinical features CSF Findings

Meninges of CNS

- Dura mater → tough fibrous covering closely attached to the skull
- Pia mater → closely applied to brain
- Arachnoid mater → lies between pia mater and dura mater

✓ Space between bone and dura meter → Epidural space
✓ Space between durameter and arachnoid meter → Subdural space
✓ Space between pia mater and arachnoid mater → subarachnoid space → contains the CSF.



Meningitis

Definition → Inflammatory involvement of meninges.



Routes of infection

1. Blood stream → Most common
2. Adjacent focus of infection
3. By iatrogenic infection → introduction of microorganisms at operation or during lumbar puncture





Types

1. Acute Pyogenic (bacterial) Meningitis
2. Acute Lymphocytic (Viral, Aseptic) Meningitis
3. Chronic (Tuberculous and Cryptococcal) Meningitis

Features	Normal	Acute Pyogenic (Bacterial) meningitis	Acute Lymphocytic (Viral Meningitis)	Chronic (Tuberculous) Meningitis
Naked eye appearance	Clear and colorless	Cloudy or frankly purulent	Clear or slightly turbid	Cob web (fibrin coagulum on standing)
CSF pressure	60-150 mm water	Elevated (Above 180 mm water)	Elevated (Above 250 mm water)	Elevated (Above 300 mm water)
Cells	0-4 lymphocytes/ul	100-10,000 Neutrophils/ul	10-100 lymphoctes/ul	100-1000 lymphoctes/ul
Glucose	50-80 mg/dl	Markedly reduced	Normal	Reduced
Protein	15 - 45 mg/dl	Markedly Elevated	Normal	Elevated
Bacteriology	Sterile	Causative Organisms present	Sterile	Tubercle bacilli Present

(1) Acute Pyogenic (bacterial) Meningitis

Etiopathogenesis →

- Neonates: Group B streptococcus (Str. agalactiae) > E. coli
- 2 months to 3 years: Pneumococci > Meningococci > H. influenzae
- 3 years to 20 years: Meningococci
- 20 years: Pneumococci

Clinical Features

Medical emergency →

- Fever
- Severe headache
- Stiffness of the neck on forward bending.
- Vomiting
- Coma
- Occasionally convulsions

Brudzinsk's Sign

Flexion of the Neck causes involuntary flexion of the Knee and Hip Joints.

CSF findings



1. Naked eye appearance → cloudy or frankly purulent CSF
2. Elevated CSF pressure (above 180 mm water)
3. Neutrophilic leucocytosis (between 100 - 10,000/µl).
4. Raised CSF protein level (higher than 50 mg/dl).
5. Decreased CSF sugar concentration (lower than 40 mg/dl).
6. Gram's stain or CSF culture reveals causative organism.

(2) Acute Lymphocytic (Viral, Aseptic) Meningitis

Etiopathogenesis →

- Enteroviruses (Most common)
- Mumps
- ECHO viruses
- Coxsackie virus
- Epstein-Barr virus
- Herpes simplex virus-2
- Arthropode-borne viruses
- HIV

Clinical Features

- Same as in bacterial meningitis
- Benign and self-limiting clinical course of short duration
- Complete recovery occurs

CSF findings



1. Naked eye appearance → clear or slightly turbid
2. CSF pressure increased (above 250 mm water)
3. Lymphocytosis in CSF (10-100 cells/µl).
4. CSF protein →normal
5. CSF sugar concentration → normal.
6. CSF bacteriologically sterile.

(3) Chronic Meningitis

Etiopathogenesis →

1. Tuberculous meningitis
2. Cryptococcal meningitis → develops in immunocompromised persons (AIDS)

Clinical Features

- Headache
- Confusion
- Malaise
- Vomiting

CSF findings

1. Naked eye appearance → fibrin web on standing (cob web appearance)
2. Raised CSF pressure (above 300 mm water).
3. Mononuclear leucocytosis (100-1000 cells/µl).
4. Raised protein content.
5. Lowered glucose concentration
6. Special stains→

   - **ZN staining** → tuberculous meningitis.
   - **India ink preparation** → cryptococcal meningitis.



Features	Acute Pyogenic (Bacterial) meningitis	Acute Lymphocytic (Viral Meningitis)	Chronic (Tuberculous) Meningitis
Etiology			
Clinical features			
Naked eye appearance			
CSF pressure			
Cells			
Glucose			
Protein			
Bacteriology			

(24.02) TUMOURS OF CNS

University Exams

Short questions

1. Classification of brain tumors.
2. Describe gross & microscopic features of Astrocytoma.
3. Morphology of glioblastoma multiforme.
4. Describe gross & microscopic features of Oligodendroglioma.
5. Describe gross & microscopic features of Medulloblastoma.
6. Histological types of gliomas.
7. Describe gross & microscopic features of meningioma.
8. Describe gross & microscopic features of Schwannoma (Neurilemoma)
9. Describe gross & microscopic features of Ependymoma

Overview

- Classification of Tumours of CNS

• Pilocytic astrocytoma • Glioblastoma Multiforme • Oligodendroglioma • Ependymoma • Meningioma • Medulloblastoma • Schwannoma	Introduction Site Gross Microscopy

Classification of Tumours of CNS

Primary tumours	Metastatic tumours
Originate in brain or spinal cord	Spread to brain from another primary site of cancer

Metastatic CNS tumours

- **Metastasis to brain →** most common brain tumor
- Most common primary malignancy metastatizing to brain → **lung carcinoma > breast > skin (melanoma) > kidney (genitourinary) and GIT cancers.**



Important Primary brain tumours

- Gliomas →
    - Astrocytoma → Pilocytic astrocytoma, Glioblastoma Multiforme
    - Oligodendroglioma
    - Ependymoma
- Meningiomas
- Medulloblastoma
- Nerve Sheath Tumours (Schwannomas)

Gliomas

Tumours arising from neuroglia

Glioma types→
1. Astrocytomas → From astrocytes
2. Oligodendroglioma → From oligodendrocytes
3. Ependymoma → From ependyma

Astrocytomas

Most common type of gliomas

WHO classification →
1. WHO Grade I (Pilocytic) Astrocytoma → Most common in children and good prognosis
2. WHO Grade II (Fibrillary) Astrocytoma
3. WHO Grade III (Anaplastic) Astrocytoma
4. WHO Grade IV Astrocytoma (Glioblastoma Multiforme)→ Most common in adult and Poor prognosis

WHO Grade I Astrocytoma (Pilocytic Astrocytoma)

Introduction →



- Occur in Children
- Surgical treatment is curative in 80 to 100% of cases
- Good prognosis

Site →

Cerebellum and brainstem



Gross →

- Usually well circumscribed
- Commonly cystic (myxoid or mucoid appearance)

Microscopy →

- Rosenthal fibers→ intracytoplasmic inclusions which are condensed mass of glial fibrillary acidic protein (GFAP)

WHO Grade IV Astrocytoma (Glioblastoma Multiforme)



Introduction →

6th decade of life

Site →

Cerebral hemispheres (temoral and frontal lobes)

Gross →

- Variegated appearance with haemorrhages and necrosis.
- Also known as butterfly tumor since it crosses midline.

Microscopy →

- Tumour cells are poorly-differentiated round cells, pleomorphic cells and giant cells.
- Mitoses are frequent
- Pseudopalisading necrosis → serpentine necrotic areas with tumor cells at edges of necrotic border



Oligodendroglioma

Site →

- Cerebral hemisphere white matter (80 to 90% supratentorial)

Gross →

- Well-circumscribed, grey-white gelatinous mass
- Foci of haemorrhages and calcification

Microscopy →

- Tumour cells have round to oval nuclei surrounded by a clear halo of cytoplasm → Fried Egg appearance
- Endothelial cell hyperplasia
- Foci of calcification



Ependymoma

Site →

- In first two decades of life → **fourth ventricle**
- In adults → **spinal cord** (associated with neurofibromatosis type 2)

Gross→

- Well-demarcated tumour

Microscopy →

- Tumour is composed of uniform epithelial (ependymal) cells forming rosettes and perivascular pseudorosettes.
- Blepharoplasts representing basal bodies of cilia seen in cytoplasm of tumour cells



Meningioma

Introduction →

- 2nd to 6th decades of life
- Female preponderance.

Site →

- Arise from cap cell layer of arachnoid

Gross →

- Well-circumscribed mass of varying size (1-10 cm in diameter)
- Solitary





Microscopy →

- Conspicuous whorled pattern of tumour cells
- Some of the whorls contain psammoma bodies → calcification of central core of whorls

Medulloblastoma



Introduction →

- Most common primary malignant brain tumour in childhood.

Site →

- Occur exclusively in the cerebellum.

Gross→

- Protrudes into the fourth ventricle as a soft, grey-white mass or invades the surface of the cerebellum



Microscopy →

a) Small round blue tumor cells
b) Homer-Wright pseudo-rosettes→ groups of tumor cells arranged in a circle around a fibrillary center

Nerve Sheath Tumours

1. Schwannoma (neurilemmoma)
2. Neurofibroma

Schwannomas (Neurilemmomas)

Introduction →

- Solitary nodule
- Multiple schwannomas occur in von Recklinghausen's disease
- Association with Neurofibromatosis type 2
- Invariably benign

Site →

Arise from cranial and spinal nerve roots.

- Most common → Acoustic schwannoma → schwannoma of 8th nerve

Gross →

- Encapsulated, solid tumour
- Produces eccentric enlargement of nerve root
- Does not infiltrate nerve



Microscopy →

- Two areas seen histologically →

  a) Antoni A pattern → dense and compact cellular areas show palisaded nuclei called Verocay bodies
  b) Antoni B pattern → Loose acellular areas

	Pilocytic astrocytoma	Glioblastoma Multiforme	Oligodendroglioma	Ependymoma
MC site	Cerebellum	Cerebral hemisphere	Cerebral Cortex (white matter)	4th Ventricle
Gross	• Well circumscribed • Cystic	Butterfly tumor	Gelatinous mass	Well demarcated
Microscopy	Rosenthal fibres	• Pleomorphic • Pseudopalisading necrosis • Mitosis	• Fried egg cell • Calcification	Rosettes, Pseudorosettes Blepharoplasts

	Meningioma	Medulloblastoma	Schwannomas
MC site	• Arise from Arachnoid • Lateral Cerebral Convexities	Cerebellum	8th nerve (Acoustic)
Gross	• Well circumscribed	Grey- white mass	• Encapsulated • Eccentric • Does not infiltrate nerve
Microscopy	• Whorled pattern • Psammoma bodies	– Home Wright Rosettes	• Antoni A → Cellular areas (Verocay body) • Antoni B → Loose areas

(25.01) TUMORS OF SALIVARY GLANDS

University Exams

Short Questions

1. Classify Salivary gland tumours
2. Name 3 benign Salivary gland tumours. Write Etiology, Gross, Microscopy and clinical features of Pleomorphic Adenoma
3. Discuss Etiology, Gross, Microscopy and clinical features of Warthin's Tumour
4. Write Gross and Microscopy of Mucoepidermoid Carcinoma

Overview

- Classification

1. Pleomorphic Adenoma
2. Warthin's Tumour
3. Mucoepidermoid Carcinoma

} Introduction, Age, Gender, Etiology, Site, Gross, Microscopy, Clinical features

Classification

Benign	Malignant
Pleomorphic adenoma (mixed tumor)	Mucoepidermoid carcinoma
Warthin tumor	Adenocarcinoma
Oncocytoma	Acinic cell carcinoma
Basal cell adenoma	Adenoid cystic carcinoma
	Malignant mixed tumor

(1) Pleomorphic Adenoma

Introduction →

- Also known as → Mixed Salivary Tumour
- Most common benign tumor of the salivary glands
- Pleomorphic → variable cell type seen in this lesion

Age →

3rd to 5th decades of life

Gender →

F > M

Etiology →

Exposure to radiation increases the risk

Site

Major salivary gland → superficial lobe of the parotid (60%) > submandibular glands > sublingual gland > Minor salivary gland

Gross

- Solitary
- 2-5 cm in diameter
- Circumscribed
- Pseudoencapsulated
- Multilobulated, firm mass,
- Bosselated surface.
- C/S → lobules of grey-white circumscribed tumour having semitranslucent parenchyma

Microscopy



- Pleomorphic or 'mixed' appearance → epithelial elements present in a stromal matrix of mucoid, myxoid and chondroid tissue
- Epithelial element → ducts, acini, tubules, sheets and strands of cuboidal and myoepithelial cells.
- Mesenchymal elements → pseudocartilage which is matrix of myxoid, chondroid and mucoid material.

Clinical features



1. Often located below and in front of ear
2. Painless
3. Slow-growing
4. Rarely, a carcinoma may arise in pleomorphic adenoma → carcinoma ex pleomorphic adenoma or a malignant mixed tumor.
5. Pleomorphic adenoma is recurrent

Main factors responsible for recurrance →

- Incomplete surgical removal due to proximity to facial nerve
- Multiple foci of tumour
- Pseudoencapsulation
- Implantation in surgical field

Warthin's Tumour

Introduction →

- Also known as →
  - Papillary Cystadenoma Lymphomatosum
  - Adenolymphoma
- Second most common salivary gland neoplasm.
- 8% of all parotid neoplasms

Age →

4th to 7th decades of life

Gender →

M > F

Etiology →

Smoking increases the risk

Site

Almost exclusively arises in the parotid gland

Gross

- Encapsulated
- Round or oval with smooth surface
- C/S → characteristic slit-like or cystic spaces, containing milky fluid and having papillary projections.



WARTHIN'S TUMOR
Gross description

Microscopy

Tumour shows 2 components:

1. Epithelial parenchyma →
   - Glandular and cystic structures having papillary arrangement
   - Lined by characteristic eosinophilic epithelium.
2. Lymphoid stroma →
   - Present under the epithelium in form of prominent lymphoid tissue
   - Often with germinal centres.



Clinical features

1. Often located below and in front of ear
2. Painless
3. Slow-growing

Mucoepidermoid Carcinoma

Introduction →

- Most common Primary malignant tumor of salivary gland.
- Constitute about 15% of all salivary gland tumors.

Age →

children

Gender →

M = F

Etiology

Balanced (11;19) (q21; p13) chromosomal translocation.
↓
Creates a fusion gene composed of portions of the CRTC1 and MAML2 genes
↓
Activation of the Notch pathway
↓
Uncontrolled mitosis
↓
Mucoepidermoid Carcinoma

Site

- Most common → Parotids (60–70%)
- Account for a major fraction of minor salivary gland tumors.

Gross



- Size → as large as 8 cm in diameter.
- Appear circumscribed, but do not show well-defined capsules
- Infiltrative at margins.

- C/S →
    - Pale and gray-white
    - Contain small, mucin-containing cysts.

Microscopy



Tumor cells arranged in cords, sheets, or cystic structures

Composed of variable mixtures of →
1. Squamous cells
2. Mucus-secreting cells
3. Intermediate cells

Clinical features

- Usually Painless
- Sometimes painful
- Otorrhea
- Dysphagia
- Trismus

	Pleomorphic Adenoma	Warthin's Tumour	Mucoepidermoid Carcinoma
Introduction			
Age			
Gender			
Etiology			
Site			
Gross			
Microscopy			
Clinical features			

(26.01) TUMOURS OF SKIN

University Exams

Short Questions

1. Discuss briefly about precancerous lesions of skin.
2. Write Predisposing factors, Pathogenesis, Gross and microscopy of Basal cell carcinoma (rodent ulcer)
3. Discuss Predisposing factors, Pathogenesis, growth patterns and morphology of Malignant melanoma
4. Etiology of skin cancer
5. Etiology, goss and microscopy of squamous cell carcinoma of skin.
6. Whal is ABCD of melanoma?
7. Sites of malignant melanoma.
8. Gross and Microscopy of malignant melanoma

Overview

1. Squamous Cell Carcinoma
2. Basal Cell Carcinoma
3. Melanoma

} Introduction, Predisposing factors, Pathogenesis, Site, Gross, Microscopy, Clinical features, Prognosis

Squamous Cell Carcinoma

Introduction →

Second most common tumor arising on sun-exposed sites

Predisposing conditions

1. Xeroderma pigmentosum
2. Epidermodysplasia verruciformis induced by HPV
3. Solar keratosis
4. Chronic inflammatory conditions such as chronic ulcers and draining osteomyelitis
5. Old burn scars (Marjolin's ulcers)
6. Chemical burns
7. Psoriasis
8. HIV infection
9. Ionising radiation
10. Industrial carcinogens (coal tars, oils etc)
11. In the case of cancer of oral cavity, chewing betel nuts and tobacco.

Pathogenesis

Predisposing conditions → Mainly Prolonged sun exposure and immunosuppression
↓
p53 mutation
↓
DNA damage
↓
dysregulation of signalling pathway
↓
Uncontrolled mitosis
↓
Squamous Cell Carcinoma

Site

Squamous cell carcinoma may arise on any part of the skin and mucous membranes lined by squamous epithelium.

Most common →
- Face
- Pinna of ears
- Back of hands
- Mucocutaneous junctions such as on lips, anal canal and glans penis.

Gross

2 patterns:
1. Ulcerated growth with elevated and indurated margin → More common
2. Fungating or polypoid verrucous lesion→ Less common



Microscopy

- Tumors cells arrangement → Irregular downward proliferation of epidermal cells into dermis.
- Depending upon grade of malignancy→ masses of epidermal cells show atypical features →
  - Variation in cell size and shape
  - Nuclear hyperchromatism
  - Absence of intercellular bridges
  - Individual cell keratinisation
  - Occurrence of atypical mitotic figures





1. Well differentiated squamous cell carcinomas →
   - Composed of polygonal squamous tumor cells arranged in lobules
   - Produce large amounts of keratin
   - Some of this keratin form squamous pearls → Centres contain laminated, keratin material



2. Moderately differentiated squamous cell carcinoma →
   - Consist of anaplastic squamous cells
   - Single-cell keratinization (dyskeratosis)



3. Poorly differentiated squamous cell carcinomas →
   - Consist of highly anaplastic cells.
   - Fewer or no Keratin pearls

Clinical features

- Appear as sharply defined, red, scaling plaques.

Prognosis

- Overall prognosis → excellent.
- Superficial invasive tumour may metastasise locally.
- Prognosis of deeply invading tumour depends upon TNM stage

Basal Cell Carcinoma

Introduction →

- Also known as → Rodent Ulcer
- Locally invasive
- Slow-growing tumour
- Rarely metastasises

Predisposing factors

1. Light-skinned people who have little melanin.
2. Prolonged exposure to strong sunlight like in those living in Australia and New Zealand.
3. Inherited defect in DNA repair mechanism in xeroderma pigmentosum.
4. Nevoid basal cell carcinoma syndrome→ autosomal dominant condition in which multiple basal cell carcinomas appear at a young age (under 20 years).

Pathogenesis

Basal Cell Carcinoma
↓
Primitive adnexal carcinomas
↓
Indubitable hair follicle derivation

Site

- Exclusively on hairy skin
- Most common location (90%) → Face → above a line from lobe of ear to corner of mouth



Gross

- Nodulo-ulcerative → small nodule undergoes central ulceration with pearly, rolled margins.
- Tumour enlarges in size by burrowing and by destroying tissues locally like a rodent → 'rodent ulcer'

Microscopy

Resemble normal basal cell layer of epidermis and show peripheral palisading.

Tumors cells arrangement →
- In nests
- Attached to epidermis and protrude into subjacent papillary dermis.
- At periphery of each nest columnar cells arranged radially with their long axes in parallel alignment → peripheral palisading.


Tumor cells with peripheral palisading

Tumor cells→
- Resemble normal basal cell layer of epidermis
- Deeply basophilic epithelial cells
- Have a large, oval, or elongated nucleus
- Narrow rim of cytoplasm

Clinical features

- Erodes underlying tissue like a rodent
- Appear as pearly papules often containing prominent, dilated subepidermal blood vessels.
- Locally invasive
- Metastasis very rare.

Prognosis

Overall prognosis → excellent for cases that have not spread to other sites

Malignant Melanoma

Introduction →

- Most rapidly spreading malignant tumour of skin

Risk factors

- Persistent change in appearance of a mole.
- Presence of pre-existing naevus
- Family history of melanoma in a patient of atypical mole.
- Higher age of patient.
- More than 50 moles 2 mm or more in diameter.
- Molecular studies in familial and hereditary cases revealed germline mutation in CDKN2A gene which encodes for cyclin-dependent kinase inhibitor, mutational loss of PTEN gene and mutation in several other tumour suppressor genes but not p53.

Pathogenesis

Cell of origin: Melanocytes

Sun Exposure →
- Melanomas develop on sun-exposed surfaces → upper back in men and the back and legs in women.
- Lightly pigmented individuals are at greater risk than darkly pigmented individuals.
- Higher incidence in New Zealand and Australia where sun exposure in high.

Sites

Skin →
- Trunk (in men)
- Legs (in women)
- Other locations → face, soles, palms and nail-beds.

Other sites→ Oral and anogenital mucosal surfaces, esophagus, leptomeninges, eye, and the substantia nigra.

Gross

2 growth phases →
1. Initial radial growth
2. Later vertical growth phase

1) Radial growth phase

- Melanoma spread horizontally within epidermis and superficial dermis.

- Tumor cells lack capacity to metastasize.
- Clinicopathologic classes →

1. Lentigo maligna →

   - Indolent lesion on face of older men
   - Remain inradial growth phase for several decades.

2. Superficial spreading →

   - Most common type of melanoma
   - Involves skin exposed to sun

3. Acral/mucosal lentiginous melanoma→

   - Not related to sun exposure
   - Seen in palm, sole or mucosa.

2) Vertical growth phase



- After a variable and unpredictable period → melanoma from radial phase develops a vertical growth phase.
- During this phase→
    - Tumor cells invade downward into deeper dermis.
    - Melanoma may appear as a nodule
    - Metastatic potential.
    - Risk of metastasis correlates with the depth of invasion (Breslow thickness)



Microscopy

Tumors cells arrangement →
- Arranged in solid masses,sheets, islands, etc.
- Invades upper epidermis and deeper dermis

Tumor cells →
- Larger than normal melanocytes
- Large nuclei with irregular contours
- Clumping of chromatin at the periphery of nuclear membrane
- Prominent red nucleoli.
- Mitotic figures Common



Melanin pigment →
- May be present (melanotic) or absent (amelanotic melanoma) without any prognostic influence.
- If present, tends to be in form of uniform fine granules

Tumor infiltrating lymphocytes (TILS) →
- Seen in tissue surrounding the tumor

Depending upon the depth of invasion into dermis → Clark has described following 5 levels:
- Level I: Malignant melanoma cells confined to epidermis
- Level II: Extension into papillary dermis.
- Level III: Extension upto interface between papillary and reticular dermis.
- Level IV: Invasion of reticular dermis.
- Level V: Invasion of subcutaneous fat.



- Special stain for melanin → Masson–Fontana or DOPA-oxidase reaction.
- Tumor marker for Malignant melanoma → HMB 45

Clinical Features

- Usually asymptomatic
- May present with itching or pain at site of lesion.

Warning signs of melanoma → ABCD

- Asymmetry
- Irregular Borders
- Variegated **C**olor
- Increased **D**iameter

Prognosis

Depth of invasion is most important factor

Prognostic factors	Favourable prognosis
Tumor depth (the Breslow thickness)	Tumor depth of less than 1.7 mm
Number of mitoses	No or very few mitoses
Evidence of tumor regression (may be due to the host immune response)	Absence of regression
Presence and number of tumor infiltrating lymphocytes (TILs)	Brisk TIL response
Gender	Female gender
Location (central body or extremity)	Location on an extremity

	Squamous Cell Carcinoma	Basal Cell Carcinoma	Melanoma
Introduction			
Predisposing factors			
Pathogenesis			
Site			
Gross			
Microscopy			
Clinical features			
Prognosis			

Dr. Priyanka Sachdev is a distinguished medical professional with a background in both academia and clinical practice. She completed her MBBS from MGM Medical College, Indore, and further pursued MD from Tata Memorial Hospital, Mumbai, India. Dr. Sachdev's expertise spans a wide range of pathology topics, and she is renowned for her dedication to medical education. Through her extensive experience and commitment to excellence, Dr. Sachdev has become a trusted authority in the field, guiding and inspiring countless medical students on their academic journeys.Dr. Priyanka Sachdev's influential presence extends beyond traditional education channels. As the founder of the widely acclaimed YouTube channel and Instagram page, **MedLiveByDrPriyanka**, she has revolutionized medical education, reaching millions of students worldwide. Through engaging and informative content, Dr. Sachdev has empowered aspiring medical professionals with invaluable insights and practical knowledge, making complex medical concepts accessible to learners of all levels. Her dedication to supporting the global medical community has earned her widespread admiration and gratitude from students and professionals alike.

Salient features of the Book

- **ALIGNED WITH UNIVERSITY EXAM PATTERN:** Designed to mirror the structure of university exams paper I and II, providing comprehensive coverage of essential topics.
- **TWO-VOLUME SET:** A fully colored book presented in two volumes, offering a visually engaging and comprehensive resource for pathology studies.
- **CLEAR LINE DIAGRAMS:** Includes detailed line diagrams to aid understanding and facilitate easy reproduction of concepts during exams.
- **Effective Mnemonics and Flow Charts:** Utilizes mnemonics and flow charts to simplify complex topics, aiding in retention and understanding for students.
- **TOPIC-WISE QUESTION BANK:** Features topic-wise questions sourced from various university exams, providing valuable practice and insight into exam patterns.
- **EXTENSIVE COVERAGE:** Covers 26 units across more than 180 topics, ensuring thorough preparation for pathology exams and beyond.
- **SYNCED VIDEO COURSE AVAILABLE:** Access to a comprehensive video course by Dr. Priyanka Sachdev, fully synchronized with the content of the book. This video course, available to purchase separately, complements the book by offering in-depth explanations and visual demonstrations to enhance understanding and retention of key concepts.











ISBN 978-93-340-4816-2

@medlivebydrpriyanka